

# Frank W Lewis

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8701729/frank-w-lewis-publications-by-year.pdf>

**Version:** 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

37  
papers

1,286  
citations

16  
h-index

35  
g-index

41  
ext. papers

1,457  
ext. citations

4.6  
avg, IF

4.16  
L-index

#	Paper	IF	Citations
37	Design and evaluation of bi-functional iron chelators for protection of dopaminergic neurons from toxicants. <i>Archives of Toxicology</i> , <b>2020</b> , 94, 3105-3123	5.8	6
36	Novel 1-hydroxypyridin-2-one metal chelators prevent and rescue ubiquitin proteasomal-related neuronal injury in an in vitro model of Parkinson's disease. <i>Archives of Toxicology</i> , <b>2020</b> , 94, 813-831	5.8	4
35	Thermodynamic parameters of Am(III), Cm(III) and Eu(III) extraction by CyMe4-BTPhen in cyclohexanone from HNO <sub>3</sub> solutions. <i>Journal of Chemical Thermodynamics</i> , <b>2020</b> , 141, 105955	2.9	4
34	Exploring the Subtle Effect of Aliphatic Ring Size on Minor Actinide-Extraction Properties and Metal Ion Speciation in Bis-1,2,4-Triazine Ligands. <i>Chemistry - A European Journal</i> , <b>2020</b> , 26, 428-437	4.8	15
33	Separation of the Minor Actinides Americium(III) and Curium(III) by Hydrophobic and Hydrophilic BTPhen ligands: Exploiting Differences in their Rates of Extraction and Effective Separations at Equilibrium. <i>Solvent Extraction and Ion Exchange</i> , <b>2018</b> , 36, 115-135	2.5	14
32	1,2,4-Triazines in the Synthesis of Bipyridine Bisphenolate ONNO Ligands and Their Highly Luminescent Tetradentate Pt(II) Complexes for Solution-Processable OLEDs. <i>Inorganic Chemistry</i> , <b>2018</b> , 57, 3825-3832	5.1	23
31	Modelling of the Am(III) / Cm(III) kinetic separation effect observed during metal ion extraction by bis-(1,2,4)-triazine ligands. <i>Separation Science and Technology</i> , <b>2018</b> , 53, 277-285	2.5	8
30	Plutonium coordination and redox chemistry with the CyMe-BTPhen polydentate N-donor extractant ligand. <i>Chemical Communications</i> , <b>2018</b> , 54, 12582-12585	5.8	6
29	Spin-polarized electron transmission through chiral halocamphor molecules. <i>Journal of Physics B: Atomic, Molecular and Optical Physics</i> , <b>2018</b> , 51, 21LT01	1.3	4
28	An Overview of Multifunctional Metal Chelators as Potential Treatments for Neurodegenerative Diseases <b>2017</b> , 399-414		0
27	Anomalously Large Chiral Sensitivity in the Dissociative Electron Attachment of 10-Iodocamphor. <i>Physical Review Letters</i> , <b>2016</b> , 116, 093201	7.4	11
26	Solvent Extraction and Fluorescence Spectroscopic Investigation of the Selective Am(III) Complexation with TS-BTPhen. <i>Solvent Extraction and Ion Exchange</i> , <b>2016</b> , 34, 126-140	2.5	32
25	Rapid selective separation of americium/curium from simulated nuclear forensic matrices using triazine ligands. <i>Radiochimica Acta</i> , <b>2015</b> , 103, 687-694	1.9	7
24	Hydrophilic sulfonated bis-1,2,4-triazine ligands are highly effective reagents for separating actinides(III) from lanthanides(III) selective formation of aqueous actinide complexes. <i>Chemical Science</i> , <b>2015</b> , 6, 4812-4821	9.4	84
23	Protection from neurodegeneration in the 6-hydroxydopamine (6-OHDA) model of Parkinson's with novel 1-hydroxypyridin-2-one metal chelators. <i>Metallomics</i> , <b>2015</b> , 7, 867-76	4.5	33
22	Efficient masking of corrosion and fission products such as Ni(II) and Pd(II) in the presence of the minor actinide Am(III) using hydrophilic anionic or cationic bis-triazines. <i>Chemical Communications</i> , <b>2015</b> , 51, 9189-92	5.8	8
21	Role of ligands in catalytic water oxidation by mononuclear ruthenium complexes. <i>Coordination Chemistry Reviews</i> , <b>2015</b> , 304-305, 88-101	23.2	58

20	TS-BTPhen as a promising hydrophilic complexing agent for selective Am(III) separation by solvent extraction. <i>Nukleonika</i> , <b>2015</b> , 60, 815-820	1	11
19	Synthesis and Evaluation of a Novel Hydrophilic 6,6'-Bis(1,2,4-triazin-3-yl)-2,2'-bipyridine Ligand for Separating Actinide(III) from Lanthanide(III). <i>Synlett</i> , <b>2015</b> , 27, 1-5	2.2	3
18	An improved synthesis of 10-isobornylsultone. <i>Tetrahedron: Asymmetry</i> , <b>2014</b> , 25, 1150-1152		2
17	Use of soft heterocyclic N-donor ligands to separate actinides and lanthanides. <i>Inorganic Chemistry</i> , <b>2013</b> , 52, 3414-28	5.1	204
16	The Circuitous Journey from Malonamides to BTPhens. <i>Strategies and Tactics in Organic Synthesis</i> , <b>2013</b> , 177-202	0.2	6
15	Direct Selective Extraction of Actinides (III) from PUREX Raffinate using a Mixture of CyMe4BTBP and TODGA as 1-cycle SANEX Solvent Part III: Demonstration of a Laboratory-Scale Counter-Current Centrifugal Contactor Process. <i>Solvent Extraction and Ion Exchange</i> , <b>2013</b> , 31, 519-537	2.5	59
14	Lanthanide speciation in potential SANEX and GANEX actinide/lanthanide separations using tetra-N-donor extractants. <i>Inorganic Chemistry</i> , <b>2013</b> , 52, 3429-44	5.1	78
13	BTBPs versus BTPhens: some reasons for their differences in properties concerning the partitioning of minor actinides and the advantages of BTPhens. <i>Inorganic Chemistry</i> , <b>2013</b> , 52, 4993-5005	5.1	103
12	Synthesis and Evaluation of Lipophilic BTBP Ligands for An/Ln Separation in Nuclear Waste Treatment: The Effect of Alkyl Substitution on Extraction Properties and Implications for Ligand Design. <i>European Journal of Organic Chemistry</i> , <b>2012</b> , 2012, 1509-1519	3.2	40
11	The diastereoselective Meth-Cohn epoxidation of camphor-derived vinyl sulfones. <i>Tetrahedron: Asymmetry</i> , <b>2012</b> , 23, 643-649		1
10	Influence of diluent alkyl substitution on the extraction of Am(III) and Eu(III) by a 6,6'-bis(1,2,4-triazin-3-yl)-2,2'-bipyridine ligand dissolved in alkylated cyclohexanone diluents. <i>Radiochimica Acta</i> , <b>2012</b> , 100, 747-752	1.9	10
9	From BTBPs to BTPhens: The Effect of Ligand Pre-Organization on the Extraction Properties of Quadridentate Bis-Triazine Ligands. <i>Procedia Chemistry</i> , <b>2012</b> , 7, 231-238		37
8	Complexation of lanthanides, actinides and transition metal cations with a 6-(1,2,4-triazin-3-yl)-2,2'-bipyridine ligand: implications for actinide(III)/lanthanide(III) partitioning. <i>Dalton Transactions</i> , <b>2012</b> , 41, 9209-19	4.3	34
7	The Separation of Americium(III) from Europium(III) by Two New 6,6'-Bis(1,2,4-triazin-3-yl)-2,2'-bipyridines in Different Diluents. <i>Solvent Extraction and Ion Exchange</i> , <b>2011</b> , 29, 551-576	2.5	22
6	Highly efficient separation of actinides from lanthanides by a phenanthroline-derived bis-triazine ligand. <i>Journal of the American Chemical Society</i> , <b>2011</b> , 133, 13093-102	16.4	250
5	Preliminary investigations on novel camphor-derived chiral sulfones: completely stereoselective formation of tricyclic hydroxy sulfones from 8- and 10-functionalized camphor derivatives. <i>Tetrahedron</i> , <b>2011</b> , 67, 7517-7528	2.4	11
4	Development of Highly Selective Ligands for Separations of Actinides from Lanthanides in the Nuclear Fuel Cycle. <i>Synlett</i> , <b>2011</b> , 2011, 2609-2632	2.2	51
3	Interaction of 6,6'-bis(5,5,8,8-tetramethyl-5,6,7,8-tetrahydro-1,2,4-benzotriazin-3-yl)-2,2'-bipyridine (CyMe4-BTTP) with some trivalent ions such as lanthanide(III) ions and americium(III). <i>Dalton Transactions</i> , <b>2010</b> , 2010, 5172-82	4.3	31

- 2 Synthesis of  $\beta$ -Amino Alcohols via the Reduction of Lactamides Derived from Ethyl (2S)-Lactate with Borane-Methyl Sulfide. *Synlett*, **2009**, 2009, 1923-1928 2.2 2
- 1 Unexpected formation of 10-iodo- and 10-chlorocamphor under halosulfonylation conditions, and convenient routes to 10-chloro- and 10-bromocamphor. *Tetrahedron: Asymmetry*, **2009**, 20, 1531-1535 14