

# Hiroaki Taguchi

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8705007/publications.pdf>

Version: 2024-02-01

14  
papers

339  
citations

1163117

8  
h-index

1125743

13  
g-index

14  
all docs

14  
docs citations

14  
times ranked

281  
citing authors

#	ARTICLE	IF	CITATIONS
1	Naturally Occurring Proteolytic Antibodies. <i>Journal of Biological Chemistry</i> , 2004, 279, 39611-39619.	3.4	77
2	Autoantibody-catalyzed Hydrolysis of Amyloid $\beta$ Peptide. <i>Journal of Biological Chemistry</i> , 2008, 283, 4714-4722.	3.4	70
3	Phosphonate Ester Probes for Proteolytic Antibodies. <i>Journal of Biological Chemistry</i> , 2001, 276, 28314-28320.	3.4	60
4	Overview and outlook of Toll-like receptor ligand-antigen conjugate vaccines. <i>Therapeutic Delivery</i> , 2012, 3, 749-760.	2.2	45
5	Covalent reactivity of phosphonate monophenyl esters with serine proteinases: an overlooked feature of presumed transition state analogs. <i>Archives of Biochemistry and Biophysics</i> , 2002, 402, 281-288.	3.0	25
6	A mechanism-based probe for gp120-Hydrolyzing antibodies. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2002, 12, 3167-3170.	2.2	14
7	6-(4-Amino-2-butyl-imidazoquinolyl)-norleucine: Toll-like receptor 7 and 8 agonist amino acid for self-adjuvanting peptide vaccine. <i>Amino Acids</i> , 2016, 48, 1319-1329.	2.7	12
8	Role of the constant region domain in the structural diversity of human antibody light chains. <i>FASEB Journal</i> , 2017, 31, 1668-1677.	0.5	9
9	Effects of Physical Damage in the Intermediate Phase on the Progression of Amyloid $\beta$ Fibrillization. <i>Chemistry - an Asian Journal</i> , 2019, 14, 4140-4145.	3.3	9
10	Finding and characterizing a catalytic antibody light chain, H34, capable of degrading the PD-1 molecule. <i>RSC Chemical Biology</i> , 2021, 2, 220-229.	4.1	7
11	A new algorithm to convert a normal antibody into the corresponding catalytic antibody. <i>Science Advances</i> , 2020, 6, eaay6441.	10.3	6
12	Identification of the minimal region of peptide derived from ADP-ribosylation factor1 (ARF1) that inhibits IgE-mediated mast cell activation. <i>Molecular Immunology</i> , 2019, 105, 32-37.	2.2	4
13	Development of an activity-based probe for amyloid $\beta$ -hydrolyzing antibodies. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2016, 26, 2210-2213.	2.2	1
14	A unique method for antibody to possess the catalytic function (3 rd report). <i>FASEB Journal</i> , 2018, 32, lb77.	0.5	0