
## Felix Goeser

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8708502/publications.pdf Version: 2024-02-01



FELLY COESED

#	Article	IF	CITATIONS
1	Compartment-specific distribution of human intestinal innate lymphoid cells is altered in HIV patients under effective therapy. PLoS Pathogens, 2017, 13, e1006373.	2.1	85
2	Polo-Like Kinase 1 is a Therapeutic Target in High-Risk Neuroblastoma. Clinical Cancer Research, 2011, 17, 731-741.	3.2	67
3	Antibiotic resistance in healthcareâ€related and nosocomial spontaneous bacterial peritonitis. European Journal of Clinical Investigation, 2017, 47, 44-52.	1.7	50
4	The ratio of calprotectin to total protein as a diagnostic and prognostic marker for spontaneous bacterial peritonitis in patients with liver cirrhosis and ascites. Clinical Chemistry and Laboratory Medicine, 2015, 53, 2031-9.	1.4	35
5	Fecal Microbiota Transplant in Patients With Re current Clostridium Difficile Infection. Deutsches Ärzteblatt International, 2016, 113, 583-9.	0.6	35
6	The impact of technical and clinical factors on fecal microbiota transfer outcomes for the treatment of recurrent <i>Clostridioides difficile</i> infections in Germany. United European Gastroenterology Journal, 2019, 7, 716-722.	1.6	24
7	Fecal microbiota transfer for refractory intestinal graftâ€versusâ€host disease — Experience from two German tertiary centers. European Journal of Haematology, 2021, 107, 229-245.	1.1	20
8	Relative Ascites Polymorphonuclear Cell Count Indicates Bacterascites and Risk of Spontaneous Bacterial Peritonitis. Digestive Diseases and Sciences, 2017, 62, 2558-2568.	1.1	16
9	Alterations of the NK cell pool in HIV/HCV co-infection. PLoS ONE, 2017, 12, e0174465.	1.1	13
10	Hypoxia impairs anti-viral activity of natural killer (NK) cells but has little effect on anti-fibrotic NK cell functions in hepatitis C virus infection. Journal of Hepatology, 2015, 63, 1334-1344.	1.8	11
11	HIV mono-infection is associated with an impaired anti-HCV activity of NK cells. Aids, 2015, 30, 1.	1.0	10
12	A genetic variant in tollâ€like receptor 5 is linked to chemokine levels and hepatocellular carcinoma in steatohepatitis. Liver International, 2021, 41, 2139-2148.	1.9	6
13	IL-28B Genetic Variants Determine the Extent of Monocyte-Induced Activation of NK Cells in Hepatitis C. PLoS ONE, 2016, 11, e0162068.	1.1	4
14	Neither black nor white: do altered intestinal microbiota reflect chronic liver disease severity?. Gut, 2021, 70, gutjnl-2020-321424.	6.1	4
15	8. Mikrobiom und Lebererkrankungen. , 2016, , .		2
16	Clinical effectiveness of bidirectional fecal microbiota transfer in the treatment of recurrent Clostridioides difficile infections. Digestive and Liver Disease, 2021, 53, 706-711.	0.4	1
17	Research update for articles published in EJCI in 2017. European Journal of Clinical Investigation, 2019, 49, e13163.	1.7	0
18	Renal replacement therapy as bridging-to-recovery in refractory hepatorenal syndrome. Zeitschrift Fur Gastroenterologie, 2021, 59, 331-335.	0.2	0