

# Xin-Ming Chen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8718755/publications.pdf>

Version: 2024-02-01

38  
papers

1,451  
citations

304743

22  
h-index

315739

38  
g-index

39  
all docs

39  
docs citations

39  
times ranked

2005  
citing authors

#	ARTICLE	IF	CITATIONS
1	Thioredoxin interacting protein (TXNIP) regulates tubular autophagy and mitophagy in diabetic nephropathy through the mTOR signaling pathway. <i>Scientific Reports</i> , 2016, 6, 29196.	3.3	106
2	The renal cortical fibroblast in renal tubulointerstitial fibrosis. <i>International Journal of Biochemistry and Cell Biology</i> , 2006, 38, 1-5.	2.8	100
3	Transforming growth factor- $\beta$ 2/connective tissue growth factor axis in the kidney. <i>International Journal of Biochemistry and Cell Biology</i> , 2008, 40, 9-13.	2.8	94
4	TGF- $\beta$ 1 induces IL-8 and MCP-1 through a connective tissue growth factor-independent pathway. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 290, F703-F709.	2.7	84
5	Blockade of KCa3.1 Ameliorates Renal Fibrosis Through the TGF- $\beta$ 1/Smad Pathway in Diabetic Mice. <i>Diabetes</i> , 2013, 62, 2923-2934.	0.6	77
6	Role of Kr $\beta$ 4ppel-like factor 6 in transforming growth factor- $\beta$ 1-induced epithelial-mesenchymal transition of proximal tubule cells. <i>American Journal of Physiology - Renal Physiology</i> , 2008, 295, F1388-F1396.	2.7	76
7	MiRNA-200b represses transforming growth factor- $\beta$ 1-induced EMT and fibronectin expression in kidney proximal tubular cells. <i>American Journal of Physiology - Renal Physiology</i> , 2013, 304, F1266-F1273.	2.7	74
8	High Glucose-Induced Thioredoxin-Interacting Protein in Renal Proximal Tubule Cells Is Independent of Transforming Growth Factor- $\beta$ 1. <i>American Journal of Pathology</i> , 2007, 171, 744-754.	3.8	71
9	KCa3.1 mediates dysfunction of tubular autophagy in diabetic kidneys via PI3k/Akt/mTOR signaling pathways. <i>Scientific Reports</i> , 2016, 6, 23884.	3.3	60
10	Thioredoxin-interacting protein mediates dysfunction of tubular autophagy in diabetic kidneys through inhibiting autophagic flux. <i>Laboratory Investigation</i> , 2014, 94, 309-320.	3.7	50
11	Transcription Factors Kr $\beta$ 4ppel-Like Factor 6 and Peroxisome Proliferator-Activated Receptor- $\beta$ 3 Mediate High Glucose-Induced Thioredoxin-Interacting Protein. <i>American Journal of Pathology</i> , 2009, 175, 1858-1867.	3.8	48
12	Increased sphingosine 1-phosphate mediates inflammation and fibrosis in tubular injury in diabetic nephropathy. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2016, 43, 56-66.	1.9	48
13	Transforming growth factor- $\beta$ 1 differentially mediates fibronectin and inflammatory cytokine expression in kidney tubular cells. <i>American Journal of Physiology - Renal Physiology</i> , 2006, 291, F1070-F1077.	2.7	46
14	KCa3.1 mediates activation of fibroblasts in diabetic renal interstitial fibrosis. <i>Nephrology Dialysis Transplantation</i> , 2014, 29, 313-324.	0.7	44
15	The role of Sgk-1 in the upregulation of transport proteins by PPAR- $\alpha$ agonists in human proximal tubule cells. <i>Nephrology Dialysis Transplantation</i> , 2008, 24, 1130-1141.	0.7	40
16	Mesenchymal Stem Cell-Derived Extracellular Vesicles to the Rescue of Renal Injury. <i>International Journal of Molecular Sciences</i> , 2021, 22, 6596.	4.1	37
17	High glucose induces macrophage inflammatory protein-3 $\alpha$ in renal proximal tubule cells via a transforming growth factor- $\beta$ 1 dependent mechanism. <i>Nephrology Dialysis Transplantation</i> , 2007, 22, 3147-3153.	0.7	34
18	Inhibition of KCa3.1 suppresses TGF- $\beta$ 1 induced MCP-1 expression in human proximal tubular cells through Smad3, p38 and ERK1/2 signaling pathways. <i>International Journal of Biochemistry and Cell Biology</i> , 2014, 47, 1-10.	2.8	27

#	ARTICLE	IF	CITATIONS
19	The roles of Kruppel-like factor 6 and peroxisome proliferator-activated receptor- $\beta$ in the regulation of macrophage inflammatory protein-3 $\alpha$ at early onset of diabetes. <i>International Journal of Biochemistry and Cell Biology</i> , 2011, 43, 383-392.	2.8	26
20	Preparation of Inert Polystyrene Latex Particles as MicroRNA Delivery Vectors by Surfactant-Free RAFT Emulsion Polymerization. <i>Biomacromolecules</i> , 2016, 17, 965-973.	5.4	26
21	Fluorescent Labeling and Biodistribution of Latex Nanoparticles Formed by Surfactant-Free RAFT Emulsion Polymerization. <i>Macromolecular Bioscience</i> , 2017, 17, 1600366.	4.1	26
22	MicroRNA as novel biomarkers and therapeutic targets in diabetic kidney disease: An update. <i>FASEB BioAdvances</i> , 2019, 1, 375-388.	2.4	25
23	RIPK3 blockade attenuates tubulointerstitial fibrosis in a mouse model of diabetic nephropathy. <i>Scientific Reports</i> , 2020, 10, 10458.	3.3	24
24	Metformin attenuates folic acid induced renal fibrosis in mice. <i>Journal of Cellular Physiology</i> , 2018, 233, 7045-7054.	4.1	23
25	The role of Kruppel-like factor 4 in transforming growth factor- $\beta$ -induced inflammatory and fibrotic responses in human proximal tubule cells. <i>Clinical and Experimental Pharmacology and Physiology</i> , 2015, 42, 680-686.	1.9	21
26	RIPK3 blockade attenuates kidney fibrosis in a folic acid model of renal injury. <i>FASEB Journal</i> , 2020, 34, 10286-10298.	0.5	20
27	Metformin Attenuates Renal Fibrosis in a Mouse Model of Adenine-Induced Renal Injury Through Inhibiting TGF- $\beta$ 1 Signaling Pathways. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 603802.	3.7	19
28	High Glucose Induces CCL20 in Proximal Tubular Cells via Activation of the KCa3.1 Channel. <i>PLoS ONE</i> , 2014, 9, e95173.	2.5	17
29	The differential regulation of Smad7 in kidney tubule cells by connective tissue growth factor and transforming growth factor-beta1. <i>Nephrology</i> , 2007, 12, 267-274.	1.6	16
30	Role of the potassium channel KCa3.1 in diabetic nephropathy. <i>Clinical Science</i> , 2014, 127, 423-433.	4.3	15
31	The KCa3.1 blocker TRAM34 reverses renal damage in a mouse model of established diabetic nephropathy. <i>PLoS ONE</i> , 2018, 13, e0192800.	2.5	15
32	Mesenchymal Stem Cell-Derived Exosomes: Toward Cell-Free Therapeutic Strategies in Chronic Kidney Disease. <i>Frontiers in Medicine</i> , 2022, 9, 816656.	2.6	14
33	RIPK3: A New Player in Renal Fibrosis. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 502.	3.7	12
34	KCa3.1 Mediates Dysregulation of Mitochondrial Quality Control in Diabetic Kidney Disease. <i>Frontiers in Cell and Developmental Biology</i> , 2021, 9, 573814.	3.7	10
35	KCa3.1. <i>Current Opinion in Nephrology and Hypertension</i> , 2015, 24, 61-66.	2.0	9
36	The Mitochondrial Kinase PINK1 in Diabetic Kidney Disease. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1525.	4.1	9

#	ARTICLE	IF	CITATIONS
37	A single-domain i-body, AD-114, attenuates renal fibrosis through blockade of CXCR4. JCI Insight, 2022, 7, .	5.0	5
38	KCa3.1 in diabetic kidney disease. Current Opinion in Nephrology and Hypertension, 2022, 31, 129-134.	2.0	3