

Stefania Grimaudo

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8724820/stefania-grimaudo-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

99
papers

3,864
citations

35
h-index

60
g-index

114
ext. papers

4,461
ext. citations

5
avg, IF

4.58
L-index

#	Paper	IF	Citations
99	PSD3 downregulation confers protection against fatty liver disease.. <i>Nature Metabolism</i> , 2022 , 4, 60-75	14.6	1
98	PCSK9 rs11591147 R46L loss-of-function variant protects against liver damage in individuals with NAFLD. <i>Liver International</i> , 2021 , 41, 321-332	7.9	10
97	NR1H4 rs35724 G>C variant modulates liver damage in nonalcoholic fatty liver disease. <i>Liver International</i> , 2021 , 41, 2712-2719	7.9	3
96	Coffee Restores Expression of lncRNAs Involved in Steatosis and Fibrosis in a Mouse Model of NAFLD. <i>Nutrients</i> , 2021 , 13,	6.7	3
95	PNPLA3 rs738409 C>G Variant Predicts Fibrosis Progression by Noninvasive Tools in Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 1979-1981	6.9	4
94	The "Janus" Role of C/EBPs Family Members in Cancer Progression. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	10
93	Aminopyrine breath test predicts liver-related events and death in HCV-related cirrhosis on SVR after DAA therapy. <i>Liver International</i> , 2020 , 40, 530-538	7.9	1
92	Neurotensin up-regulation is associated with advanced fibrosis and hepatocellular carcinoma in patients with MAFLD. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2020 , 1865, 158765	5	4
91	Hepatitis C virus eradication by direct antiviral agents abates oxidative stress in patients with advanced liver fibrosis. <i>Liver International</i> , 2020 , 40, 2820-2827	7.9	9
90	Optimizing Sequential Systemic Therapies for Advanced Hepatocellular Carcinoma: A Decision Analysis. <i>Cancers</i> , 2020 , 12,	6.6	10
89	SARS-CoV-2 Viral Load, IFN γ Polymorphisms and the Course of COVID-19: An Observational Study. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	21
88	Association Between PNPLA3 rs738409 C>G Variant and Liver-Related Outcomes in Patients With Nonalcoholic Fatty Liver Disease. <i>Clinical Gastroenterology and Hepatology</i> , 2020 , 18, 935-944.e3	6.9	55
87	Serum coding and non-coding RNAs as biomarkers of NAFLD and fibrosis severity. <i>Liver International</i> , 2019 , 39, 1742-1754	7.9	37
86	Role of Myeloid-Epithelial-Reproductive Tyrosine Kinase and Macrophage Polarization in the Progression of Atherosclerotic Lesions Associated With Nonalcoholic Fatty Liver Disease. <i>Frontiers in Pharmacology</i> , 2019 , 10, 604	5.6	10
85	FXR rs35724 G>C variant modulates cholesterol levels, carotid atherosclerosis and liver damage in non-alcoholic fatty liver. <i>Digestive and Liver Disease</i> , 2019 , 51, e26	3.3	3
84	NS5A Gene Analysis by Next Generation Sequencing in HCV Nosocomial Transmission Clusters of HCV Genotype 1b Infected Patients. <i>Cells</i> , 2019 , 8,	7.9	9
83	STAT5 and STAT5 Inhibitors in Hematological Malignancies. <i>Anti-Cancer Agents in Medicinal Chemistry</i> , 2019 , 19, 2036-2046	2.2	1

82	Hepatitis C virus eradication by direct-acting antiviral agents improves carotid atherosclerosis in patients with severe liver fibrosis. <i>Journal of Hepatology</i> , 2018 , 69, 18-24	13.4	68
81	Prevalence and severity of nonalcoholic fatty liver disease by transient elastography: Genetic and metabolic risk factors in a general population. <i>Liver International</i> , 2018 , 38, 2060-2068	7.9	39
80	Causal relationship of hepatic fat with liver damage and insulin resistance in nonalcoholic fatty liver. <i>Journal of Internal Medicine</i> , 2018 , 283, 356-370	10.8	140
79	Protein phosphatase 1 regulatory subunit 3B gene variation protects against hepatic fat accumulation and fibrosis in individuals at high risk of nonalcoholic fatty liver disease. <i>Hepatology Communications</i> , 2018 , 2, 666-675	6	30
78	Fibronectin Type III Domain-Containing Protein 5 rs3480 A>G Polymorphism, Irisin, and Liver Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017 , 102, 2660-2669	5.6	30
77	Telomerase reverse transcriptase germline mutations and hepatocellular carcinoma in patients with nonalcoholic fatty liver disease. <i>Cancer Medicine</i> , 2017 , 6, 1930-1940	4.8	29
76	Interferon lambda 4 rs368234815 TT>G variant is associated with liver damage in patients with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2017 , 66, 1885-1893	11.2	59
75	MBOAT7 rs641738 variant and hepatocellular carcinoma in non-cirrhotic individuals. <i>Scientific Reports</i> , 2017 , 7, 4492	4.9	131
74	Effects of Pimozide Derivatives on pSTAT5 in K562 Cells. <i>ChemMedChem</i> , 2017 , 12, 1183-1190	3.7	16
73	MBOAT7 locus rs641738 variant predisposes to hepatocellular carcinoma in nonalcoholic fatty liver. <i>Digestive and Liver Disease</i> , 2016 , 48, e7-e8	3.3	2
72	MERTK rs4374383 polymorphism affects the severity of fibrosis in non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2016 , 64, 682-90	13.4	79
71	Novel iodoacetamido benzoheterocyclic derivatives with potent antileukemic activity are inhibitors of STAT5 phosphorylation. <i>European Journal of Medicinal Chemistry</i> , 2016 , 108, 39-52	6.8	6
70	Clinical Course and Genetic Susceptibility of Primary Biliary Cirrhosis: Analysis of a Prospective Cohort. <i>Hepatitis Monthly</i> , 2016 , 16, e31681	1.8	3
69	TM6SF2 rs58542926 is not associated with steatosis and fibrosis in large cohort of patients with genotype 1 chronic hepatitis C. <i>Liver International</i> , 2016 , 36, 198-204	7.9	12
68	Effects of Eradicating Hepatitis C Virus Infection in Patients With Cirrhosis Differ With Stage of Portal Hypertension. <i>Gastroenterology</i> , 2016 , 151, 130-139.e2	13.3	107
67	Transmembrane 6 superfamily member 2 gene variant disentangles nonalcoholic steatohepatitis from cardiovascular disease. <i>Hepatology</i> , 2015 , 61, 506-14	11.2	311
66	The hepatic expression of vitamin D receptor is inversely associated with the severity of liver damage in genotype 1 chronic hepatitis C patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015 , 100, 193-200	5.6	21
65	The UCP2 -866G>A promoter region polymorphism is associated with nonalcoholic steatohepatitis. <i>Liver International</i> , 2015 , 35, 1574-80	7.9	35

64	PNPLA3 rs738409 I748M is associated with steatohepatitis in 434 non-obese subjects with hepatitis C. <i>Alimentary Pharmacology and Therapeutics</i> , 2015 , 41, 939-48	6.1	16
63	Synthesis and antiproliferative activity of 3-(2-chloroethyl)-5-methyl-6-phenyl-8-(trifluoromethyl)-5,6-dihydropyrazolo[3,4-f][1,2,3,5]tetrazepin-4-(3H)-one	3.8	9
	<i>European Journal of Medicinal Chemistry</i> , 2015 , 96, 98-104		
62	Statin use and non-alcoholic steatohepatitis in at risk individuals. <i>Journal of Hepatology</i> , 2015 , 63, 705-12	3.4	227
61	Inhibition of activated STAT5 in Bcr/Abl expressing leukemia cells with new pimozide derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2014 , 24, 4568-4574	2.9	21
60	Novel antiproliferative chimeric compounds with marked histone deacetylase inhibitory activity. <i>ACS Medicinal Chemistry Letters</i> , 2014 , 5, 973-8	4.3	10
59	Glucokinase regulatory protein gene polymorphism affects liver fibrosis in non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2014 , 9, e87523	3.7	83
58	Effects of IL28B rs12979860 CC genotype on metabolic profile and sustained virologic response in patients with genotype 1 chronic hepatitis C. <i>Clinical Gastroenterology and Hepatology</i> , 2013 , 11, 311-7.e19	6.9	27
57	Role of IL-28B and inosine triphosphatase polymorphisms in efficacy and safety of Peg-Interferon and ribavirin in chronic hepatitis C compensated cirrhosis with and without oesophageal varices. <i>Journal of Viral Hepatitis</i> , 2013 , 20, 113-21	3.4	16
56	Association of vitamin D serum levels and its common genetic determinants, with severity of liver fibrosis in genotype 1 chronic hepatitis C patients. <i>Journal of Viral Hepatitis</i> , 2013 , 20, 486-93	3.4	42
55	Insulin-like growth factor-I, inflammatory proteins, and fibrosis in subjects with nonalcoholic fatty liver disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, E304-8	5.6	48
54	The new iodoacetamidobenzofuran derivative TR120 decreases STAT5 expression and induces antitumor effects in imatinib-sensitive and imatinib-resistant BCR-ABL-expressing leukemia cells. <i>Anti-Cancer Drugs</i> , 2013 , 24, 384-93	2.4	6
53	PNPLA3 GG genotype and carotid atherosclerosis in patients with non-alcoholic fatty liver disease. <i>PLoS ONE</i> , 2013 , 8, e74089	3.7	47
52	A natural-like synthetic small molecule impairs bcr-abl signaling cascades and induces megakaryocyte differentiation in erythroleukemia cells. <i>PLoS ONE</i> , 2013 , 8, e57650	3.7	11
51	IL28B polymorphisms influence stage of fibrosis and spontaneous or interferon-induced viral clearance in thalassemia patients with hepatitis C virus infection. <i>Haematologica</i> , 2012 , 97, 679-86	6.6	41
50	IL28B and PNPLA3 polymorphisms affect histological liver damage in patients with non-alcoholic fatty liver disease. <i>Journal of Hepatology</i> , 2012 , 56, 1356-62	13.4	72
49	Vitamin D levels and IL28B polymorphisms are related to rapid virological response to standard of care in genotype 1 chronic hepatitis C. <i>Antiviral Therapy</i> , 2012 , 17, 823-31	1.6	28
48	High liver RBP4 protein content is associated with histological features in patients with genotype 1 chronic hepatitis C and with nonalcoholic steatohepatitis. <i>Digestive and Liver Disease</i> , 2011 , 43, 404-10	3.3	24
47	Genetic predisposition to thrombophilia in inflammatory bowel disease. <i>Journal of Clinical Gastroenterology</i> , 2011 , 45, e25-9	3	4

46	3-Aryl-2-[1H-benzotriazol-1-yl]acrylonitriles: a novel class of potent tubulin inhibitors. <i>European Journal of Medicinal Chemistry</i> , 2011 , 46, 4151-67	6.8	31
45	Synthesis of substituted 3-amino-N-phenyl-1H-indazole-1-carboxamides endowed with antiproliferative activity. <i>European Journal of Medicinal Chemistry</i> , 2011 , 46, 168-74	6.8	14
44	Synthesis of novel antimitotic agents based on 2-amino-3-aryl-5-(hetero)arylethynyl thiophene derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2011 , 21, 2746-51	2.9	26
43	Synthesis, antiproliferative activity, and mechanism of action of a series of 2-[[2E)-3-phenylprop-2-enoyl]amino}benzamides. <i>European Journal of Medicinal Chemistry</i> , 2011 , 46, 2786-96	6.8	25
42	Substituted 2-(3,4,5-trimethoxybenzoyl)-benzo[b]thiophene derivatives as potent tubulin polymerization inhibitors. <i>Bioorganic and Medicinal Chemistry</i> , 2010 , 18, 5114-22	3.4	35
41	Design, synthesis and structure-activity relationship of 2-(3,4,5-trimethoxybenzoyl)-benzo[b]furan derivatives as a novel class of inhibitors of tubulin polymerization. <i>Bioorganic and Medicinal Chemistry</i> , 2009 , 17, 6862-71	3.4	64
40	Synthesis and antiproliferative activity of 3-amino-N-phenyl-1H-indazole-1-carboxamides. <i>European Journal of Medicinal Chemistry</i> , 2009 , 44, 165-78	6.8	18
39	Identification of biphenyl-based hybrid molecules able to decrease the intracellular level of Bcl-2 protein in Bcl-2 overexpressing leukemia cells. <i>Journal of Medicinal Chemistry</i> , 2009 , 52, 6936-40	8.3	68
38	Synthesis and biological evaluation of 1-methyl-2-(3,4,5-trimethoxybenzoyl)-3-aminoindoles as a new class of antimitotic agents and tubulin inhibitors. <i>Journal of Medicinal Chemistry</i> , 2008 , 51, 1464-8	8.3	78
37	Superficial venous thrombosis: prevalence of common genetic risk factors and their role on spreading to deep veins. <i>Thrombosis Research</i> , 2008 , 123, 194-9	8.2	41
36	Novel terphenyls and 3,5-diaryl isoxazole derivatives endowed with growth supporting and antiapoptotic properties. <i>Journal of Medicinal Chemistry</i> , 2008 , 51, 4796-803	8.3	31
35	Galangin increases the cytotoxic activity of imatinib mesylate in imatinib-sensitive and imatinib-resistant Bcr-Abl expressing leukemia cells. <i>Cancer Letters</i> , 2008 , 265, 289-97	9.9	48
34	Antiproliferative agents that interfere with the cell cycle at the G1→S transition: further development and characterization of a small library of stilbene-derived compounds. <i>ChemMedChem</i> , 2008 , 3, 345-55	3.7	8
33	Design, synthesis, and biological evaluation of thiophene analogues of chalcones. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 5367-76	3.4	82
32	Synthesis and biological evaluation of 2-(3,4,5-trimethoxybenzoyl)-3-N,N-dimethylamino benzo[b]furan derivatives as inhibitors of tubulin polymerization. <i>Bioorganic and Medicinal Chemistry</i> , 2008 , 16, 8419-26	3.4	34
31	Synthesis and biological evaluation of 2-amino-3-(3,4,5-trimethoxybenzoyl)-6-substituted-4,5,6,7-tetrahydrothieno[2,3-c]pyridine derivatives as antimitotic agents and inhibitors of tubulin polymerization. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 5041-5	2.9	20
30	Synthesis and biological evaluation of 2- and 3-aminobenzo[b]thiophene derivatives as antimitotic agents and inhibitors of tubulin polymerization. <i>Journal of Medicinal Chemistry</i> , 2007 , 50, 2273-7	8.3	115
29	Identification of a terphenyl derivative that blocks the cell cycle in the G0-G1 phase and induces differentiation in leukemia cells. <i>Journal of Medicinal Chemistry</i> , 2006 , 49, 3012-8	8.3	71

28	Synthesis and biological evaluation of 2-(3,4,5-trimethoxybenzoyl)-3-amino 5-aryl thiophenes as a new class of tubulin inhibitors. <i>Journal of Medicinal Chemistry</i> , 2006 , 49, 6425-8	8.3	50
27	Stilbene-based anticancer agents: resveratrol analogues active toward HL60 leukemic cells with a non-specific phase mechanism. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 3245-8	2.9	60
26	Methylenetetrahydrofolate reductase mutation in subjects with abdominal aortic aneurysm subdivided for age. <i>Clinical Hemorheology and Microcirculation</i> , 2006 , 34, 421-6	2.5	17
25	Heterocyclic and phenyl double-bond-locked combretastatin analogues possessing potent apoptosis-inducing activity in HL60 and in MDR cell lines. <i>Journal of Medicinal Chemistry</i> , 2005 , 48, 723-36	8.3	130
24	Synthesis and pharmacology of 6-substituted benzotropines: discovery of novel dopamine uptake inhibitors possessing low binding affinity to the dopamine transporter. <i>Journal of Medicinal Chemistry</i> , 2005 , 48, 3337-43	8.3	8
23	Pterostilbene and 3-hydroxypterostilbene are effective apoptosis-inducing agents in MDR and BCR-ABL-expressing leukemia cells. <i>International Journal of Biochemistry and Cell Biology</i> , 2005 , 37, 1709-26	5.6	127
22	Prolonged prothrombin time, Factor VII and activated FVII levels in chronic liver disease are partly dependent on Factor VII gene polymorphisms. <i>Digestive and Liver Disease</i> , 2005 , 37, 446-50	3.3	2
21	Studies on the apoptotic activity of natural and synthetic retinoids: discovery of a new class of synthetic terphenyls that potently support cell growth and inhibit apoptosis in neuronal and HL-60 cells. <i>Journal of Medicinal Chemistry</i> , 2005 , 48, 4293-9	8.3	26
20	Novel Stilbene-Based Antileukemic Agents Active in P-Glycoprotein Expressing and Apoptosis-Resistant Acute Leukaemia Cell Lines.. <i>Blood</i> , 2005 , 106, 4436-4436	2.2	
19	Resistance to gemcitabine in a lymphoma cell line resistant to Fas-mediated apoptosis. <i>Anticancer Research</i> , 2004 , 24, 851-7	2.3	4
18	Synthesis and biological evaluation of resveratrol and analogues as apoptosis-inducing agents. <i>Journal of Medicinal Chemistry</i> , 2003 , 46, 3546-54	8.3	190
17	A convenient synthesis of unsymmetrically substituted terphenyls of biologically active stilbenes via a double Suzuki cross-coupling protocol. <i>Tetrahedron Letters</i> , 2003 , 44, 3005-3008	2	29
16	Chemically modified tetracyclines induce cytotoxic effects against J774 tumour cell line by activating the apoptotic pathway. <i>International Immunopharmacology</i> , 2003 , 3, 63-73	5.8	23
15	Retinoic acid and analogs as potent inducers of differentiation and apoptosis. New promising chemopreventive and chemotherapeutic agents in oncology. <i>Pure and Applied Chemistry</i> , 2001 , 73, 1437-1444	2.1	6
14	Effects of chemically modified tetracyclines (CMTs) in sensitive, multidrug resistant and apoptosis resistant leukaemia cell lines. <i>British Journal of Pharmacology</i> , 2001 , 133, 306-14	8.6	20
13	Heterocycle-containing retinoids. Discovery of a novel isoxazole retinoid possessing potent apoptotic activity in multidrug and drug-induced apoptosis-resistant cells. <i>Journal of Medicinal Chemistry</i> , 2001 , 44, 2308-18	8.3	76
12	Anti-inflammatory effects of chemically modified tetracyclines by the inhibition of nitric oxide and interleukin-12 synthesis in J774 cell line. <i>International Immunopharmacology</i> , 2001 , 1, 1765-76	5.8	50
11	Pig liver esterase (PLE)-mediated resolution of N-substituted 4-benzoyloxy-3-carbomethoxypiperidines: a convenient preparation of 4-hydroxy- and 4-benzoyloxy-3-carbomethoxypiperidines in enantiomerically pure form. <i>Tetrahedron: Asymmetry</i> , 2000 , 11, 1897-1905		6

10	Programmed cell death (PCD) associated with the stilbene motif of arotinoids: discovery of novel apoptosis inducer agents possessing activity on multidrug resistant tumor cells. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2000 , 10, 2669-73	2.9	12
9	Structure-activity relationship studies of novel heteroretinoids: induction of apoptosis in the HL-60 cell line by a novel isoxazole-containing heteroretinoid. <i>Journal of Medicinal Chemistry</i> , 1999 , 42, 4961-9	8.3	39
8	The CD95/CD95 ligand system is not the major effector in anticancer drug-mediated apoptosis. <i>Cell Death and Differentiation</i> , 1998 , 5, 735-42	12.7	46
7	Reactivity of aminopyrroles: Protonation. <i>Journal of Heterocyclic Chemistry</i> , 1996 , 33, 161-168	1.9	10
6	Reaction of amino substituted heterocycles with one heteroatom in a five-membered ring as enamines. A revision. <i>Journal of Heterocyclic Chemistry</i> , 1995 , 32, 985-989	1.9	16
5	Organometallic complexes with biological molecules, IV. Di- and tri-organotin(IV) amoxicillin derivatives: Solid-state and solution-phase spectroscopic investigations. <i>Applied Organometallic Chemistry</i> , 1995 , 9, 227-239	3.1	25
4	3-Triazenoindoles. Synthesis and antileukemic activity. <i>European Journal of Medicinal Chemistry</i> , 1994 , 29, 889-891	6.8	11
3	Polycondensed Nitrogen Heterocycles. Part 26. Aminopyrrolo[1,2-f]phenanthridines by Decomposition and Cyclization of 2-Aryl-1-(3-azidophenyl)pyrroles. <i>Heterocycles</i> , 1994 , 37, 1549	0.8	6
2	Synthesis of 3-Triazenopyrroles. <i>Synthetic Communications</i> , 1993 , 23, 1627-1631	1.7	5
1	Identification and cloning of a umu locus in <i>Streptomyces coelicolor</i> A3(2). <i>Mutation Research-Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1991 , 262, 183-8		9