

# Jianming Li

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8727518/jianming-li-publications-by-year.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

68

papers

9,509

citations

40

h-index

74

g-index

74

ext. papers

11,183

ext. citations

11.9

avg, IF

6.09

L-index

#	Paper	IF	Citations
68	Versatile Physiological Functions of Plant GSK3-Like Kinases. <i>Genes</i> , <b>2021</b> , 12,	4.2	3
67	The Evolutionarily Conserved Serine Residues in BRI1 LRR Motifs Are Critical for Protein Secretion. <i>Frontiers in Plant Science</i> , <b>2020</b> , 11, 32	6.2	4
66	Regulation of Three Key Kinases of Brassinosteroid Signaling Pathway. <i>International Journal of Molecular Sciences</i> , <b>2020</b> , 21,	6.3	9
65	Comparative Transcriptomic Analysis to Identify Brassinosteroid Response Genes. <i>Plant Physiology</i> , <b>2020</b> , 184, 1072-1082	6.6	3
64	The Crucial Role of Demannosylating Asparagine-Linked Glycans in ERADicating Misfolded Glycoproteins in the Endoplasmic Reticulum. <i>Frontiers in Plant Science</i> , <b>2020</b> , 11, 625033	6.2	4
63	PAWH1 and PAWH2 are plant-specific components of an Arabidopsis endoplasmic reticulum-associated degradation complex. <i>Nature Communications</i> , <b>2019</b> , 10, 3492	17.4	11
62	Communications Between the Endoplasmic Reticulum and Other Organelles During Abiotic Stress Response in Plants. <i>Frontiers in Plant Science</i> , <b>2019</b> , 10, 749	6.2	30
61	An H3K27me3 demethylase-HSFA2 regulatory loop orchestrates transgenerational thermomemory in Arabidopsis. <i>Cell Research</i> , <b>2019</b> , 29, 379-390	24.7	76
60	Trimming of N-Glycans by the Golgi-Localized $\beta$ 1,2-Mannosidases, MNS1 and MNS2, Is Crucial for Maintaining RSW2 Protein Abundance during Salt Stress in Arabidopsis. <i>Molecular Plant</i> , <b>2018</b> , 11, 678-690	11.4	24
59	Brassinosteroid Signaling Recruits Histone 3 Lysine-27 Demethylation Activity to FLOWERING LOCUS C Chromatin to Inhibit the Floral Transition in Arabidopsis. <i>Molecular Plant</i> , <b>2018</b> , 11, 1135-1146	14.4	35
58	A Temperature-Sensitive Misfolded bri1-301 Receptor Requires Its Kinase Activity to Promote Growth. <i>Plant Physiology</i> , <b>2018</b> , 178, 1704-1719	6.6	12
57	A natural tandem array alleviates epigenetic repression of IPA1 and leads to superior yielding rice. <i>Nature Communications</i> , <b>2017</b> , 8, 14789	17.4	85
56	The SnRK2 kinases modulate miRNA accumulation in Arabidopsis. <i>PLoS Genetics</i> , <b>2017</b> , 13, e1006753	6	56
55	OsREM4.1 Interacts with OsSERK1 to Coordinate the Interlinking between Abscisic Acid and Brassinosteroid Signaling in Rice. <i>Developmental Cell</i> , <b>2016</b> , 38, 201-13	10.2	72
54	Endoplasmic reticulum-associated N-glycan degradation of cold-upregulated glycoproteins in response to chilling stress in Arabidopsis. <i>New Phytologist</i> , <b>2016</b> , 212, 282-96	9.8	38
53	Overexpression of receptor-like kinase ERECTA improves thermotolerance in rice and tomato. <i>Nature Biotechnology</i> , <b>2015</b> , 33, 996-1003	44.5	107
52	EBS7 is a plant-specific component of a highly conserved endoplasmic reticulum-associated degradation system in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2015</b> , 112, 12205-10	11.5	32

51	Genetic and epigenetic control of plant heat responses. <i>Frontiers in Plant Science</i> , <b>2015</b> , 6, 267	6.2	166
50	Endoplasmic reticulum-mediated protein quality control in Arabidopsis. <i>Frontiers in Plant Science</i> , <b>2014</b> , 5, 162	6.2	68
49	An in vivo investigation of amino acid residues critical for the lectin function of Arabidopsis calreticulin 3. <i>Molecular Plant</i> , <b>2013</b> , 6, 985-7	14.4	6
48	A conserved basic residue cluster is essential for the protein quality control function of the Arabidopsis calreticulin 3. <i>Plant Signaling and Behavior</i> , <b>2013</b> , 8, e23864	2.5	5
47	Warm temperatures induce transgenerational epigenetic release of RNA silencing by inhibiting siRNA biogenesis in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2013</b> , 110, 9171-6	11.5	82
46	Auxin controls seed dormancy through stimulation of abscisic acid signaling by inducing ARF-mediated ABI3 activation in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2013</b> , 110, 15485-90	11.5	263
45	The Protein Quality Control of Plant Receptor-Like Kinases in the Endoplasmic Reticulum. <i>Signaling and Communication in Plants</i> , <b>2012</b> , 275-307	1	4
44	Formation of complex extrachromosomal T-DNA structures in <i>Agrobacterium tumefaciens</i> -infected plants. <i>Plant Physiology</i> , <b>2012</b> , 160, 511-22	6.6	35
43	Characterization of temperature-sensitive mutants reveals a role for receptor-like kinase SCRAMBLED/STRUBBELIG in coordinating cell proliferation and differentiation during Arabidopsis leaf development. <i>Plant Journal</i> , <b>2012</b> , 72, 707-20	6.9	28
42	Evolutionarily conserved glycan signal to degrade aberrant brassinosteroid receptors in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2012</b> , 109, 11437-42	11.5	41
41	The Arabidopsis homolog of the mammalian OS-9 protein plays a key role in the endoplasmic reticulum-associated degradation of misfolded receptor-like kinases. <i>Molecular Plant</i> , <b>2012</b> , 5, 929-40	14.4	47
40	Expression of RNA-interference/antisense transgenes by the cognate promoters of target genes is a better gene-silencing strategy to study gene functions in rice. <i>PLoS ONE</i> , <b>2011</b> , 6, e17444	3.7	26
39	Characterization of cp3 reveals a new bri1 allele, bri1-120, and the importance of the LRR domain of BRI1 mediating BR signaling. <i>BMC Plant Biology</i> , <b>2011</b> , 11, 8	5.3	22
38	Conserved endoplasmic reticulum-associated degradation system to eliminate mutated receptor-like kinases in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 870-5	11.5	94
37	Direct involvement of leucine-rich repeats in assembling ligand-triggered receptor-coreceptor complexes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2011</b> , 108, 8073-4	11.5	13
36	Activation-tagged suppressors of a weak brassinosteroid receptor mutant. <i>Molecular Plant</i> , <b>2010</b> , 3, 260-8	14.4	17
35	A direct docking mechanism for a plant GSK3-like kinase to phosphorylate its substrates. <i>Journal of Biological Chemistry</i> , <b>2010</b> , 285, 24646-53	5.4	43
34	BAK7 displays unequal genetic redundancy with BAK1 in brassinosteroid signaling and early senescence in Arabidopsis. <i>Molecules and Cells</i> , <b>2010</b> , 29, 259-66	3.5	33

33	Regulation of the nuclear activities of brassinosteroid signaling. <i>Current Opinion in Plant Biology</i> , <b>2010</b> , 13, 540-7	9.9	55
32	Functional characterization of Arabidopsis thaliana transthyretin-like protein. <i>BMC Plant Biology</i> , <b>2010</b> , 10, 30	5.3	30
31	Regulation of Arabidopsis brassinosteroid signaling by atypical basic helix-loop-helix proteins. <i>Plant Cell</i> , <b>2009</b> , 21, 3781-91	11.6	131
30	Mutations of an alpha1,6 mannosyltransferase inhibit endoplasmic reticulum-associated degradation of defective brassinosteroid receptors in Arabidopsis. <i>Plant Cell</i> , <b>2009</b> , 21, 3792-802	11.6	65
29	A plant-specific calreticulin is a key retention factor for a defective brassinosteroid receptor in the endoplasmic reticulum. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2009</b> , 106, 13612-7	11.5	95
28	BIN2 functions redundantly with other Arabidopsis GSK3-like kinases to regulate brassinosteroid signaling. <i>Plant Physiology</i> , <b>2009</b> , 150, 710-21	6.6	137
27	Analysis of phosphorylation of the BRI1/BAK1 complex in arabidopsis reveals amino acid residues critical for receptor formation and activation of BR signaling. <i>Molecules and Cells</i> , <b>2009</b> , 27, 183-90	3.5	45
26	Multiple mechanism-mediated retention of a defective brassinosteroid receptor in the endoplasmic reticulum of Arabidopsis. <i>Plant Cell</i> , <b>2008</b> , 20, 3418-29	11.6	147
25	Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. <i>Cell Host and Microbe</i> , <b>2008</b> , 4, 17-27	23.4	410
24	Regulation of the Arabidopsis GSK3-like kinase BRASSINOSTEROID-INSENSITIVE 2 through proteasome-mediated protein degradation. <i>Molecular Plant</i> , <b>2008</b> , 1, 338-46	14.4	125
23	Regulation of brassinosteroid signaling. <i>Trends in Plant Science</i> , <b>2007</b> , 12, 37-41	13.1	136
22	Allele-specific suppression of a defective brassinosteroid receptor reveals a physiological role of UGGT in ER quality control. <i>Molecular Cell</i> , <b>2007</b> , 26, 821-30	17.6	148
21	Brassinosteroid-independent function of BRI1/CLV1 chimeric receptors. <i>Functional Plant Biology</i> , <b>2006</b> , 33, 723-730	2.7	7
20	Brassinosteroid signaling: from receptor kinases to transcription factors. <i>Current Opinion in Plant Biology</i> , <b>2005</b> , 8, 526-31	9.9	40
19	BRL1 and BRL3 are novel brassinosteroid receptors that function in vascular differentiation in Arabidopsis. <i>Development (Cambridge)</i> , <b>2004</b> , 131, 5341-51	6.6	391
18	The Arabidopsis transthyretin-like protein is a potential substrate of BRASSINOSTEROID-INSENSITIVE 1. <i>Plant Cell</i> , <b>2004</b> , 16, 2406-17	11.6	91
17	CLAVATA1 dominant-negative alleles reveal functional overlap between multiple receptor kinases that regulate meristem and organ development. <i>Plant Cell</i> , <b>2003</b> , 15, 1198-211	11.6	155
16	Brassinosteroids <b>2003</b> , 214-219		

15	Brassinosteroid Signal Transduction: A Mix of Conservation and Novelty. <i>Journal of Plant Growth Regulation</i> , <b>2003</b> , 22, 298-312	4.7	25
14	Brassinosteroids signal through two receptor-like kinases. <i>Current Opinion in Plant Biology</i> , <b>2003</b> , 6, 494-9	9.9	29
13	The GSK3-like kinase BIN2 phosphorylates and destabilizes BZR1, a positive regulator of the brassinosteroid signaling pathway in Arabidopsis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , <b>2002</b> , 99, 10185-90	11.5	469
12	Two putative BIN2 substrates are nuclear components of brassinosteroid signaling. <i>Plant Physiology</i> , <b>2002</b> , 130, 1221-9	6.6	168
11	BES1 accumulates in the nucleus in response to brassinosteroids to regulate gene expression and promote stem elongation. <i>Cell</i> , <b>2002</b> , 109, 181-91	56.2	858
10	BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling. <i>Cell</i> , <b>2002</b> , 110, 203-12	56.2	861
9	Regulation of brassinosteroid signaling by a GSK3/SHAGGY-like kinase. <i>Science</i> , <b>2002</b> , 295, 1299-301	33.3	451
8	BIN2, a new brassinosteroid-insensitive locus in Arabidopsis. <i>Plant Physiology</i> , <b>2001</b> , 127, 14-22	6.6	356
7	Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich repeat receptor serine/threonine kinase. <i>Plant Physiology</i> , <b>2000</b> , 123, 1247-56	6.6	376
6	Perception of brassinosteroids by the extracellular domain of the receptor kinase BRI1. <i>Science</i> , <b>2000</b> , 288, 2360-3	33.3	381
5	Arabidopsis det2 is defective in the conversion of (24R)-24-methylcholest-4-En-3-one to (24R)-24-methyl-5alpha-cholestan-3-one in brassinosteroid biosynthesis. <i>Plant Physiology</i> , <b>1999</b> , 120, 833-40	6.6	134
4	The phosphoenolpyruvate/phosphate translocator is required for phenolic metabolism, palisade cell development, and plastid-dependent nuclear gene expression. <i>Plant Cell</i> , <b>1999</b> , 11, 1609-22	11.6	228
3	Preparation of DNA from Arabidopsis. <i>Methods in Molecular Biology</i> , <b>1998</b> , 82, 55-60	1.4	24
2	A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. <i>Cell</i> , <b>1997</b> , 90, 929-38	56.2	1285
1	The pc-1 phenotype of <i>Chlamydomonas reinhardtii</i> results from a deletion mutation in the nuclear gene for NADPH:protochlorophyllide oxidoreductase. <i>Plant Molecular Biology</i> , <b>1996</b> , 30, 15-37	4.6	60