

Xavier Subirats

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8734375/publications.pdf>

Version: 2024-02-01

40
papers

748
citations

516215

16
h-index

552369

26
g-index

41
all docs

41
docs citations

41
times ranked

745
citing authors

#	ARTICLE	IF	CITATIONS
1	On the Effect of Organic Solvent Composition on the pH of Buffered HPLC Mobile Phases and the p <i>K</i> _a of Analytes: A Review. Separation and Purification Reviews, 2007, 36, 231-255.	2.8	104
2	Retention models for ionizable compounds in reversed-phase liquid chromatography. Journal of Chromatography A, 2009, 1216, 1756-1775.	1.8	62
3	Viral Uncoating Is Directional: Exit of the Genomic RNA in a Common Cold Virus Starts with the Poly-(A) Tail at the 3' End. PLoS Pathogens, 2013, 9, e1003270.	2.1	43
4	Retention of ionisable compounds on high-performance liquid chromatography. Journal of Chromatography A, 2004, 1059, 33-42.	1.8	40
5	Octanol-Water Partition Constant. , 2020, , 183-208.		36
6	Retention of ionisable compounds on high-performance liquid chromatography XVII. Journal of Chromatography A, 2007, 1138, 203-215.	1.8	35
7	Recent developments in capillary and chip electrophoresis of bioparticles: Viruses, organelles, and cells. Electrophoresis, 2011, 32, 1579-1590.	1.3	34
8	Retention of ionisable compounds on high-performance liquid chromatography. Journal of Chromatography A, 2006, 1121, 170-177.	1.8	31
9	Liposomal Nanocontainers as Models for Viral Infection: Monitoring Viral Genomic RNA Transfer through Lipid Membranes. Journal of Virology, 2011, 85, 8368-8375.	1.5	26
10	Characterization of hydrophilic interaction liquid chromatography retention by a linear free energy relationship. Comparison to reversed- and normal-phase retentions. Analytica Chimica Acta, 2019, 1092, 132-143.	2.6	26
11	Characterization of rhinovirus subviral particles via capillary electrophoresis, electron microscopy and gas-phase electrophoretic mobility molecular analysis: Part I. Electrophoresis, 2012, 33, 1833-1841.	1.3	23
12	Nitromethane as solvent in capillary electrophoresis. Journal of Chromatography A, 2005, 1079, 246-253.	1.8	22
13	Retention of ionisable compounds on high-performance liquid chromatography XVIII: pH variation in mobile phases containing formic acid, piperazine, tris, boric acid or carbonate as buffering systems and acetonitrile as organic modifier. Journal of Chromatography A, 2009, 1216, 2491-2498.	1.8	22
14	Critical comparison of shake-flask, potentiometric and chromatographic methods for lipophilicity evaluation (log P _{o/w}) of neutral, acidic, basic, amphoteric, and zwitterionic drugs. European Journal of Pharmaceutical Sciences, 2018, 122, 331-340.	1.9	21
15	Comparison of methanol and acetonitrile as solvents for the separation of sertindole and its major metabolites by capillary zone electrophoresis. Electrophoresis, 2005, 26, 3315-3324.	1.3	20
16	Lipophilicity of amphoteric and zwitterionic compounds: A comparative study of determination methods. Talanta, 2017, 162, 293-299.	2.9	20
17	Retention-pH profiles of acids and bases in hydrophilic interaction liquid chromatography. Analytica Chimica Acta, 2019, 1050, 176-184.	2.6	18
18	Phenothiazines solution complexity: Determination of p <i>K</i> _a and solubility-pH profiles exhibiting sub-micellar aggregation at 25 and 37°C. European Journal of Pharmaceutical Sciences, 2016, 93, 163-176.	1.9	15

#	ARTICLE	IF	CITATIONS
19	HILIC characterization: Estimation of phase volumes and composition for a zwitterionic column. <i>Analytica Chimica Acta</i> , 2020, 1130, 39-48.	2.6	15
20	Microemulsion electrokinetic chromatography as a suitable tool for lipophilicity determination of acidic, neutral, and basic compounds. <i>Electrophoresis</i> , 2016, 37, 2010-2016.	1.3	11
21	Chasing the elusive hold-up time from an LFER approach. <i>Journal of Chromatography A</i> , 2018, 1571, 176-184.	1.8	11
22	Retention of ionisable compounds on high-performance liquid chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 5445-5448.	1.8	10
23	Characterization of rhinovirus subviral A particles via capillary electrophoresis, electron microscopy and gas phase electrophoretic mobility molecular analysis: Part II. <i>Electrophoresis</i> , 2013, 34, 1600-1609.	1.3	10
24	New discrimination tools for harvest year and varieties of white wines based on hydrophilic interaction liquid chromatography with amperometric detection. <i>Talanta</i> , 2019, 201, 104-110.	2.9	10
25	Liposomal Leakage Induced by Virus-Derived Peptides, Viral Proteins, and Entire Virions: Rapid Analysis by Chip Electrophoresis. <i>Analytical Chemistry</i> , 2010, 82, 8146-8152.	3.2	9
26	Capillary Electrophoresis, Gas-Phase Electrophoretic Mobility Molecular Analysis, and Electron Microscopy: Effective Tools for Quality Assessment and Basic Rhinovirus Research. <i>Methods in Molecular Biology</i> , 2015, 1221, 101-128.	0.4	9
27	High-throughput logPo/w determination from UHPLC measurements: Revisiting the chromatographic hydrophobicity index. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2016, 127, 26-31.	1.4	9
28	Revisiting blood-brain barrier: A chromatographic approach. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 98-109.	1.4	9
29	Ionizable Drug Self-Associations and the Solubility Dependence on pH: Detection of Aggregates in Saturated Solutions Using Mass Spectrometry (ESI-Q-TOF-MS/MS). <i>Molecular Pharmaceutics</i> , 2021, 18, 2311-2321.	2.3	9
30	Volume and composition of semi-adsorbed stationary phases in hydrophilic interaction liquid chromatography. Comparison of water adsorption in common stationary phases and eluents. <i>Journal of Chromatography A</i> , 2021, 1656, 462543.	1.8	8
31	Solute-Solvent Interactions in Hydrophilic Interaction Liquid Chromatography: Characterization of the Retention in a Silica Column by the Abraham Linear Free Energy Relationship Model. <i>Journal of Solution Chemistry</i> , 2022, 51, 1081-1100.	0.6	6
32	Capillary electrophoresis for drug analysis and physicochemical characterization. <i>Handbook of Analytical Separations</i> , 2020, , 633-666.	0.8	5
33	High throughput determination log Po/w/pKa/log Do/w of drugs by combination of UHPLC and CE methods. <i>ADMET and DMPK</i> , 2014, 2, .	1.1	5
34	Methods for pKa Determination (I): Potentiometry, Spectrophotometry, and Capillary Electrophoresis. , 2015, , .		3
35	Enhanced voltammetric performance of sensors based on oxidized 2D layered black phosphorus. <i>Talanta</i> , 2022, 238, 123036.	2.9	3
36	Methods for pKa Determination (II): Sparingly Soluble Compounds and High-Throughput Approaches. , 2015, , .		2

#	ARTICLE	IF	CITATIONS
37	Potentiometric CheqSol and standardized shake-flask solubility methods are complimentary tools in physicochemical profiling. European Journal of Pharmaceutical Sciences, 2020, 148, 105305.	1.9	2
38	Solubility-pH profiles of a free base and its salt: sibutramine as a case study. ADMET and DMPK, 2017, 5, 253-256.	1.1	2
39	Buffers for Reversed-Phase Liquid Chromatography. , 2015, , .		0
40	Lipophilicity determination of acidic compounds: MEEKC as a reliable high-throughput methodology. ADMET and DMPK, 0, , .	1.1	0