

Teh-Hui Kao

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8736656/teh-hui-kao-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85
papers

6,639
citations

40
h-index

81
g-index

87
ext. papers

7,098
ext. citations

9.5
avg, IF

5.35
L-index

#	Paper	IF	Citations
85	Isolation and characterization of rat and human glyceraldehyde-3-phosphate dehydrogenase cDNAs: genomic complexity and molecular evolution of the gene. <i>Nucleic Acids Research</i> , 1985 , 13, 2485-502	20.1	1690
84	S proteins control rejection of incompatible pollen in <i>Petunia inflata</i> . <i>Nature</i> , 1994 , 367, 560-3	50.4	354
83	Identification of the pollen determinant of S-RNase-mediated self-incompatibility. <i>Nature</i> , 2004 , 429, 302-5	50.4	327
82	A GFP-MAP4 reporter gene for visualizing cortical microtubule rearrangements in living epidermal cells. <i>Plant Cell</i> , 1998 , 10, 1927-40	11.6	320
81	Changes in root cap pH are required for the gravity response of the <i>Arabidopsis</i> root. <i>Plant Cell</i> , 2001 , 13, 907-21	11.6	229
80	Collaborative non-self recognition system in S-RNase-based self-incompatibility. <i>Science</i> , 2010 , 330, 796-9	33.3	211
79	The molecular and genetic bases of S-RNase-based self-incompatibility. <i>Plant Cell</i> , 2004 , 16 Suppl, S72-83	11.6	199
78	A GFP-MAP4 Reporter Gene for Visualizing Cortical Microtubule Rearrangements in Living Epidermal Cells. <i>Plant Cell</i> , 1998 , 10, 1927	11.6	199
77	<i>Petunia</i> phospholipase c1 is involved in pollen tube growth. <i>Plant Cell</i> , 2006 , 18, 1438-53	11.6	178
76	Molecular recognition and response in pollen and pistil interactions. <i>Annual Review of Cell and Developmental Biology</i> , 2000 , 16, 333-64	12.6	170
75	Identification and characterization of components of a putative <i>petunia</i> S-locus F-box-containing E3 ligase complex involved in S-RNase-based self-incompatibility. <i>Plant Cell</i> , 2006 , 18, 2531-53	11.6	132
74	S-allele sequence diversity in natural populations of <i>Solanum carolinense</i> (Horsenettle). <i>Heredity</i> , 1995 , 75 (Pt 4), 405-15	3.6	96
73	Cytochrome oxidase subunit II gene of rice has an insertion sequence within the intron. <i>Nucleic Acids Research</i> , 1984 , 12, 7305-15	20.1	96
72	Biochemical models for S-RNase-based self-incompatibility. <i>Molecular Plant</i> , 2008 , 1, 575-85	14.4	92
71	Characterization of Ribonuclease Activity of Three S-Allele-Associated Proteins of <i>Petunia inflata</i> . <i>Plant Physiology</i> , 1991 , 96, 61-8	6.6	86
70	The flanking regions of two <i>Petunia inflata</i> S alleles are heterogeneous and contain repetitive sequences. <i>Plant Molecular Biology</i> , 1992 , 18, 725-37	4.6	86
69	PRK1, a receptor-like kinase of <i>Petunia inflata</i> , is essential for postmeiotic development of pollen. <i>Plant Journal</i> , 1996 , 9, 613-624	6.9	76

68	Self-incompatibility in <i>Petunia inflata</i> : isolation and characterization of cDNAs encoding three S-allele-associated proteins. <i>Sexual Plant Reproduction</i> , 1990 , 3, 130		72
67	Characterization of a pollen-expressed gene encoding a putative pectin esterase of <i>Petunia inflata</i> . <i>Plant Molecular Biology</i> , 1994 , 25, 539-44	4.6	70
66	Evidence that intragenic recombination contributes to allelic diversity of the S-RNase gene at the self-incompatibility (S) locus in <i>Petunia inflata</i> . <i>Plant Physiology</i> , 2001 , 125, 1012-22	6.6	68
65	Chromosome walking in the <i>Petunia inflata</i> self-incompatibility (S-) locus and gene identification in an 881-kb contig containing S2-RNase. <i>Plant Molecular Biology</i> , 2004 , 54, 727-42	4.6	64
64	RNase X2, a pistil-specific ribonuclease from <i>Petunia inflata</i> , shares sequence similarity with solanaceous S proteins. <i>Plant Molecular Biology</i> , 1992 , 20, 1131-41	4.6	63
63	Pea cytochrome oxidase subunit II gene has no intron and generates two mRNA transcripts with different 5' termini. <i>Nucleic Acids Research</i> , 1985 , 13, 3195-212	20.1	63
62	Comparison of <i>Petunia inflata</i> S-Locus F-box protein (Pi SLF) with Pi SLF like proteins reveals its unique function in S-RNase based self-incompatibility. <i>Plant Cell</i> , 2007 , 19, 3593-609	11.6	62
61	Characterization of a predominantly pistil-expressed gene encoding a gamma-thionin-like protein of <i>Petunia inflata</i> . <i>Plant Molecular Biology</i> , 1994 , 26, 459-64	4.6	61
60	S-alleles are retained and expressed in a self-compatible cultivar of <i>Petunia hybrida</i> . <i>Molecular Genetics and Genomics</i> , 1991 , 230, 353-8		61
59	Genetic mapping and molecular characterization of the self-incompatibility (S) locus in <i>Petunia inflata</i> . <i>Plant Molecular Biology</i> , 2003 , 53, 565-80	4.6	60
58	Transcriptome analysis reveals the same 17 S-locus F-box genes in two haplotypes of the self-incompatibility locus of <i>Petunia inflata</i> . <i>Plant Cell</i> , 2014 , 26, 2873-88	11.6	57
57	Cloning and sequencing of cDNAs encoding two self-incompatibility associated proteins in <i>Solanum chacoense</i> . <i>Molecular Genetics and Genomics</i> , 1990 , 224, 341-6		54
56	Ethylene Synthesis and Floral Senescence following Compatible and Incompatible Pollinations in <i>Petunia inflata</i> . <i>Plant Physiology</i> , 1992 , 99, 38-45	6.6	54
55	Rice chloroplast DNA molecules are heterogeneous as revealed by DNA sequences of a cluster of genes. <i>Nucleic Acids Research</i> , 1987 , 15, 611-30	20.1	53
54	S-RNase-based self-incompatibility in <i>Petunia inflata</i> . <i>Annals of Botany</i> , 2011 , 108, 637-46	4.1	50
53	Duplication of the S-locus F-box gene is associated with breakdown of pollen function in an S-haplotype identified in a natural population of self-incompatible <i>Petunia axillaris</i> . <i>Plant Molecular Biology</i> , 2005 , 57, 141-53	4.6	48
52	Identification of self-incompatibility (S-) locus linked pollen cDNA markers in <i>Petunia inflata</i> . <i>Genome</i> , 2000 , 43, 619-627	2.4	47
51	Identification of major lysine residues of S(3)-RNase of <i>Petunia inflata</i> involved in ubiquitin-26S proteasome-mediated degradation in vitro. <i>Plant Journal</i> , 2008 , 54, 1094-104	6.9	46

50	Breakdown of self-incompatibility in a natural population of <i>Petunia axillaris</i> (Solanaceae) in Uruguay containing both self-incompatible and self-compatible plants. <i>Sexual Plant Reproduction</i> , 1999 , 12, 6-13		45
49	Sequence of an S-protein of <i>Lycopersicon peruvianum</i> and comparison with other solanaceous S-proteins. <i>Sexual Plant Reproduction</i> , 1992 , 5, 256		45
48	Genome sequence of a cellulose-producing bacterium, <i>Gluconacetobacter hansenii</i> ATCC 23769. <i>Journal of Bacteriology</i> , 2010 , 192, 4256-7	3.5	43
47	Breakdown of self-incompatibility in a natural population of <i>Petunia axillaris</i> caused by a modifier locus that suppresses the expression of an S-RNase gene. <i>Sexual Plant Reproduction</i> , 2003 , 15, 255-263		41
46	Identification and characterization of non-cellulose-producing mutants of <i>Gluconacetobacter hansenii</i> generated by Tn5 transposon mutagenesis. <i>Journal of Bacteriology</i> , 2013 , 195, 5072-83	3.5	40
45	Breakdown of self-incompatibility in a natural population of <i>Petunia axillaris</i> caused by loss of pollen function. <i>Plant Physiology</i> , 2003 , 131, 1903-12	6.6	40
44	The emerging complexity of self-incompatibility (S-) loci. <i>Sexual Plant Reproduction</i> , 1999 , 12, 1-5		40
43	Isolation and characterization of kinase interacting protein 1, a pollen protein that interacts with the kinase domain of PRK1, a receptor-like kinase of petunia. <i>Plant Physiology</i> , 2001 , 126, 1480-92	6.6	37
42	Identification of the self-incompatibility locus F-box protein-containing complex in <i>Petunia inflata</i> . <i>Plant Reproduction</i> , 2014 , 27, 31-45	3.9	35
41	Construction of a binary bacterial artificial chromosome library of <i>Petunia inflata</i> and the isolation of large genomic fragments linked to the self-incompatibility (S-) locus. <i>Genome</i> , 2000 , 43, 820-826	2.4	35
40	S-RNase-mediated self-incompatibility. <i>Journal of Experimental Botany</i> , 2003 , 54, 115-22	7	34
39	The sensitivity of barley aleurone tissue to gibberellin is heterogeneous and may be spatially determined. <i>Plant Physiology</i> , 1999 , 120, 361-70	6.6	34
38	Four previously identified <i>Petunia inflata</i> S-locus F-box genes are involved in pollen specificity in self-incompatibility. <i>Molecular Plant</i> , 2014 , 7, 567-9	14.4	30
37	Self-incompatibility in <i>Petunia inflata</i> : the relationship between a self-incompatibility locus F-box protein and its non-self S-RNases. <i>Plant Cell</i> , 2013 , 25, 470-85	11.6	30
36	Insight into S-RNase-based self-incompatibility in <i>Petunia</i> : recent findings and future directions. <i>Frontiers in Plant Science</i> , 2015 , 6, 41	6.2	29
35	Identification and characterization of PiORP1, a <i>Petunia</i> oxysterol-binding-protein related protein involved in receptor-kinase mediated signaling in pollen, and analysis of the ORP gene family in <i>Arabidopsis</i> . <i>Plant Molecular Biology</i> , 2006 , 61, 553-65	4.6	28
34	Identification and characterization of a cellulose binding heptapeptide revealed by phage display. <i>Biomacromolecules</i> , 2013 , 14, 1795-805	6.9	27
33	Distribution of self-compatible and self-incompatible populations of <i>Petunia axillaris</i> (Solanaceae) outside Uruguay. <i>Journal of Plant Research</i> , 2006 , 119, 419-30	2.6	27

32	Rice mitochondrial genome contains a rearranged chloroplast gene cluster. <i>Molecular Genetics and Genomics</i> , 1988 , 213, 247-53		27
31	Formation and properties of a covalent complex between elongation factor Tu and Phe-tRNA bearing a photoaffinity probe on its 3-(3-amino-3-carboxypropyl)uridine residue. <i>Journal of Molecular Biology</i> , 1983 , 166, 383-405	6.5	27
30	Gametophytic self-incompatibility: biochemical, molecular genetic, and evolutionary aspects. <i>International Review of Cytology</i> , 1992 , 140, 449-83		21
29	Cloning and sequencing of cDNAs encoding two S proteins of a self-compatible cultivar of <i>Petunia hybrida</i> . <i>Plant Molecular Biology</i> , 1992 , 19, 523-8	4.6	20
28	Identification of self-incompatibility (S-) locus linked pollen cDNA markers in <i>Petunia inflata</i> . <i>Genome</i> , 2000 , 43, 619-627	2.4	19
27	A Mutant S 3 RNase of <i>Petunia inflata</i> Lacking RNase Activity Has an Allele-Specific Dominant Negative Effect on Self-Incompatibility Interactions. <i>Plant Cell</i> , 1997 , 9, 85	11.6	18
26	CRISPR/Cas9-mediated knockout of PiSSK1 reveals essential role of S-locus F-box protein-containing SCF complexes in recognition of non-self S-RNases during cross-compatible pollination in self-incompatible <i>Petunia inflata</i> . <i>Plant Reproduction</i> , 2018 , 31, 129-143	3.9	18
25	Characterization of a Pollen-Expressed Receptor-Like Kinase Gene of <i>Petunia inflata</i> and the Activity of Its Encoded Kinase. <i>Plant Cell</i> , 1994 , 6, 709	11.6	16
24	Isolation and characterization of two cellulose morphology mutants of <i>Gluconacetobacter hansenii</i> ATCC23769 producing cellulose with lower crystallinity. <i>PLoS ONE</i> , 2015 , 10, e0119504	3.7	15
23	Ectopic expression of S-RNase of <i>Petunia inflata</i> in pollen results in its sequestration and non-cytotoxic function. <i>Sexual Plant Reproduction</i> , 2009 , 22, 263-75		15
22	S-Locus F-Box Proteins Are Solely Responsible for S-RNase-Based Self-Incompatibility of Pollen. <i>Plant Cell</i> , 2018 , 30, 2959-2972	11.6	15
21	Molecular mechanisms of self-incompatibility. <i>Current Opinion in Biotechnology</i> , 1996 , 7, 150-154	11.4	14
20	Construction of a binary bacterial artificial chromosome library of <i>Petunia inflata</i> and the isolation of large genomic fragments linked to the self-incompatibility (S-) locus. <i>Genome</i> , 2000 , 43, 820-826	2.4	14
19	Pollen S-locus F-box proteins of <i>Petunia</i> involved in S-RNase-based self-incompatibility are themselves subject to ubiquitin-mediated degradation. <i>Plant Journal</i> , 2015 , 83, 213-23	6.9	13
18	The amino terminal F-box domain of <i>Petunia inflata</i> S-locus F-box protein is involved in the S-RNase-based self-incompatibility mechanism. <i>AoB PLANTS</i> , 2011 , 2011, plr016	2.9	13
17	Isolation and nucleotide sequence of the pea cytochrome oxidase subunit I gene. <i>Plant Molecular Biology</i> , 1989 , 13, 121-4	4.6	13
16	AcsA-AcsB: The core of the cellulose synthase complex from <i>Gluconacetobacter hansenii</i> ATCC23769. <i>Enzyme and Microbial Technology</i> , 2016 , 82, 58-65	3.8	11
15	Carbohydrate Moiety of the <i>Petunia inflata</i> S 3 Protein Is Not Required for Self-Incompatibility Interactions between Pollen and Pistil. <i>Plant Cell</i> , 1994 , 6, 1933	11.6	11

14	Self-incompatibility in <i>Petunia</i> : a self/nonself-recognition mechanism employing S-locus F-box proteins and S-RNase to prevent inbreeding. <i>Wiley Interdisciplinary Reviews: Developmental Biology</i> , 2012 , 1, 267-75	5.9	10
13	Processing of cellulose synthase (AcsAB) from <i>Gluconacetobacter hansenii</i> 23769. <i>Archives of Biochemistry and Biophysics</i> , 2013 , 529, 92-8	4.1	9
12	Differentiation in the status of self-incompatibility among <i>Calibrachoa</i> species (Solanaceae). <i>Journal of Plant Research</i> , 2002 , 115, 185-93	2.6	9
11	All 17 S-locus F-box proteins of the S2 - and S3 -haplotypes of <i>Petunia inflata</i> are assembled into similar SCF complexes with a specific function in self-incompatibility. <i>Plant Journal</i> , 2016 , 87, 606-16	6.9	9
10	Sequence of the chloroplast-encoded atpB-atpE-trnM gene clusters from rice. <i>Nucleic Acids Research</i> , 1987 , 15, 4358-9	20.1	7
9	Expression of two S-ribonucleases of <i>Petunia inflata</i> using baculovirus expression system. <i>Biochemical and Biophysical Research Communications</i> , 1992 , 187, 299-304	3.4	6
8	Use of Domain-Swapping to Identify Candidate Amino Acids Involved in Differential Interactions between Two Allelic Variants of Type-1 S-Locus F-Box Protein and S3-RNase in <i>Petunia inflata</i> . <i>Plant and Cell Physiology</i> , 2018 , 59, 234-247	4.9	5
7	Changes in Root Cap pH Are Required for the Gravity Response of the Arabidopsis Root. <i>Plant Cell</i> , 2001 , 13, 907	11.6	5
6	Functional characterization of two chimeric proteins between a <i>Petunia inflata</i> S-locus F-box protein, PiSLF2, and a PiSLF-like protein, PiSLFLb-S2. <i>Plant Molecular Biology</i> , 2010 , 74, 279-92	4.6	4
5	Sequence analysis of the <i>Petunia inflata</i> S-locus region containing 17 S-Locus F-Box genes and the S-RNase gene involved in self-incompatibility. <i>Plant Journal</i> , 2020 , 104, 1348-1368	6.9	4
4	Cytological study of pollen tube growth and early seed development in <i>Petunia inflata</i> 1997 , 40, 212-219		3
3	Interaction of PRK1 receptor-like kinase with a putative eIF2B beta-subunit in tobacco. <i>Molecules and Cells</i> , 2000 , 10, 626-32	3.5	3
2	Disputed Ancestry: Comments on a Model for the Origin of Incompatibility in Flowering Plants [with Reply]. <i>Plant Cell</i> , 1995 , 7, 661	11.6	2
1	S-RNase-Based Self-Incompatibility in <i>Petunia</i> : A Complex Non-Self Recognition System Between Pollen and Pistil 2014 , 289-303		