## Teh-Hui Kao

## List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/8736656/teh-hui-kao-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

85 6,639 40 81 g-index

87 7,098 9.5 5.35 ext. papers ext. citations avg, IF L-index

#	Paper	IF	Citations
85	Sequence analysis of the Petunia inflata S-locus region containing 17 S-Locus F-Box genes and the S-RNase gene involved in self-incompatibility. <i>Plant Journal</i> , <b>2020</b> , 104, 1348-1368	6.9	4
84	Use of Domain-Swapping to Identify Candidate Amino Acids Involved in Differential Interactions between Two Allelic Variants of Type-1 S-Locus F-Box Protein and S3-RNase in Petunia inflata. <i>Plant and Cell Physiology</i> , <b>2018</b> , 59, 234-247	4.9	5
83	CRISPR/Cas9-mediated knockout of PiSSK1 reveals essential role of S-locus F-box protein-containing SCF complexes in recognition of non-self S-RNases during cross-compatible pollination in self-incompatible Petunia inflata. <i>Plant Reproduction</i> , <b>2018</b> , 31, 129-143	3.9	18
82	S-Locus F-Box Proteins Are Solely Responsible for S-RNase-Based Self-Incompatibility of Pollen. <i>Plant Cell</i> , <b>2018</b> , 30, 2959-2972	11.6	15
81	AcsA-AcsB: The core of the cellulose synthase complex from Gluconacetobacter hansenii ATCC23769. Enzyme and Microbial Technology, <b>2016</b> , 82, 58-65	3.8	11
80	All 17 S-locus F-box proteins of the S2 - and S3 -haplotypes of Petunia inflata are assembled into similar SCF complexes with a specific function in self-incompatibility. <i>Plant Journal</i> , <b>2016</b> , 87, 606-16	6.9	9
79	Insight into S-RNase-based self-incompatibility in Petunia: recent findings and future directions. <i>Frontiers in Plant Science</i> , <b>2015</b> , 6, 41	6.2	29
78	Pollen S-locus F-box proteins of Petunia involved in S-RNase-based self-incompatibility are themselves subject to ubiquitin-mediated degradation. <i>Plant Journal</i> , <b>2015</b> , 83, 213-23	6.9	13
77	Isolation and characterization of two cellulose morphology mutants of Gluconacetobacter hansenii ATCC23769 producing cellulose with lower crystallinity. <i>PLoS ONE</i> , <b>2015</b> , 10, e0119504	3.7	15
76	Identification of the self-incompatibility locus F-box protein-containing complex in Petunia inflata. <i>Plant Reproduction</i> , <b>2014</b> , 27, 31-45	3.9	35
75	Transcriptome analysis reveals the same 17 S-locus F-box genes in two haplotypes of the self-incompatibility locus of Petunia inflata. <i>Plant Cell</i> , <b>2014</b> , 26, 2873-88	11.6	57
74	Four previously identified Petunia inflata S-locus F-box genes are involved in pollen specificity in self-incompatibility. <i>Molecular Plant</i> , <b>2014</b> , 7, 567-9	14.4	30
73	S-RNase-Based Self-Incompatibility in Petunia: A Complex Non-Self Recognition System Between Pollen and Pistil <b>2014</b> , 289-303		
72	Identification and characterization of a cellulose binding heptapeptide revealed by phage display. <i>Biomacromolecules</i> , <b>2013</b> , 14, 1795-805	6.9	27
71	Processing of cellulose synthase (AcsAB) from Gluconacetobacter hansenii 23769. <i>Archives of Biochemistry and Biophysics</i> , <b>2013</b> , 529, 92-8	4.1	9
70	Identification and characterization of non-cellulose-producing mutants of Gluconacetobacter hansenii generated by Tn5 transposon mutagenesis. <i>Journal of Bacteriology</i> , <b>2013</b> , 195, 5072-83	3.5	40
69	Self-incompatibility in Petunia inflata: the relationship between a self-incompatibility locus F-box protein and its non-self S-RNases. <i>Plant Cell</i> , <b>2013</b> , 25, 470-85	11.6	30

## (2004-2012)

68	Self-incompatibility in Petunia: a self/nonself-recognition mechanism employing S-locus F-box proteins and S-RNase to prevent inbreeding. <i>Wiley Interdisciplinary Reviews: Developmental Biology</i> , <b>2012</b> , 1, 267-75	5.9	10
67	The amino terminal F-box domain of Petunia inflata S-locus F-box protein is involved in the S-RNase-based self-incompatibility mechanism. <i>AoB PLANTS</i> , <b>2011</b> , 2011, plr016	2.9	13
66	S-RNase-based self-incompatibility in Petunia inflata. <i>Annals of Botany</i> , <b>2011</b> , 108, 637-46	4.1	50
65	Genome sequence of a cellulose-producing bacterium, Gluconacetobacter hansenii ATCC 23769. Journal of Bacteriology, <b>2010</b> , 192, 4256-7	3.5	43
64	Collaborative non-self recognition system in S-RNase-based self-incompatibility. <i>Science</i> , <b>2010</b> , 330, 796	5 <del>-9</del> 3.3	211
63	Functional characterization of two chimeric proteins between a Petunia inflata S-locus F-box protein, PiSLF2, and a PiSLF-like protein, PiSLFLb-S2. <i>Plant Molecular Biology</i> , <b>2010</b> , 74, 279-92	4.6	4
62	Ectopic expression of S-RNase of Petunia inflata in pollen results in its sequestration and non-cytotoxic function. <i>Sexual Plant Reproduction</i> , <b>2009</b> , 22, 263-75		15
61	Identification of major lysine residues of S(3)-RNase of Petunia inflata involved in ubiquitin-26S proteasome-mediated degradation in vitro. <i>Plant Journal</i> , <b>2008</b> , 54, 1094-104	6.9	46
60	Biochemical models for S-RNase-based self-incompatibility. <i>Molecular Plant</i> , <b>2008</b> , 1, 575-85	14.4	92
59	Comparison of Petunia inflata S-Locus F-box protein (Pi SLF) with Pi SLF like proteins reveals its unique function in S-RNase based self-incompatibility. <i>Plant Cell</i> , <b>2007</b> , 19, 3593-609	11.6	62
58	Identification and characterization of components of a putative petunia S-locus F-box-containing E3 ligase complex involved in S-RNase-based self-incompatibility. <i>Plant Cell</i> , <b>2006</b> , 18, 2531-53	11.6	132
57	Petunia phospholipase c1 is involved in pollen tube growth. <i>Plant Cell</i> , <b>2006</b> , 18, 1438-53	11.6	178
56	Identification and characterization of PiORP1, a Petunia oxysterol-binding-protein related protein involved in receptor-kinase mediated signaling in pollen, and analysis of the ORP gene family in Arabidopsis. <i>Plant Molecular Biology</i> , <b>2006</b> , 61, 553-65	4.6	28
55	Distribution of self-compatible and self-incompatible populations of Petunia axillaris (Solanaceae) outside Uruguay. <i>Journal of Plant Research</i> , <b>2006</b> , 119, 419-30	2.6	27
54	Duplication of the S-locus F-box gene is associated with breakdown of pollen function in an S-haplotype identified in a natural population of self-incompatible Petunia axillaris. <i>Plant Molecular Biology</i> , <b>2005</b> , 57, 141-53	4.6	48
53	The molecular and genetic bases of S-RNase-based self-incompatibility. <i>Plant Cell</i> , <b>2004</b> , 16 Suppl, S72-8	8 <b>3</b> 1.6	199
52	Chromosome walking in the Petunia inflata self-incompatibility (S-) locus and gene identification in an 881-kb contig containing S2-RNase. <i>Plant Molecular Biology</i> , <b>2004</b> , 54, 727-42	4.6	64
51	Identification of the pollen determinant of S-RNase-mediated self-incompatibility. <i>Nature</i> , <b>2004</b> , 429, 302-5	50.4	327

50	Genetic mapping and molecular characterization of the self-incompatibility (S) locus in Petunia inflata. <i>Plant Molecular Biology</i> , <b>2003</b> , 53, 565-80	4.6	60
49	Breakdown of self-incompatibility in a natural population of Petunia axillaris caused by a modifier locus that suppresses the expression of an S-RNase gene. <i>Sexual Plant Reproduction</i> , <b>2003</b> , 15, 255-263		41
48	Breakdown of self-incompatibility in a natural population of Petunia axillaris caused by loss of pollen function. <i>Plant Physiology</i> , <b>2003</b> , 131, 1903-12	6.6	40
47	S-RNase-mediated self-incompatibility. <i>Journal of Experimental Botany</i> , <b>2003</b> , 54, 115-22	7	34
46	Differentiation in the status of self-incompatibility among Calibrachoa species (Solanaceae). Journal of Plant Research, <b>2002</b> , 115, 185-93	2.6	9
45	Changes in root cap pH are required for the gravity response of the Arabidopsis root. <i>Plant Cell</i> , <b>2001</b> , 13, 907-21	11.6	229
44	Isolation and characterization of kinase interacting protein 1, a pollen protein that interacts with the kinase domain of PRK1, a receptor-like kinase of petunia. <i>Plant Physiology</i> , <b>2001</b> , 126, 1480-92	6.6	37
43	Changes in Root Cap pH Are Required for the Gravity Response of the Arabidopsis Root. <i>Plant Cell</i> , <b>2001</b> , 13, 907	11.6	5
42	Evidence that intragenic recombination contributes to allelic diversity of the S-RNase gene at the self-incompatibility (S) locus in Petunia inflata. <i>Plant Physiology</i> , <b>2001</b> , 125, 1012-22	6.6	68
41	Interaction of PRK1 receptor-like kinase with a putative elF2B beta-subunit in tobacco. <i>Molecules and Cells</i> , <b>2000</b> , 10, 626-32	3.5	3
40	Molecular recognition and response in pollen and pistil interactions. <i>Annual Review of Cell and Developmental Biology</i> , <b>2000</b> , 16, 333-64	12.6	170
39	Construction of a binary bacterial artificial chromosome library of Petunia inflata and the isolation of large genomic fragments linked to the self-incompatibility (S-) locus. <i>Genome</i> , <b>2000</b> , 43, 820-826	2.4	35
38	Identification of self-incompatibility (S-) locus linked pollen cDNA markers in Petunia inflata. <i>Genome</i> , <b>2000</b> , 43, 619-627	2.4	47
37	Identification of self-incompatibility (S-) locus linked pollen cDNA markers in Petunia inflata. <i>Genome</i> , <b>2000</b> , 43, 619-627	2.4	19
36	Construction of a binary bacterial artificial chromosome library of Petunia inflata and the isolation of large genomic fragments linked to the self-incompatibility (S-) locus. <i>Genome</i> , <b>2000</b> , 43, 820-826	2.4	14
35	The sensitivity of barley aleurone tissue to gibberellin is heterogeneous and may Be spatially determined. <i>Plant Physiology</i> , <b>1999</b> , 120, 361-70	6.6	34
34	The emerging complexity of self-incompatibility (S-) loci. Sexual Plant Reproduction, 1999, 12, 1-5		40
33	Breakdown of self-incompatibility in a natural population of Petunia axillaris (Solanaceae) in Uruguay containing both self-incompatible and self-compatible plants. <i>Sexual Plant Reproduction</i> , <b>1999</b> , 12, 6-13		45

32	A GFP-MAP4 reporter gene for visualizing cortical microtubule rearrangements in living epidermal cells. <i>Plant Cell</i> , <b>1998</b> , 10, 1927-40	11.6	320
31	A GFP-MAP4 Reporter Gene for Visualizing Cortical Microtubule Rearrangements in Living Epidermal Cells. <i>Plant Cell</i> , <b>1998</b> , 10, 1927	11.6	199
30	A Mutant S 3 RNase of Petunia inflata Lacking RNase Activity Has an Allele-Specific Dominant Negative Effect on Self-Incompatibility Interactions. <i>Plant Cell</i> , <b>1997</b> , 9, 85	11.6	18
29	Cytological study of pollen tube growth and early seed development inPetunia inflata <b>1997</b> , 40, 212-2	19	3
28	Molecular mechanisms of self-incompatibility. Current Opinion in Biotechnology, 1996, 7, 150-154	11.4	14
27	PRK1, a receptor-like kinase of Petunia inflata, is essential for postmeiotic development of pollen. <i>Plant Journal</i> , <b>1996</b> , 9, 613-624	6.9	76
26	S-allele sequence diversity in natural populations of Solanum carolinense (Horsenettle). <i>Heredity</i> , <b>1995</b> , 75 ( Pt 4), 405-15	3.6	96
25	Disputed Ancestry: Comments on a Model for the Origin of Incompatibility in Flowering Plants [with Reply]. <i>Plant Cell</i> , <b>1995</b> , 7, 661	11.6	2
24	Carbohydrate Moiety of the Petunia inflata S 3 Protein Is Not Required for Self-Incompatibility Interactions between Pollen and Pistil. <i>Plant Cell</i> , <b>1994</b> , 6, 1933	11.6	11
23	Characterization of a predominantly pistil-expressed gene encoding a gamma-thionin-like protein of Petunia inflata. <i>Plant Molecular Biology</i> , <b>1994</b> , 26, 459-64	4.6	61
22	Characterization of a pollen-expressed gene encoding a putative pectin esterase of Petunia inflata. <i>Plant Molecular Biology</i> , <b>1994</b> , 25, 539-44	4.6	70
21	S proteins control rejection of incompatible pollen in Petunia inflata. <i>Nature</i> , <b>1994</b> , 367, 560-3	50.4	354
20	Characterization of a Pollen-Expressed Receptor-Like Kinase Gene of Petunia inflata and the Activity of Its Encoded Kinase. <i>Plant Cell</i> , <b>1994</b> , 6, 709	11.6	16
19	Gametophytic self-incompatibility: biochemical, molecular genetic, and evolutionary aspects. <i>International Review of Cytology</i> , <b>1992</b> , 140, 449-83		21
18	Expression of two S-ribonucleases of Petunia inflata using baculovirus expression system. <i>Biochemical and Biophysical Research Communications</i> , <b>1992</b> , 187, 299-304	3.4	6
17	The flanking regions of two Petunia inflata S alleles are heterogeneous and contain repetitive sequences. <i>Plant Molecular Biology</i> , <b>1992</b> , 18, 725-37	4.6	86
16	Cloning and sequencing of cDNAs encoding two S proteins of a self-compatible cultivar of Petunia hybrida. <i>Plant Molecular Biology</i> , <b>1992</b> , 19, 523-8	4.6	20
15	RNase X2, a pistil-specific ribonuclease from Petunia inflata, shares sequence similarity with solanaceous S proteins. <i>Plant Molecular Biology</i> , <b>1992</b> , 20, 1131-41	4.6	63

14	Sequence of an S-protein of Lycopersicon peruvianum and comparison with other solanaceous S-proteins. <i>Sexual Plant Reproduction</i> , <b>1992</b> , 5, 256		45
13	Ethylene Synthesis and Floral Senescence following Compatible and Incompatible Pollinations in Petunia inflata. <i>Plant Physiology</i> , <b>1992</b> , 99, 38-45	6.6	54
12	S-alleles are retained and expressed in a self-compatible cultivar of Petunia hybrida. <i>Molecular Genetics and Genomics</i> , <b>1991</b> , 230, 353-8		61
11	Characterization of Ribonuclease Activity of Three S-Allele-Associated Proteins of Petunia inflata. <i>Plant Physiology</i> , <b>1991</b> , 96, 61-8	6.6	86
10	Cloning and sequencing of cDNAs encoding two self-incompatibility associated proteins in Solanum chacoense. <i>Molecular Genetics and Genomics</i> , <b>1990</b> , 224, 341-6		54
9	Self-incompatibility in Petunia inflata: isolation and characterization of cDNAs encoding three S-allele-associated proteins. <i>Sexual Plant Reproduction</i> , <b>1990</b> , 3, 130		72
8	Isolation and nucleotide sequence of the pea cytochrome oxidase subunit I gene. <i>Plant Molecular Biology</i> , <b>1989</b> , 13, 121-4	4.6	13
7	Rice mitochondrial genome contains a rearranged chloroplast gene cluster. <i>Molecular Genetics and Genomics</i> , <b>1988</b> , 213, 247-53		27
6	Sequence of the chloroplast-encoded atpB-atpE-trnM gene clusters from rice. <i>Nucleic Acids Research</i> , <b>1987</b> , 15, 4358-9	20.1	7
5	Rice chloroplast DNA molecules are heterogeneous as revealed by DNA sequences of a cluster of genes. <i>Nucleic Acids Research</i> , <b>1987</b> , 15, 611-30	20.1	53
4	Pea cytochrome oxidase subunit II gene has no intron and generates two mRNA transcripts with different 5V termini. <i>Nucleic Acids Research</i> , <b>1985</b> , 13, 3195-212	20.1	63
3	Isolation and characterization of rat and human glyceraldehyde-3-phosphate dehydrogenase cDNAs: genomic complexity and molecular evolution of the gene. <i>Nucleic Acids Research</i> , <b>1985</b> , 13, 248.	5- <del>3</del> 02	1690
2	Cytochrome oxidase subunit II gene of rice has an insertion sequence within the intron. <i>Nucleic Acids Research</i> , <b>1984</b> , 12, 7305-15	20.1	96
1	Formation and properties of a covalent complex between elongation factor Tu and Phe-tRNA bearing a photoaffinity probe on its 3-(3-amino-3-carboxypropyl)uridine residue. <i>Journal of Molecular Biology</i> 1983, 166, 383-405	6.5	27