

# Noriyuki Kitagawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8743189/publications.pdf>

Version: 2024-02-01

34  
papers

328  
citations

1039880

9  
h-index

996849

15  
g-index

35  
all docs

35  
docs citations

35  
times ranked

383  
citing authors

#	ARTICLE	IF	CITATIONS
1	Hemoglobin concentration and incident metabolic syndrome: a population-based large-scale cohort study. <i>Endocrine</i> , 2015, 50, 390-396.	1.1	38
2	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021, 12, 783363.	1.5	32
3	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021, 40, 1613-1620.	2.3	22
4	High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019, 66, 369-377.	0.7	19
5	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 3729.	1.7	18
6	Prognostic significance of day-by-day variability of home blood pressure on progression to macroalbuminuria in patients with diabetes. <i>Journal of Hypertension</i> , 2018, 36, 1068-1075.	0.3	17
7	Creatinine/(cystatin C $\times$ body weight) ratio is associated with skeletal muscle mass index. <i>Endocrine Journal</i> , 2020, 67, 733-740.	0.7	16
8	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 72.	1.7	16
9	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021, 13, 2335.	1.7	15
10	Impact of extracellular $\leftrightarrow$ intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1202-1211.	1.1	11
11	Predictive power of home blood pressure measurement for cardiovascular outcomes in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2021, 44, 348-354.	1.5	11
12	Postprandial hyperglycemia was ameliorated by taking metformin 30 min before a meal than taking metformin with a meal; a randomized, open-label, crossover pilot study. <i>Endocrine</i> , 2016, 52, 271-276.	1.1	10
13	Threshold value of home pulse pressure predicting arterial stiffness in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Journal of Clinical Hypertension</i> , 2018, 20, 472-477.	1.0	10
14	Dipeptidyl peptidase-4 inhibitors have adverse effects for the proliferation of human T cells. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2018, 63, 106-112.	0.6	10
15	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020, 12, 2174.	1.7	10
16	High-sensitivity cardiac troponin T is associated with coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2015, 9, 209-214.	0.7	9
17	Isolated high home systolic blood pressure in patients with type 2 diabetes is a prognostic factor for the development of diabetic nephropathy: KAMOGAWA-HBP study. <i>Diabetes Research and Clinical Practice</i> , 2019, 158, 107920.	1.1	8
18	The Association Between Taste Impairment and Serum Zinc Concentration in Adult Patients With Type 2 Diabetes. <i>Canadian Journal of Diabetes</i> , 2018, 42, 520-524.	0.4	6

#	ARTICLE	IF	CITATIONS
19	Which Measurement of Blood Pressure Is More Associated With Albuminuria in Patients With Type 2 Diabetes: Central Blood Pressure or Peripheral Blood Pressure?. <i>Journal of Clinical Hypertension</i> , 2016, 18, 790-795.	1.0	5
20	Prognostic significance of home pulse pressure for progression of diabetic nephropathy: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2018, 41, 363-371.	1.5	5
21	Handgrip measurement as a useful benchmark for locomotive syndrome in patients with type 2 diabetes mellitus: A KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2020, 11, 1602-1611.	1.1	5
22	Eating Speed Is Associated with the Presence of Sarcopenia in Older Patients with Type 2 Diabetes: A Cross-Sectional Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2022, 14, 759.	1.7	5
23	Serum levels of mac-2 binding protein are associated with diabetic microangiopathy and macroangiopathy in people with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001189.	1.2	4
24	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 1488.	1.7	4
25	Diabetic nephropathy ameliorated in patients with normal home blood pressure compared to those with isolated high home systolic blood pressure: A 5-year prospective cohort study among patients with type 2 diabetes mellitus. <i>Diabetes and Vascular Disease Research</i> , 2022, 19, 147916412210981.	0.9	4
26	Impact of Eating Speed on Muscle Mass in Older Patients With Type 2 Diabetes: A Prospective Study of KAMOGAWA-DM Cohort. <i>Frontiers in Nutrition</i> , 0, 9, .	1.6	4
27	Impact of Isolated High Home Systolic Blood Pressure and Diabetic Nephropathy in Patients with Type 2 Diabetes Mellitus: A 5-Year Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2021, 10, 1929.	1.0	3
28	Development of application to automatically calculate mean amplitude of glycaemic excursions using intermittently scanned continuous glucose monitoring data. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 2155-2160.	2.2	3
29	Effect of Teriparatide on Bone Mineral Density and Trabecular Bone Score in Type 2 Diabetic Patients with Osteoporosis: A Retrospective Cohort Study. <i>Medicina (Lithuania)</i> , 2022, 58, 481.	0.8	3
30	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Experimental Gerontology</i> , 2022, 159, 111671.	1.2	2
31	Correlation between Liver Stiffness by Two-Dimensional Shear Wave Elastography and Waist Circumference in Japanese Local Citizens with Abdominal Obesity. <i>Journal of Clinical Medicine</i> , 2021, 10, 1971.	1.0	1
32	Association of Estimated Salt and Miso Intake with the Prevalence of Obesity in People with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021, 13, 3014.	1.7	1
33	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022, 11, 1623.	1.8	1
34	Reply to: Serum high-sensitivity cardiac troponin T and coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2016, 10, e3-e4.	0.7	0