

Noriyuki Kitagawa

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8743189/noriyuki-kitagawa-publications-by-citations.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

28

papers

142

citations

7

h-index

10

g-index

35

ext. papers

235

ext. citations

4.4

avg, IF

2.34

L-index

#	Paper	IF	Citations
28	Hemoglobin concentration and incident metabolic syndrome: a population-based large-scale cohort study. <i>Endocrine</i> , 2015 , 50, 390-6	4	26
27	Prognostic significance of day-by-day variability of home blood pressure on progression to macroalbuminuria in patients with diabetes. <i>Journal of Hypertension</i> , 2018 , 36, 1068-1075	1.9	14
26	High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocrine Journal</i> , 2019 , 66, 369-377	2.9	10
25	Threshold value of home pulse pressure predicting arterial stiffness in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Journal of Clinical Hypertension</i> , 2018 , 20, 472-477	2.3	9
24	Creatinine/(cystatin C/body weight) ratio is associated with skeletal muscle mass index. <i>Endocrine Journal</i> , 2020 , 67, 733-740	2.9	9
23	Dipeptidyl peptidase-4 inhibitors have adverse effects for the proliferation of human T cells. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2018 , 63, 106-112	3.1	8
22	Isolated high home systolic blood pressure in patients with type 2 diabetes is a prognostic factor for the development of diabetic nephropathy: KAMOGAWA-HBP study. <i>Diabetes Research and Clinical Practice</i> , 2019 , 158, 107920	7.4	7
21	Short energy intake is associated with muscle mass loss in older patients with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort. <i>Clinical Nutrition</i> , 2021 , 40, 1613-1620	5.9	7
20	Postprandial hyperglycemia was ameliorated by taking metformin 30 min before a meal than taking metformin with a meal; a randomized, open-label, crossover pilot study. <i>Endocrine</i> , 2016 , 52, 271-6	4	6
19	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with a Low Prevalence of Sarcopenia in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 13,	6.7	6
18	High-sensitivity cardiac troponin T is associated with coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2015 , 9, 209-14	2.8	5
17	Eating Fast Is Associated with Nonalcoholic Fatty Liver Disease in Men But Not in Women with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2020 , 12,	6.7	5
16	Predictive power of home blood pressure measurement for cardiovascular outcomes in patients with type 2 diabetes: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2021 , 44, 348-354	4.7	5
15	Prognostic significance of home pulse pressure for progression of diabetic nephropathy: KAMOGAWA-HBP study. <i>Hypertension Research</i> , 2018 , 41, 363-371	4.7	4
14	Which Measurement of Blood Pressure Is More Associated With Albuminuria in Patients With Type 2 Diabetes: Central Blood Pressure or Peripheral Blood Pressure?. <i>Journal of Clinical Hypertension</i> , 2016 , 18, 790-5	2.3	4
13	Handgrip measurement as a useful benchmark for locomotive syndrome in patients with type 2 diabetes mellitus: A KAMOGAWA-DM cohort study. <i>Journal of Diabetes Investigation</i> , 2020 , 11, 1602-1611	3.9	3
12	The Association Between Taste Impairment and Serum Zinc Concentration in Adult Patients With Type 2 Diabetes. <i>Canadian Journal of Diabetes</i> , 2018 , 42, 520-524	2.1	3

11	Impact of Isolated High Home Systolic Blood Pressure and Diabetic Nephropathy in Patients with Type 2 Diabetes Mellitus: A 5-Year Prospective Cohort Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
10	Vitamin Intake and Loss of Muscle Mass in Older People with Type 2 Diabetes: A Prospective Study of the KAMOGAWA-DM Cohort. <i>Nutrients</i> , 2021 , 13,	6.7	2
9	Association between Geriatric Nutrition Risk Index and The Presence of Sarcopenia in People with Type 2 Diabetes Mellitus: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
8	Impact of extracellular-to-intracellular fluid volume ratio on albuminuria in patients with type 2 diabetes: A cross-sectional and longitudinal cohort study. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1202-1211	3.9	1
7	Habitual Miso (Fermented Soybean Paste) Consumption Is Associated with Glycemic Variability in Patients with Type 2 Diabetes: A Cross-Sectional Study. <i>Nutrients</i> , 2021 , 13,	6.7	1
6	Diabetic nephropathy ameliorated in patients with normal home blood pressure compared to those with isolated high home systolic blood pressure: A 5-year prospective cohort study among patients with type 2 diabetes mellitus.. <i>Diabetes and Vascular Disease Research</i> , 2022 , 19, 14791641221098193	3.3	1
5	Sarcopenia Is Associated With a Risk of Mortality in People With Type 2 Diabetes Mellitus. <i>Frontiers in Endocrinology</i> , 2021 , 12, 783363	5.7	0
4	Characterization of Peripheral Blood TCR in Patients with Type 1 Diabetes Mellitus by BD Rhapsody™ VDJ CDR3 Assay. <i>Cells</i> , 2022 , 11, 1623	7.9	0
3	Reply to: Serum high-sensitivity cardiac troponin T and coronary artery calcification. <i>Journal of Cardiovascular Computed Tomography</i> , 2016 , 10, e3-4	2.8	
2	Relationship between eosinophils counts and muscle mass decline in older people with type 2 diabetes: A prospective study of the KAMOGAWA-DM cohort.. <i>Experimental Gerontology</i> , 2022 , 159, 111671	4.5	1
1	Development of application to automatically calculate mean amplitude of glycaemic excursions using intermittently scanned continuous glucose monitoring data. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 2155-2160	6.7	