Masoud Tabari Kouchaksaraei

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8744781/publications.pdf

Version: 2024-02-01

687363 940533 17 423 13 16 g-index citations h-index papers 17 17 17 500 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	Drought Effects on Morpho-Physiological and Biochemical Traits in Persian Oak and Black Poplar Seedlings. Forests, 2022, 13, 399.	2.1	8
2	Drought and Pathogen Effects on Survival, Leaf Physiology, Oxidative Damage, and Defense in Two Middle Eastern Oak Species. Forests, 2021, 12, 247.	2.1	15
3	Dual inoculations of arbuscular mycorrhizal fungi and plant growth-promoting rhizobacteria boost drought resistance and essential oil yield of common myrtle. Forest Ecology and Management, 2021, 497, 119478.	3. 2	34
4	Differential physiological and biochemical responses of <i>Quercus infectoria</i> and <i>Q. libani</i> to drought and charcoal disease. Physiologia Plantarum, 2020, 168, 876-892.	5.2	14
5	Assessment of anticipated performance index of some deciduous plant species under dust air pollution. Environmental Science and Pollution Research, 2020, 27, 38987-38994.	5. 3	25
6	The effect of biochar amendment on the growth, morphology and physiology of Quercus castaneifolia seedlings under water-deficit stress. European Journal of Forest Research, 2019, 138, 967-979.	2.5	29
7	The response of English yew (Taxus baccata L.) to climate change in the Caspian Hyrcanian Mixed Forest ecoregion. Regional Environmental Change, 2019, 19, 1495-1506.	2.9	26
8	The impact of nanoparticles zero-valent iron (nZVI) and rhizosphere microorganisms on the phytoremediation ability of white willow and its response. Environmental Science and Pollution Research, 2019, 26, 10776-10789.	5. 3	57
9	Change in biochemical parameters of Persian oak (Quercus brantii Lindl.) seedlings inoculated by pathogens of charcoal disease under water deficit conditions. Trees - Structure and Function, 2018, 32, 1595-1608.	1.9	16
10	<i>Cantharellus alborufescens</i> and <i>C. ferruginascens</i> (Cantharellaceae, Basidiomycota) New to Iran. Cryptogamie, Mycologie, 2018, 39, 299-310.	1.0	7
11	Growth and physiological responses of <i>Quercus brantii</i> seedlings inoculated with <i>Biscogniauxia mediterranea</i> and <i>Obolarina persica</i> under drought stress. Forest Pathology, 2017, 47, e12353.	1.1	29
12	Growth, morphology and gas exchange responses of two-year-old <i>Quercus castaneifolia</i> seedlings to flooding stress. Scandinavian Journal of Forest Research, 2016, 31, 458-466.	1.4	6
13	Effect of SiO ₂ nanoparticles on drought resistance in hawthorn seedlings. Forest Research Papers, 2015, 76, 350-359.	0.2	88
14	Utility of ITS region sequence and structure for molecular identification of Tilia species from Hyrcanian forests, Iran. Plant Systematics and Evolution, 2012, 298, 947-961.	0.9	22
15	Long-term impact of municipal sewage irrigation on treated soil and black locust trees in a semi-arid suburban area of Iran. Journal of Environmental Sciences, 2009, 21, 1438-1445.	6.1	19
16	Influence of Two Ground-Based Skidding Systems on Soil Compaction Under Different Slope and Gradient Conditions. International Journal of Forest Engineering, 2008, 19, 9-16.	0.8	28
17	Restoration of Oak Forests in Soils Compacted by Human and Livestock. Pakistan Journal of Biological Sciences, 2007, 10, 1536-1539.	0.5	0