

# Erik H Poelman

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8753532/publications.pdf>

Version: 2024-02-01

108  
papers

4,831  
citations

87843

38  
h-index

114418

63  
g-index

115  
all docs

115  
docs citations

115  
times ranked

3947  
citing authors

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Plasticity in induced resistance to sequential attack by multiple herbivores in <i>Brassica nigra</i> . <i>Oecologia</i> , 2022, 198, 11-20.  | 0.9 | 7         |
| 2  | The Ecology of Hyperparasitoids. <i>Annual Review of Entomology</i> , 2022, 67, 143-161.  | 5.7 | 11        |
| 3  | Impact of parasitoid-associated polydnviruses on plant-mediated herbivore interactions. <i>Current Opinion in Insect Science</i> , 2022, 49, 56-62.   | 2.2 | 5         |
| 4  | Plant defense strategies against attack by multiple herbivores. <i>Trends in Plant Science</i> , 2022, 27, 528-535.   | 4.3 | 23        |
| 5  | Flowers prepare thyselfes: leaf and root herbivores induce specific changes in floral phytochemistry with consequences for plant interactions with florivores. <i>New Phytologist</i> , 2022, 233, 2548-2560. | 3.5 | 6         |
| 6  | Effects of ozone stress on flowering phenology, plant-pollinator interactions and plant reproductive success. <i>Environmental Pollution</i> , 2021, 272, 115953.   | 3.7 | 21        |
| 7  | Insect egg-killing: a new front on the evolutionary arms-race between brassicaceous plants and pierid butterflies. <i>New Phytologist</i> , 2021, 230, 341-353.   | 3.5 | 27        |
| 8  | Insect species richness affects plant responses to multi-herbivore attack. <i>New Phytologist</i> , 2021, 231, 2333-2345.   | 3.5 | 14        |
| 9  | Predictability of Biotic Stress Structures Plant Defence Evolution. <i>Trends in Ecology and Evolution</i> , 2021, 36, 444-456.   | 4.2 | 48        |
| 10 | Intraspecific variation in plant-associated herbivore communities is phylogenetically structured in Brassicaceae. <i>Ecology Letters</i> , 2021, 24, 2314-2327.   | 3.0 | 8         |
| 11 | Plant-phenotypic changes induced by parasitoid ichnoviruses enhance the performance of both unparasitized and parasitized caterpillars. <i>Molecular Ecology</i> , 2021, 30, 4567-4583.                       | 2.0 | 7         |
| 12 | The enemy of my enemy is not always my friend: Negative effects of carnivorous arthropods on plants. <i>Functional Ecology</i> , 2021, 35, 2365-2375.   | 1.7 | 10        |
| 13 | Parasitism by endoparasitoid wasps alters the internal but not the external microbiome in host caterpillars. <i>Animal Microbiome</i> , 2021, 3, 73.  | 1.5 | 12        |
| 14 | Plant defence to sequential attack is adapted to prevalent herbivores. <i>Nature Plants</i> , 2021, 7, 1347-1353.   | 4.7 | 32        |
| 15 | Plant age at the time of ozone exposure affects flowering patterns, biotic interactions and reproduction of wild mustard. <i>Scientific Reports</i> , 2021, 11, 23448.  | 1.6 | 5         |
| 16 | Microbial Symbionts of Parasitoids. <i>Annual Review of Entomology</i> , 2020, 65, 171-190.   | 5.7 | 44        |
| 17 | Exploiting chemical ecology to manage hyperparasitoids in biological control of arthropod pests. <i>Pest Management Science</i> , 2020, 76, 432-443.  | 1.7 | 39        |
| 18 | Adverse weather conditions impede odor-guided foraging of parasitoids and reduce their host-finding success. <i>Agriculture, Ecosystems and Environment</i> , 2020, 301, 107066.                              | 2.5 | 8         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | Adapted dandelions trade dispersal for germination upon root herbivore attack. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2020, 287, 20192930.   | 1.2 | 7         |
| 20 | Spatial scale, neighbouring plants and variation in plant volatiles interactively determine the strength of host-parasitoid relationships. <i>Oikos</i> , 2020, 129, 1429-1439.                              | 1.2 | 8         |
| 21 | Gone with the wind: low availability of volatile information limits foraging efficiency in downwind-flying parasitoids. <i>Animal Behaviour</i> , 2020, 165, 59-70.  | 0.8 | 6         |
| 22 | Plant ontogeny determines strength and associated plant fitness consequences of plant-mediated interactions between herbivores and flower visitors. <i>Journal of Ecology</i> , 2020, 108, 1046-1060.        | 1.9 | 15        |
| 23 | Variation in parasitoid attraction to herbivore-infested plants and alternative host plant cover mediate tritrophic interactions at the landscape scale. <i>Landscape Ecology</i> , 2020, 35, 907-919.       | 1.9 | 6         |
| 24 | Optimal plant defence under competition for light and nutrients: an evolutionary modelling approach. <i>In Silico Plants</i> , 2020, 2, .  | 0.8 | 4         |
| 25 | Intraspecific variation in herbivore-induced plant volatiles influences the spatial range of plant-parasitoid interactions. <i>Oikos</i> , 2019, 128, 77-86.   | 1.2 | 31        |
| 26 | Settling on leaves or flowers: herbivore feeding site determines the outcome of indirect interactions between herbivores and pollinators. <i>Oecologia</i> , 2019, 191, 887-896.                             | 0.9 | 10        |
| 27 | Tritrophic interactions: bridging species, communities and ecosystems. <i>Ecology Letters</i> , 2019, 22, 2151-2167.   | 3.0 | 77        |
| 28 | Leaf metabolic signatures induced by real and simulated herbivory in black mustard ( <i>Brassica nigra</i> ). <i>Metabolomics</i> , 2019, 15, 130.   | 1.4 | 29        |
| 29 | Floral plasticity: Herbivore-specific induced changes in flower traits with contrasting effects on pollinator visitation. <i>Plant, Cell and Environment</i> , 2019, 42, 1882-1896.                          | 2.8 | 55        |
| 30 | Ecological significance of light quality in optimizing plant defence. <i>Plant, Cell and Environment</i> , 2019, 42, 1065-1077.  | 2.8 | 12        |
| 31 | Cross-seasonal legacy effects of arthropod community on plant fitness in perennial plants. <i>Journal of Ecology</i> , 2019, 107, 2451-2463.   | 1.9 | 10        |
| 32 | Ecology of Plastic Flowers. <i>Trends in Plant Science</i> , 2019, 24, 725-740.  | 4.3 | 38        |
| 33 | Getting confused: learning reduces parasitoid foraging efficiency in some environments with non-host-infested plants. <i>Oecologia</i> , 2019, 189, 919-930.   | 0.9 | 12        |
| 34 | Hyperparasitoids exploit herbivore-induced plant volatiles during host location to assess host quality and non-host identity. <i>Oecologia</i> , 2019, 189, 699-709.   | 0.9 | 19        |
| 35 | Plant-mediated effects of ozone on herbivores depend on exposure duration and temperature. <i>Scientific Reports</i> , 2019, 9, 19891.   | 1.6 | 14        |
| 36 | Understanding insect foraging in complex habitats by comparing trophic levels: insights from specialist host-parasitoid-hyperparasitoid systems. <i>Current Opinion in Insect Science</i> , 2019, 32, 54-60. | 2.2 | 36        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Ecological interactions shape the adaptive value of plant defence: Herbivore attack versus competition for light. <i>Functional Ecology</i> , 2019, 33, 129-138.   | 1.7 | 28        |
| 38 | Parasitic wasp-associated symbiont affects plant-mediated species interactions between herbivores. <i>Ecology Letters</i> , 2018, 21, 957-967.   | 3.0 | 34        |
| 39 | Female response to predation risk alters conspecific male behaviour during pre-copulatory mate guarding. <i>Ethology</i> , 2018, 124, 122-130.   | 0.5 | 3         |
| 40 | Associative learning of host presence in non-host environments influences parasitoid foraging. <i>Ecological Entomology</i> , 2018, 43, 318-325.   | 1.1 | 7         |
| 41 | Elucidating the interaction between light competition and herbivore feeding patterns using functional structural plant modelling. <i>Annals of Botany</i> , 2018, 121, 1019-1031.                                | 1.4 | 27        |
| 42 | Dealing with mutualists and antagonists: Specificity of plant-mediated interactions between herbivores and flower visitors, and consequences for plant fitness. <i>Functional Ecology</i> , 2018, 32, 1022-1035. | 1.7 | 39        |
| 43 | Symbiotic polydnavirus and venom reveal parasitoid to its hyperparasitoids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2018, 115, 5205-5210.                       | 3.3 | 54        |
| 44 | Covariation and phenotypic integration in chemical communication displays: biosynthetic constraints and eco-evolutionary implications. <i>New Phytologist</i> , 2018, 220, 739-749.                              | 3.5 | 101       |
| 45 | Order of herbivore arrival on wild cabbage populations influences subsequent arthropod community development. <i>Oikos</i> , 2018, 127, 1482-1493.   | 1.2 | 30        |
| 46 | Dynamic Plant-Plant-Herbivore Interactions Govern Plant Growth-Defence Integration. <i>Trends in Plant Science</i> , 2017, 22, 329-337.  | 4.3 | 40        |
| 47 | Herbivore-induced plant volatiles and tritrophic interactions across spatial scales. <i>New Phytologist</i> , 2017, 216, 1054-1063.  | 3.5 | 147       |
| 48 | Intraspecific chemical diversity among neighbouring plants correlates positively with plant size and herbivore load but negatively with herbivore damage. <i>Ecology Letters</i> , 2017, 20, 87-97.              | 3.0 | 50        |
| 49 | Response of <i>Brassica oleracea</i> to temporal variation in attack by two herbivores affects preference and performance of a third herbivore. <i>Ecological Entomology</i> , 2017, 42, 803-815.                | 1.1 | 14        |
| 50 | When does it pay off to prime for defense? A modeling analysis. <i>New Phytologist</i> , 2017, 216, 782-797.   | 3.5 | 39        |
| 51 | Density-mediated indirect interactions alter host foraging behaviour of parasitoids without altering foraging efficiency. <i>Ecological Entomology</i> , 2016, 41, 562-571.                                      | 1.1 | 6         |
| 52 | Modification of plant-induced responses by an insect ecosystem engineer influences the colonization behaviour of subsequent shelter-users. <i>Journal of Ecology</i> , 2016, 104, 1096-1105.                     | 1.9 | 20        |
| 53 | Feeding guild of non-host community members affects host foraging efficiency of a parasitic wasp. <i>Ecology</i> , 2016, 97, 1388-1399.  | 1.5 | 20        |
| 54 | Development of a solitary koinobiont hyperparasitoid in different instars of its primary and secondary hosts. <i>Journal of Insect Physiology</i> , 2016, 90, 36-42.   | 0.9 | 5         |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Herbivore species identity rather than diversity of the non-host community determines foraging behaviour of the parasitoid wasp <i>Cotesia glomerata</i> . <i>Entomologia Experimentalis Et Applicata</i> , 2016, 161, 20-30. | 0.7 | 8         |
| 56 | Plant-mediated interactions between two herbivores differentially affect a subsequently arriving third herbivore in populations of wild cabbage. <i>Plant Biology</i> , 2016, 18, 981-991.                                    | 1.8 | 31        |
| 57 | Complexity of Plant Volatile-Mediated Interactions Beyond the Third Trophic Level. <i>Signaling and Communication in Plants</i> , 2016, , 211-225.  | 0.5 | 6         |
| 58 | Intrinsic competition between primary hyperparasitoids of the solitary endoparasitoid <i>Cotesia rubecula</i> . <i>Ecological Entomology</i> , 2016, 41, 292-300.   | 1.1 | 4         |
| 59 | Community structure and abundance of insects in response to early season aphid infestation in wild cabbage populations. <i>Ecological Entomology</i> , 2016, 41, 378-388.   | 1.1 | 15        |
| 60 | Flexible parasitoid behaviour overcomes constraint resulting from position of host and nonhost herbivores. <i>Animal Behaviour</i> , 2016, 113, 125-135.  | 0.8 | 13        |
| 61 | Keystone Herbivores and the Evolution of Plant Defenses. <i>Trends in Plant Science</i> , 2016, 21, 477-485.  | 4.3 | 83        |
| 62 | From induced resistance to defence in plant-insect interactions. <i>Entomologia Experimentalis Et Applicata</i> , 2015, 157, 11-17.   | 0.7 | 63        |
| 63 | Early herbivore alert matters: plant-mediated effects of egg deposition on higher trophic levels benefit plant fitness. <i>Ecology Letters</i> , 2015, 18, 927-936.   | 3.0 | 45        |
| 64 | Fitness consequences of indirect plant defence in the annual weed, <i>Sinapis arvensis</i> . <i>Functional Ecology</i> , 2015, 29, 1019-1025.   | 1.7 | 45        |
| 65 | Ozone affects growth and development of <i>Pieris brassicae</i> on the wild host plant <i>Brassica nigra</i> . <i>Environmental Pollution</i> , 2015, 199, 119-129.   | 3.7 | 39        |
| 66 | Altered Volatile Profile Associated with Precopulatory Mate Guarding Attracts Spider Mite Males. <i>Journal of Chemical Ecology</i> , 2015, 41, 187-193.  | 0.9 | 9         |
| 67 | Editorial overview: Ecology: Ecology of plant insect interactions: the role of plant chemistry. <i>Current Opinion in Insect Science</i> , 2015, 8, iv-vi.  | 2.2 | 0         |
| 68 | Parasitism overrides herbivore identity allowing hyperparasitoids to locate their parasitoid host using herbivore-induced plant volatiles. <i>Molecular Ecology</i> , 2015, 24, 2886-2899.                                    | 2.0 | 40        |
| 69 | Drought stress affects plant metabolites and herbivore preference but not host location by its parasitoids. <i>Oecologia</i> , 2015, 177, 701-713.  | 0.9 | 75        |
| 70 | Food plant and herbivore host species affect the outcome of intrinsic competition among parasitoid larvae. <i>Ecological Entomology</i> , 2014, 39, 693-702.  | 1.1 | 33        |
| 71 | Body Odors of Parasitized Caterpillars Give Away the Presence of Parasitoid Larvae to Their Primary Hyperparasitoid Enemies. <i>Journal of Chemical Ecology</i> , 2014, 40, 986-995.  | 0.9 | 22        |
| 72 | Caught between Parasitoids and Predators – Survival of a Specialist Herbivore on Leaves and Flowers of Mustard Plants. <i>Journal of Chemical Ecology</i> , 2014, 40, 621-631.  | 0.9 | 31        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 73 | Plant Interactions with Multiple Insect Herbivores: From Community to Genes. <i>Annual Review of Plant Biology</i> , 2014, 65, 689-713.  | 8.6 | 361       |
| 74 | Insect herbivore-associated organisms affect plant responses to herbivory. <i>New Phytologist</i> , 2014, 204, 315-321.  | 3.5 | 78        |
| 75 | Foraging behaviour by parasitoids in multiherbivore communities. <i>Animal Behaviour</i> , 2013, 85, 1517-1528.  | 0.8 | 98        |
| 76 | A test of genotypic variation in specificity of herbivore-induced responses in <i>Solidago altissima</i> L. (Asteraceae). <i>Oecologia</i> , 2013, 173, 1387-1396.   | 0.9 | 48        |
| 77 | New Synthesis: Volatiles Bring out the Animal in Plants. <i>Journal of Chemical Ecology</i> , 2013, 39, 1055-1055.   | 0.9 | 1         |
| 78 | Intrinsic Inter- and Intraspecific Competition in Parasitoid Wasps. <i>Annual Review of Entomology</i> , 2013, 58, 333-351.  | 5.7 | 247       |
| 79 | Variation in herbivore-induced plant volatiles corresponds with spatial heterogeneity in the level of parasitoid competition and parasitoid exposure to hyperparasitism. <i>Functional Ecology</i> , 2013, 27, 1107-1116.                  | 1.7 | 32        |
| 80 | Amazon poison frogs ( <i>Ranitomeya amazonica</i> ) use different phytotelm characteristics to determine their suitability for egg and tadpole deposition. <i>Evolutionary Ecology</i> , 2013, 27, 661-674.                                | 0.5 | 29        |
| 81 | Induced plant responses to microbes and insects. <i>Frontiers in Plant Science</i> , 2013, 4, 475.   | 1.7 | 42        |
| 82 | Hyperparasitoids Use Herbivore-Induced Plant Volatiles to Locate Their Parasitoid Host. <i>PLoS Biology</i> , 2012, 10, e1001435.  | 2.6 | 168       |
| 83 | Dynamics of plant secondary metabolites and consequences for food chains and community dynamics. , 2012, , 308-328.  |     | 4         |
| 84 | Plants under multiple herbivory: consequences for parasitoid search behaviour and foraging efficiency. <i>Animal Behaviour</i> , 2012, 83, 501-509.  | 0.8 | 46        |
| 85 | A taxonomic revision of the Neotropical poison frog genus <i>Ranitomeya</i> (Amphibia: Dendrobatidae). <i>Zootaxa</i> , 2011, 3083, 1.   | 0.2 | 106       |
| 86 | Indirect plant-mediated interactions among parasitoid larvae. <i>Ecology Letters</i> , 2011, 14, 670-676.  | 3.0 | 38        |
| 87 | Relative importance of plant-mediated bottom-up and top-down forces on herbivore abundance on <i>Brassica oleracea</i> . <i>Functional Ecology</i> , 2011, 25, 1113-1124.  | 1.7 | 51        |
| 88 | Parasitoid-specific induction of plant responses to parasitized herbivores affects colonization by subsequent herbivores. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011, 108, 19647-19652. | 3.3 | 82        |
| 89 | Inhibition of lipoxygenase affects induction of both direct and indirect plant defences against herbivorous insects. <i>Oecologia</i> , 2010, 162, 393-404.  | 0.9 | 64        |
| 90 | The Herbivore-Induced Plant Volatile Methyl Salicylate Negatively Affects Attraction of the Parasitoid <i>Diadegma semiclausum</i> . <i>Journal of Chemical Ecology</i> , 2010, 36, 479-489.   | 0.9 | 77        |

| #   | ARTICLE   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 91  | Development and host utilization in <i>Hyposoter ebeninus</i> (Hymenoptera: Ichneumonidae), a solitary endoparasitoid of <i>Pieris rapae</i> and <i>P. brassicae</i> caterpillars (Lepidoptera: Pieridae). <i>Biological Control</i> , 2010, 53, 312-318. | 1.4 | 24        |
| 92  | Intraspecific variation in herbivore community composition and transcriptional profiles in field-grown <i>Brassica oleracea</i> cultivars. <i>Journal of Experimental Botany</i> , 2010, 61, 807-819.   | 2.4 | 29        |
| 93  | Descriptions of the Tadpoles of Two Poison Frogs, <i>Ameerega parvula</i> and <i>Ameerega bilineata</i> (Anura: Tj ETQq1 1 0.784314 rgBT /Ove 0.2 8   | 0.2 | 8         |
| 94  | Herbivore-induced plant responses in <i>Brassica oleracea</i> prevail over effects of constitutive resistance and result in enhanced herbivore attack. <i>Ecological Entomology</i> , 2010, 35, 240-247.  | 1.1 | 91        |
| 95  | Consequences of constitutive and induced variation in plant nutritional quality for immune defence of a herbivore against parasitism. <i>Oecologia</i> , 2009, 160, 299-308.  | 0.9 | 106       |
| 96  | Field parasitism rates of caterpillars on <i>Brassica oleracea</i> plants are reliably predicted by differential attraction of <i>Cotesia</i> parasitoids. <i>Functional Ecology</i> , 2009, 23, 951-962.   | 1.7 | 87        |
| 97  | Chemical diversity in <i>Brassica oleracea</i> affects biodiversity of insect herbivores. <i>Ecology</i> , 2009, 90, 1863-1877.   | 1.5 | 120       |
| 98  | Performance of specialist and generalist herbivores feeding on cabbage cultivars is not explained by glucosinolate profiles. <i>Entomologia Experimentalis Et Applicata</i> , 2008, 127, 218-228.   | 0.7 | 103       |
| 99  | Early season herbivore differentially affects plant defence responses to subsequently colonizing herbivores and their abundance in the field. <i>Molecular Ecology</i> , 2008, 17, 3352-3365.   | 2.0 | 214       |
| 100 | Responses of <i>Brassica oleracea</i> cultivars to infestation by the aphid <i>Brevicoryne brassicae</i> : an ecological and molecular approach. <i>Plant, Cell and Environment</i> , 2008, 31, 1592-1605.  | 2.8 | 63        |
| 101 | Consequences of variation in plant defense for biodiversity at higher trophic levels. <i>Trends in Plant Science</i> , 2008, 13, 534-541.   | 4.3 | 160       |
| 102 | Space Use of Amazonian Poison Frogs: Testing the Reproductive Resource Defense Hypothesis. <i>Journal of Herpetology</i> , 2008, 42, 270-278.   | 0.2 | 27        |
| 103 | Genotypic variation in genome-wide transcription profiles induced by insect feeding: <i>Brassica oleracea</i> and <i>Pieris rapae</i> interactions. <i>BMC Genomics</i> , 2007, 8, 239.   | 1.2 | 75        |
| 104 | Offering offspring as food to cannibals: oviposition strategies of Amazonian poison frogs ( <i>Dendrobates ventrimaculatus</i> ). <i>Evolutionary Ecology</i> , 2007, 21, 215-227.  | 0.5 | 70        |
| 105 | A new species of <i>Colostethus</i> (Anura, Dendrobatidae) from French Guiana with a redescription of <i>Colostethus beebei</i> (Noble, 1923) from its type locality. <i>Phyllomedusa</i> , 2006, 5, 43.  | 0.2 | 43        |
| 106 | Consistent feeding positions of great tit parents. <i>Animal Behaviour</i> , 2006, 72, 1249-1257.   | 0.8 | 21        |
| 107 | <i>Dicyphus</i> predatory bugs pre-established on tomato plants reduce <i>Nesidiocoris tenuis</i> population growth. <i>Journal of Pest Science</i> , 0, , 1.   | 1.9 | 5         |
| 108 | Evolution of koinobiont parasitoid host regulation and consequences for indirect plant defence. <i>Evolutionary Ecology</i> , 0, , 1.   | 0.5 | 7         |