

# Bettina Hjelm Clausen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8761483/publications.pdf>

Version: 2024-02-01

19  
papers

1,261  
citations

840585

11  
h-index

887953

17  
g-index

20  
all docs

20  
docs citations

20  
times ranked

2088  
citing authors

#	ARTICLE	IF	CITATIONS
1	The Role of Tumor Necrosis Factor Following Spinal Cord Injury: A Systematic Review. <i>Cellular and Molecular Neurobiology</i> , 2023, 43, 925-950.	1.7	6
2	Interleukin-1 Mediates Ischemic Brain Injury via Induction of IL-17A in $\gamma\delta$ T Cells and CXCL1 in Astrocytes. <i>NeuroMolecular Medicine</i> , 2022, 24, 437-451.	1.8	9
3	The Inflammatory Response after Moderate Contusion Spinal Cord Injury: A Time Study. <i>Biology</i> , 2022, 11, 939.	1.3	5
4	Bone Marrow-Derived IL-1Ra Increases TNF Levels Poststroke. <i>Cells</i> , 2021, 10, 956.	1.8	3
5	Topical Administration of a Soluble TNF Inhibitor Reduces Infarct Volume After Focal Cerebral Ischemia in Mice. <i>Frontiers in Neuroscience</i> , 2019, 13, 781.	1.4	25
6	Post-stroke inflammation—target or tool for therapy?. <i>Acta Neuropathologica</i> , 2019, 137, 693-714.	3.9	286
7	Inflammation leads to distinct populations of extracellular vesicles from microglia. <i>Journal of Neuroinflammation</i> , 2018, 15, 168.	3.1	133
8	Spontaneous ischaemic stroke lesions in a dog brain: neuropathological characterisation and comparison to human ischaemic stroke. <i>Acta Veterinaria Scandinavica</i> , 2017, 59, 7.	0.5	10
9	Beneficial potential of intravenously administered IL-6 in improving outcome after murine experimental stroke. <i>Brain, Behavior, and Immunity</i> , 2017, 65, 296-311.	2.0	36
10	Fumarate decreases edema volume and improves functional outcome after experimental stroke. <i>Experimental Neurology</i> , 2017, 295, 144-154.	2.0	42
11	Genetic Ablation of Soluble TNF Does Not Affect Lesion Size and Functional Recovery after Moderate Spinal Cord Injury in Mice. <i>Mediators of Inflammation</i> , 2016, 2016, 1-15.	1.4	12
12	Conditional ablation of myeloid TNF increases lesion volume after experimental stroke in mice, possibly via altered ERK1/2 signaling. <i>Scientific Reports</i> , 2016, 6, 29291.	1.6	37
13	Cell therapy centered on IL-1Ra is neuroprotective in experimental stroke. <i>Acta Neuropathologica</i> , 2016, 131, 775-791.	3.9	68
14	Systemically administered anti-TNF therapy ameliorates functional outcomes after focal cerebral ischemia. <i>Journal of Neuroinflammation</i> , 2014, 11, 203.	3.1	79
15	CSF transthyretin neuroprotection in a mouse model of brain ischemia. <i>Journal of Neurochemistry</i> , 2010, 115, 1434-1444.	2.1	73
16	Microglia Protect Neurons against Ischemia by Synthesis of Tumor Necrosis Factor. <i>Journal of Neuroscience</i> , 2009, 29, 1319-1330.	1.7	371
17	A Role for Interferon-Gamma in Focal Cerebral Ischemia in Mice. <i>Journal of Neuropathology and Experimental Neurology</i> , 2004, 63, 942-955.	0.9	65
18	Distal middle cerebral artery occlusion does not result in depression-like behaviours. <i>F1000Research</i> , 0, 7, 1430.	0.8	0

#	ARTICLE	IF	CITATIONS
19	An exploratory investigation of "depression-like" behaviours in a model of left-sided distal middle cerebral artery occlusion in young, male C57B6 mice. F1000Research, 0, 7, 1430.	0.8	1