

Joris A M Van Der Post

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8765769/joris-a-m-van-der-post-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

203
papers

7,318
citations

44
h-index

79
g-index

218
ext. papers

8,537
ext. citations

4.7
avg, IF

5.34
L-index

#	Paper	IF	Citations
203	Induction of labour versus expectant monitoring for gestational hypertension or mild pre-eclampsia after 36 weeks' gestation (HYPITAT): a multicentre, open-label randomised controlled trial. <i>Lancet, The</i> , 2009 , 374, 979-988	4.0	509
202	Use of uterine artery Doppler ultrasonography to predict pre-eclampsia and intrauterine growth restriction: a systematic review and bivariable meta-analysis. <i>Cmaj</i> , 2008 , 178, 701-11	3.5	480
201	Aspirin plus heparin or aspirin alone in women with recurrent miscarriage. <i>New England Journal of Medicine</i> , 2010 , 362, 1586-96	59.2	356
200	Induction versus expectant monitoring for intrauterine growth restriction at term: randomised equivalence trial (DIGITAT). <i>BMJ, The</i> , 2010 , 341, c7087	5.9	302
199	Significance of (sub)clinical thyroid dysfunction and thyroid autoimmunity before conception and in early pregnancy: a systematic review. <i>Human Reproduction Update</i> , 2011 , 17, 605-19	15.8	298
198	Transgenerational effects of prenatal exposure to the 1944-45 Dutch famine. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2013 , 120, 548-53	3.7	296
197	Accuracy of circulating placental growth factor, vascular endothelial growth factor, soluble fms-like tyrosine kinase 1 and soluble endoglin in the prediction of pre-eclampsia: a systematic review and meta-analysis. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012 , 119, 778-87	3.7	185
196	Cognitive, motor, behavioural and academic performances of children born preterm: a meta-analysis and systematic review involving 64 061 children. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2018 , 125, 16-25	3.7	161
195	Microparticle subpopulations are increased in preeclampsia: possible involvement in vascular dysfunction?. <i>American Journal of Obstetrics and Gynecology</i> , 2002 , 187, 450-6	6.4	154
194	Accuracy of mean arterial pressure and blood pressure measurements in predicting pre-eclampsia: systematic review and meta-analysis. <i>BMJ, The</i> , 2008 , 336, 1117-20	5.9	148
193	Foley catheter versus vaginal prostaglandin E2 gel for induction of labour at term (PROBAAT trial): an open-label, randomised controlled trial. <i>Lancet, The</i> , 2011 , 378, 2095-103	4.0	131
192	Consequences of hyperemesis gravidarum for offspring: a systematic review and meta-analysis. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2011 , 118, 1302-13	3.7	131
191	Neonatal outcome following elective cesarean section beyond 37 weeks of gestation: a 7-year retrospective analysis of a national registry. <i>American Journal of Obstetrics and Gynecology</i> , 2010 , 202, 250.e1-8	6.4	115
190	Extensive platelet activation in preeclampsia compared with normal pregnancy: enhanced expression of cell adhesion molecules. <i>American Journal of Obstetrics and Gynecology</i> , 1997 , 176, 461-9	6.4	111
189	Outcomes after internal versus external tocodynamometry for monitoring labor. <i>New England Journal of Medicine</i> , 2010 , 362, 306-13	59.2	99
188	External cephalic version-related risks: a meta-analysis. <i>Obstetrics and Gynecology</i> , 2008 , 112, 1143-51	4.9	95
187	Changes in microparticle numbers and cellular origin during pregnancy and preeclampsia. <i>Hypertension in Pregnancy</i> , 2008 , 27, 344-60	2	93

186	Obstetric complications in Marfan syndrome. <i>International Journal of Cardiology</i> , 2006 , 110, 53-9	3.2	92
185	Predictive value of cervical length measurement and fibronectin testing in threatened preterm labor. <i>Obstetrics and Gynecology</i> , 2014 , 123, 1185-1192	4.9	80
184	Serum screening with Down's syndrome markers to predict pre-eclampsia and small for gestational age: systematic review and meta-analysis. <i>BMC Pregnancy and Childbirth</i> , 2008 , 8, 33	3.2	77
183	Maternal and perinatal outcome after expectant management of the HELLP syndrome compared with pre-eclampsia without HELLP syndrome. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 1998 , 76, 31-6	2.4	72
182	Term breech deliveries in the Netherlands: did the increased cesarean rate affect neonatal outcome? A population-based cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2014 , 93, 888-96	3.8	69
181	Molecular genetics of preeclampsia and HELLP syndrome - a review. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2012 , 1822, 1960-9	6.9	69
180	The Bishop score as a predictor of labor induction success: a systematic review. <i>American Journal of Perinatology</i> , 2013 , 30, 625-30	3.3	64
179	Differentially expressed genes in the pre-eclamptic placenta: a systematic review and meta-analysis. <i>PLoS ONE</i> , 2013 , 8, e68991	3.7	61
178	Clinical factors to predict the outcome of external cephalic version: a metaanalysis. <i>American Journal of Obstetrics and Gynecology</i> , 2008 , 199, 630.e1-7; discussion e1-5	6.4	60
177	Are tests for predicting pre-eclampsia good enough to make screening viable? A review of reviews and critical appraisal. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2009 , 88, 758-65	3.8	57
176	Expression of placental FLT1 transcript variants relates to both gestational hypertensive disease and fetal growth. <i>Hypertension</i> , 2011 , 58, 70-6	8.5	56
175	Transfusion policy after severe postpartum haemorrhage: a randomised non-inferiority trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2014 , 121, 1005-14	3.7	55
174	Effect of maintenance tocolysis with nifedipine in threatened preterm labor on perinatal outcomes: a randomized controlled trial. <i>JAMA - Journal of the American Medical Association</i> , 2013 , 309, 41-7	27.4	55
173	Leukocyte activation and circulating leukocyte-derived microparticles in preeclampsia. <i>American Journal of Reproductive Immunology</i> , 2009 , 61, 346-59	3.8	55
172	Enhanced Coagulation Activation in Preeclampsia: the Role of APC Resistance, Microparticles and Other Plasma Constituents. <i>Thrombosis and Haemostasis</i> , 2002 , 88, 415-420	7	53
171	Neonatal morbidity after induction vs expectant monitoring in intrauterine growth restriction at term: a subanalysis of the DIGITAT RCT. <i>American Journal of Obstetrics and Gynecology</i> , 2012 , 206, 344.e1-7	6.4	51
170	Impact of fetal gender on the risk of preterm birth, a national cohort study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2016 , 95, 1034-41	3.8	51
169	Microparticles and exosomes: impact on normal and complicated pregnancy. <i>American Journal of Reproductive Immunology</i> , 2007 , 58, 389-402	3.8	50

168	Nifedipine versus atosiban for threatened preterm birth (APOSTEL III): a multicentre, randomised controlled trial. <i>Lancet, The</i> , 2016 , 387, 2117-2124	4.0	49
167	Maternal prepregnancy BMI, offspring's early postnatal growth, and metabolic profile at age 5-6 years: the ABCD Study. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 3845-54	5.6	49
166	Risk factors for postpartum urinary retention: a systematic review and meta-analysis. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012 , 119, 1440-6	3.7	49
165	Cardiovascular risk estimation in women with a history of hypertensive pregnancy disorders at term: a longitudinal follow-up study. <i>BMC Pregnancy and Childbirth</i> , 2013 , 13, 126	3.2	48
164	Effects on (neuro)developmental and behavioral outcome at 2 years of age of induced labor compared with expectant management in intrauterine growth-restricted infants: long-term outcomes of the DIGITAT trial. <i>American Journal of Obstetrics and Gynecology</i> , 2012 , 206, 406.e1-7	6.4	48
163	Ethnic differences in stillbirth and early neonatal mortality in The Netherlands. <i>Journal of Epidemiology and Community Health</i> , 2011 , 65, 696-701	5.1	48
162	Can flow cytometric detection of platelet activation early in pregnancy predict the occurrence of preeclampsia? A prospective study. <i>American Journal of Obstetrics and Gynecology</i> , 1997 , 177, 434-42	6.4	47
161	Subfertility and assisted reproduction techniques are associated with poorer cardiometabolic profiles in childhood. <i>Reproductive BioMedicine Online</i> , 2015 , 30, 258-67	4	44
160	Seven placental transcripts characterize HELLP-syndrome. <i>Placenta</i> , 2008 , 29, 444-53	3.4	44
159	Transvaginal sonographic assessment of cervical length and wedging for predicting outcome of labor induction at term: a systematic review and meta-analysis. <i>Ultrasound in Obstetrics and Gynecology</i> , 2013 , 42, 500-8	5.8	42
158	Accuracy of serum uric acid determination in predicting pre-eclampsia: a systematic review. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2006 , 85, 519-25	3.8	42
157	Delivery-related risk factors for covert postpartum urinary retention after vaginal delivery. <i>International Urogynecology Journal</i> , 2016 , 27, 55-60	2	39
156	STOX1 is not imprinted and is not likely to be involved in preeclampsia. <i>Nature Genetics</i> , 2007 , 39, 279-80; author reply 280-1	36.3	39
155	Accuracy of body mass index in predicting pre-eclampsia: bivariate meta-analysis. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2007 , 114, 1477-85	3.7	39
154	Maternal prepregnancy body mass index and their children's blood pressure and resting cardiac autonomic balance at age 5 to 6 years. <i>Hypertension</i> , 2013 , 62, 641-7	8.5	38
153	The functions of microparticles in pre-eclampsia. <i>Seminars in Thrombosis and Hemostasis</i> , 2011 , 37, 146-53	5.3	38
152	Foley catheter or prostaglandin E2 inserts for induction of labour at term: an open-label randomized controlled trial (PROBAAT-P trial) and systematic review of literature. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2013 , 170, 137-45	2.4	37
151	High activity of fatty acid oxidation enzymes in human placenta: implications for fetal-maternal disease. <i>Journal of Inherited Metabolic Disease</i> , 2003 , 26, 385-92	5.4	37

150	Prediction of success of external cephalic version after 36 weeks. <i>American Journal of Perinatology</i> , 2011 , 28, 103-10	3.3	36
149	Postpartum urinary retention: a systematic review of adverse effects and management. <i>International Urogynecology Journal</i> , 2014 , 25, 1605-12	2	34
148	Ted (G.J.) Kloosterman: on intrauterine growth. The significance of prenatal care. Studies on birth weight, placental weight and placental index. <i>Placenta</i> , 2006 , 27, 1052-4	3.4	34
147	Validation of automated blood pressure recording in pregnancy. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1994 , 101, 66-9	3.7	34
146	SFlt-1 elevates blood pressure by augmenting endothelin-1-mediated vasoconstriction in mice. <i>PLoS ONE</i> , 2014 , 9, e91897	3.7	33
145	An economic analysis of induction of labour and expectant monitoring in women with gestational hypertension or pre-eclampsia at term (HYPITAT trial). <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2010 , 117, 1577-85	3.7	31
144	Pregnancy in women suffering from familial hypercholesterolemia: a harmful period for both mother and newborn?. <i>Current Opinion in Lipidology</i> , 2009 , 20, 484-90	4.4	30
143	Circulating platelet-derived and placenta-derived microparticles expose Flt-1 in preeclampsia. <i>Reproductive Sciences</i> , 2008 , 15, 1002-10	3	30
142	Quantitative fetal fibronectin testing in combination with cervical length measurement in the prediction of spontaneous preterm delivery in symptomatic women. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1965-1971	3.7	30
141	The idiopathic preterm delivery methylation profile in umbilical cord blood DNA. <i>BMC Genomics</i> , 2015 , 16, 736	4.5	29
140	Should cervical favourability play a role in the decision for labour induction in gestational hypertension or mild pre-eclampsia at term? An exploratory analysis of the HYPITAT trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2012 , 119, 1123-30	3.7	29
139	Vasopressin and oxytocin levels during normal pregnancy: effects of chronic dietary sodium restriction. <i>Journal of Endocrinology</i> , 1997 , 152, 345-54	4.7	29
138	Induction of labour at 41 weeks versus expectant management until 42 weeks (INDEX): multicentre, randomised non-inferiority trial. <i>BMJ, The</i> , 2019 , 364, l344	5.9	29
137	Effect of levothyroxine on live birth rate in euthyroid women with recurrent miscarriage and TPO antibodies (T4-LIFE study). <i>Contemporary Clinical Trials</i> , 2015 , 44, 134-138	2.3	28
136	Cost-effectiveness of induction of labour at term with a Foley catheter compared to vaginal prostaglandin E ₂ gel (PROBAAT trial). <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2013 , 120, 987-95	3.7	27
135	Cell-derived microparticles and complement activation in preeclampsia versus normal pregnancy. <i>Placenta</i> , 2007 , 28, 928-35	3.4	27
134	Microparticles and exosomes in gynecologic neoplasias. <i>Seminars in Thrombosis and Hemostasis</i> , 2010 , 36, 925-9	5.3	26
133	Disproportionate Intrauterine Growth Intervention Trial At Term: DIGITAT. <i>BMC Pregnancy and Childbirth</i> , 2007 , 7, 12	3.2	25

132	Genetic Analyses in Small-for-Gestational-Age Newborns. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018 , 103, 917-925	5.6	24
131	Labour pain with remifentanyl patient-controlled analgesia versus epidural analgesia: a randomised equivalence trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2017 , 124, 652-660	3.7	24
130	Meta-Analysis of Placental Transcriptome Data Identifies a Novel Molecular Pathway Related to Preeclampsia. <i>PLoS ONE</i> , 2015 , 10, e0132468	3.7	24
129	Induction of labour versus expectant monitoring in women with pregnancy induced hypertension or mild preeclampsia at term: the HYPITAT trial. <i>BMC Pregnancy and Childbirth</i> , 2007 , 7, 14	3.2	22
128	Expression of inflammation-related genes in endothelial cells is not directly affected by microparticles from preeclamptic patients. <i>Translational Research</i> , 2006 , 147, 310-20		22
127	Nifedipine maintenance tocolysis and perinatal outcome: an individual participant data meta-analysis. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1753-60	3.7	21
126	Value of adding second-trimester uterine artery Doppler to patient characteristics in identification of nulliparous women at increased risk for pre-eclampsia: an individual patient data meta-analysis. <i>Ultrasound in Obstetrics and Gynecology</i> , 2013 , 42, 257-67	5.8	21
125	Recurrence risk of a delivery before 34 weeks of pregnancy due to an early onset hypertensive disorder: a systematic review. <i>American Journal of Perinatology</i> , 2010 , 27, 565-71	3.3	21
124	The fetal origins of hypertension: a systematic review and meta-analysis of the evidence from animal experiments of maternal undernutrition. <i>Journal of Hypertension</i> , 2012 , 30, 2255-67	1.9	21
123	Comparison of an Auscultatory Versus an Oscillometric Ambulatory Blood Pressure Monitor in Normotensive, Hypertensive, and Preeclamptic Pregnancy. <i>Hypertension in Pregnancy</i> , 1997 , 16, 187-202 ²		21
122	Preterm birth in singleton and multiple pregnancies: evaluation of costs and perinatal outcomes. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2015 , 186, 34-41	2.4	20
121	Recurrence risk and prediction of a delivery under 34 weeks of gestation after a history of a severe hypertensive disorder. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2011 , 118, 589-95	3.7	20
120	Incidence of postpartum haemorrhage in women receiving therapeutic doses of low-molecular-weight heparin: results of a retrospective cohort study. <i>BMJ Open</i> , 2011 , 1, e000257	3	20
119	The Association Between Learning Climate and Adverse Obstetrical Outcomes in 16 Nontertiary Obstetrics-Gynecology Departments in the Netherlands. <i>Academic Medicine</i> , 2017 , 92, 1740-1748	3.9	19
118	Increased glucocerebrosidase expression and activity in preeclamptic placenta. <i>Placenta</i> , 2015 , 36, 160-9	3.4	19
117	Nifedipine as a uterine relaxant for external cephalic version: a randomized controlled trial. <i>Obstetrics and Gynecology</i> , 2008 , 112, 271-6	4.9	19
116	Circadian variation of urinary albumin excretion in pregnancy. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 1995 , 102, 107-10	3.7	19
115	Prediction models for successful external cephalic version: a systematic review. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2015 , 195, 160-7	2.4	18

114	Time to conception and time to live birth in women with unexplained recurrent miscarriage. <i>Human Reproduction</i> , 2014 , 29, 1146-52	5.7	18
113	Cost-effectiveness of fibronectin testing in a triage in women with threatened preterm labor: alleviation of pregnancy outcome by suspending tocolysis in early labor (APOSTEL-I trial). <i>BMC Pregnancy and Childbirth</i> , 2009 , 9, 38	3.2	18
112	Well being of obstetric patients on minimal blood transfusions (WOMB trial). <i>BMC Pregnancy and Childbirth</i> , 2010 , 10, 83	3.2	18
111	Prediction of pre-eclampsia: a protocol for systematic reviews of test accuracy. <i>BMC Pregnancy and Childbirth</i> , 2006 , 6, 29	3.2	18
110	Ductus venosus pulsatility index measurement reduces the false-positive rate in first-trimester screening. <i>Ultrasound in Obstetrics and Gynecology</i> , 2010 , 36, 661-7	5.8	17
109	Accuracy of fibronectin tests for the prediction of pre-eclampsia: a systematic review. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2007 , 133, 12-9	2.4	17
108	Cost-effectiveness of diagnostic testing strategies including cervical-length measurement and fibronectin testing in women with symptoms of preterm labor. <i>Ultrasound in Obstetrics and Gynecology</i> , 2018 , 51, 596-603	5.8	16
107	A model for preconceptional prediction of recurrent early-onset preeclampsia: derivation and internal validation. <i>Reproductive Sciences</i> , 2011 , 18, 1154-9	3	16
106	Comparison of Portapres with standard sphygmomanometry in pregnancy. <i>Hypertension in Pregnancy</i> , 2002 , 21, 65-76	2	16
105	Total bile acids in the maternal and fetal compartment in relation to placental ABCG2 expression in preeclamptic pregnancies complicated by HELLP syndrome. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2015 , 1852, 131-6	6.9	15
104	Early enteral tube feeding in optimizing treatment of hyperemesis gravidarum: the Maternal and Offspring outcomes after Treatment of HyperEmesis by Refeeding (MOTHER) randomized controlled trial. <i>American Journal of Clinical Nutrition</i> , 2017 , 106, 812-820	7	15
103	Intrapartum and neonatal mortality in primary midwife-led and secondary obstetrician-led care in the Amsterdam region of the Netherlands: A retrospective cohort study. <i>Midwifery</i> , 2015 , 31, 1168-76	2.8	15
102	Health-related quality of life after induction of labor versus expectant monitoring in gestational hypertension or preeclampsia at term. <i>Hypertension in Pregnancy</i> , 2011 , 30, 260-74	2	15
101	Early nasogastric tube feeding in optimising treatment for hyperemesis gravidarum: the MOTHER randomised controlled trial (Maternal and Offspring outcomes after Treatment of HyperEmesis by Refeeding). <i>BMC Pregnancy and Childbirth</i> , 2016 , 16, 22	3.2	15
100	Predicting seizures in pregnant women with epilepsy: Development and external validation of a prognostic model. <i>PLoS Medicine</i> , 2019 , 16, e1002802	11.6	14
99	Weight loss in pregnancy and cardiometabolic profile in childhood: findings from a longitudinal birth cohort. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2015 , 122, 1664-73	3.7	14
98	Morning versus evening induction of labour for improving outcomes. <i>The Cochrane Library</i> , 2013 , CD007307	3.07	14
97	Maternal health-related quality of life after induction of labor or expectant monitoring in pregnancy complicated by intrauterine growth retardation beyond 36 weeks. <i>Quality of Life Research</i> , 2011 , 20, 1427-36	3.7	14

96	Pessaries in multiple pregnancy as a prevention of preterm birth: the ProTwin Trial. <i>BMC Pregnancy and Childbirth</i> , 2009 , 9, 44	3.2	14
95	Expectant parents' preferences for mode of delivery and trade-offs of outcomes for breech presentation. <i>Patient Education and Counseling</i> , 2008 , 72, 305-10	3.1	14
94	Risk stratification with cervical length and fetal fibronectin in women with threatened preterm labor before 34 weeks and not delivering within 7 days. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2015 , 94, 715-721	3.8	13
93	Economic analysis comparing induction of labour and expectant management for intrauterine growth restriction at term (DIGITAT trial). <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2013 , 170, 358-63	2.4	13
92	Patient's willingness to opt for external cephalic version. <i>Journal of Psychosomatic Obstetrics and Gynaecology</i> , 2013 , 34, 15-21	3.6	13
91	Labour and neonatal outcome in small for gestational age babies delivered beyond 36+0 weeks: a retrospective cohort study. <i>Journal of Pregnancy</i> , 2011 , 2011, 293516	2.5	13
90	Barriers and Facilitators for the Use of a Medical Mobile App to Prevent Work-Related Risks in Pregnancy: A Qualitative Analysis. <i>JMIR Research Protocols</i> , 2017 , 6, e163	2	13
89	The impact of fetal gender and ethnicity on the risk of spontaneous preterm delivery in women with symptoms of preterm labor. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 3563-9	2	13
88	Low dose aspirin in the prevention of recurrent spontaneous preterm labour - the APRIL study: a multicenter randomized placebo controlled trial. <i>BMC Pregnancy and Childbirth</i> , 2017 , 17, 223	3.2	12
87	Microparticles of pregnant women and preeclamptic patients activate endothelial cells in the presence of monocytes. <i>American Journal of Reproductive Immunology</i> , 2012 , 67, 206-15	3.8	12
86	External validation of a prediction model for successful external cephalic version. <i>American Journal of Perinatology</i> , 2012 , 29, 231-6	3.3	12
85	Implementation of the external cephalic version in breech delivery. Dutch national implementation study of external cephalic version. <i>BMC Pregnancy and Childbirth</i> , 2010 , 10, 20	3.2	12
84	Preeclampsia is not associated with altered platelet vasopressin binding and cytosolic Ca ⁺⁺ concentration. <i>American Journal of Obstetrics and Gynecology</i> , 1993 , 169, 1169-78	6.4	12
83	Comparison of the Actim Partus test and the fetal fibronectin test in the prediction of spontaneous preterm birth in symptomatic women undergoing cervical length measurement. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2016 , 206, 220-224	2.4	12
82	Timing induction of labour at 41 or 42 weeks? A closer look at time frames of comparison: A review. <i>Midwifery</i> , 2018 , 66, 111-118	2.8	11
81	Antenatal prediction of neonatal mortality in very premature infants. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2014 , 176, 126-31	2.4	11
80	Why were the results of randomized trials on the clinical utility of fetal fibronectin negative? A systematic review of their study designs. <i>American Journal of Perinatology</i> , 2011 , 28, 145-50	3.3	11
79	Which factors play a role in clinical decision-making in external cephalic version?. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2008 , 87, 31-5	3.8	11

78	Long-term micturition problems of asymptomatic postpartum urinary retention: a prospective case-control study. <i>International Urogynecology Journal</i> , 2018 , 29, 481-488	2	10
77	Comparison of clean intermittent and transurethral indwelling catheterization for the treatment of overt urinary retention after vaginal delivery: a multicentre randomized controlled clinical trial. <i>International Urogynecology Journal</i> , 2018 , 29, 1281-1287	2	10
76	Preterm Breech Presentation: A Comparison of Intended Vaginal and Intended Cesarean Delivery. <i>Obstetrics and Gynecology</i> , 2015 , 126, 1223-1230	4.9	10
75	Which intrauterine growth restricted fetuses at term benefit from early labour induction? A secondary analysis of the DIGITAT randomised trial. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2014 , 172, 20-5	2.4	10
74	Start of induction of labour with oxytocin in the morning or in the evening. A randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2009 , 116, 562-8	3.7	10
73	Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials. <i>PLoS Medicine</i> , 2020 , 17, e1003436	11.6	10
72	Postnatal Catch-Up Growth After Suspected Fetal Growth Restriction at Term. <i>Frontiers in Endocrinology</i> , 2019 , 10, 274	5.7	9
71	Randomized comparison of nifedipine and placebo in fibronectin-negative women with symptoms of preterm labor and a short cervix (APOSTEL-I Trial). <i>American Journal of Perinatology</i> , 2015 , 32, 451-60 ^{3.3}	3.3	9
70	Live-birth rate in euthyroid women with recurrent miscarriage and thyroid peroxidase antibodies. <i>Gynecological Endocrinology</i> , 2016 , 32, 132-5	2.4	8
69	Intrapartum epidural analgesia and low Apgar score among singleton infants born at term: A propensity score matched study. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2020 , 99, 1155-1162	3.8	8
68	Assessment of perinatal outcome after sustained tocolysis in early labour (APOSTEL-II trial). <i>BMC Pregnancy and Childbirth</i> , 2009 , 9, 42	3.2	8
67	Maintenance tocolysis with nifedipine in threatened preterm labour: 2-year follow up of the offspring in the APOSTEL II trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016 , 123, 1107-14	3.7	7
66	Induction of labour or expectant monitoring in hypertensive pregnancy disorders at term: do women's postpartum cardiovascular risk factors differ between the two strategies?. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2013 , 171, 30-4	2.4	7
65	Predictive Factors for Delivery within 7 Days after Successful 48-Hour Treatment of Threatened Preterm Labor. <i>AJP Reports</i> , 2015 , 5, e141-9	1.2	7
64	World Health Organization systematic review of screening tests for preeclampsia. <i>Obstetrics and Gynecology</i> , 2005 , 105, 1151-2; author reply 1152	4.9	7
63	Higher decidual EB13 and HLA-G mRNA expression in preeclampsia: Cause or consequence of preeclampsia. <i>Human Immunology</i> , 2016 , 77, 68-70	2.3	6
62	Placental expression of heparan sulfate 3-O-sulfotransferase-3A1 in normotensive and pre-eclamptic pregnancies. <i>Placenta</i> , 2015 , 36, 1218-24	3.4	6
61	Impact of a randomized trial on maintenance tocolysis on length of hospital admission of women with threatened preterm labor in The Netherlands. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2015 , 186, 8-11	2.4	6

60	Neonatal outcome following elective cesarean section of twin pregnancies beyond 35 weeks of gestation. <i>American Journal of Obstetrics and Gynecology</i> , 2012 , 207, 480.e1-7	6.4	6
59	Subsequent Pregnancy after Preterm Prelabor Rupture of Membranes before 27 Weeks' Gestation. <i>AJP Reports</i> , 2013 , 3, 113-8	1.2	6
58	The functions of microparticles in preeclampsia. <i>Pregnancy Hypertension</i> , 2011 , 1, 59-65	2.6	6
57	Indicators for intervention during the expulsive second-stage arrest of labour. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2009 , 116, 1773-81	3.7	6
56	Internal versus external tocodynamometry during induced or augmented labour. <i>Cochrane Database of Systematic Reviews</i> , 2012 , 12, CD006947		6
55	Does use of an intrauterine catheter during labor increase risk of infection?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2012 , 25, 415-8	2	6
54	Cost-effectiveness of recurrence risk guided care versus care as usual in women who suffered from early-onset preeclampsia including HELLP syndrome in their previous pregnancy (the PreCare study). <i>BMC Pregnancy and Childbirth</i> , 2010 , 10, 60	3.2	6
53	Usability and Usefulness of a Mobile Health App for Pregnancy-Related Work Advice: Mixed-Methods Approach. <i>JMIR MHealth and UHealth</i> , 2019 , 7, e11442	5.5	6
52	Development and internal validation of a clinical prediction model for external cephalic version. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2018 , 228, 137-142	2.4	6
51	Atosiban versus fenoterol as a uterine relaxant for external cephalic version: randomised controlled trial. <i>BMJ, The</i> , 2017 , 356, i6773	5.9	5
50	The fourth sound of Korotkoff in pregnancy: a myth. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 1998 , 76, 53-9	2.4	5
49	Levothyroxine in euthyroid thyroid peroxidase antibody positive women with recurrent pregnancy loss (T4LIFE trial): a multicentre, randomised, double-blind, placebo-controlled, phase 3 trial.. <i>Lancet Diabetes and Endocrinology,the</i> , 2022 ,	18.1	5
48	The standardized 12-lead fetal electrocardiogram of the healthy fetus in mid-pregnancy: A cross-sectional study. <i>PLoS ONE</i> , 2020 , 15, e0232606	3.7	4
47	Abnormal thyroid function parameters in the second trimester of pregnancy are associated with breech presentation at term: a nested cohort study. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2016 , 199, 169-74	2.4	4
46	Internal versus external tocodynamometry during induced or augmented labour. <i>The Cochrane Library</i> , 2013 , CD006947	5.2	4
45	Pharmacokinetics of nifedipine slow-release during sustained tocolysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2015 , 53, 84-91	2	4
44	Determinants of disease course and severity in hyperemesis gravidarum. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2020 , 245, 162-167	2.4	4
43	Intrapartum and neonatal mortality in low-risk term women in midwife-led care and obstetrician-led care at the onset of labor: A national matched cohort study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2020 , 99, 546-554	3.8	4

42	Mode of childbirth and neonatal outcome after external cephalic version: A prospective cohort study. <i>Midwifery</i> , 2016 , 39, 44-8	2.8	4
41	Study protocol for a randomised trial for atosiban versus placebo in threatened preterm birth: the APOSTEL 8 study. <i>BMJ Open</i> , 2019 , 9, e029101	3	4
40	Posttraumatic stress disorder in partners following severe postpartum haemorrhage: A prospective cohort study. <i>Women and Birth</i> , 2020 , 33, 360-366	3.3	4
39	Fetal fibronectin status and cervical length in women with threatened preterm labor and the effectiveness of maintenance tocolysis. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 1556-61	2.61	3
38	Which Factors Contribute to False-Positive, False-Negative, and Invalid Results in Fetal Fibronectin Testing in Women with Symptoms of Preterm Labor?. <i>American Journal of Perinatology</i> , 2017 , 34, 234-239	3.3	3
37	Is short first-trimester crown-rump length associated with spontaneous preterm birth?. <i>Ultrasound in Obstetrics and Gynecology</i> , 2012 , 40, 636-41	5.8	3
36	HELLP syndrome preceded by intrahepatic cholestasis of pregnancy: one serious itch. <i>BMJ Case Reports</i> , 2014 , 2014,	0.9	3
35	What women want and why. Women's preferences for induction of labour or expectant management in late-term pregnancy. <i>Women and Birth</i> , 2021 , 34, 250-256	3.3	3
34	Prescribing patterns of antenatal corticosteroids in women with threatened preterm labor. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2015 , 192, 47-53	2.4	2
33	Severe postpartum hemorrhage increases risk of posttraumatic stress disorder: a prospective cohort study. <i>Journal of Psychosomatic Obstetrics and Gynaecology</i> , 2021 , 42, 335-345	3.6	2
32	Intrapartum and neonatal mortality among low-risk women in midwife-led versus obstetrician-led care in the Amsterdam region of the Netherlands: a propensity score matched study. <i>BMJ Open</i> , 2018 , 8, e018845	3	2
31	488: Economic analysis of fetal fibrinectin testing and/or cervical length measurement in women with threatened preterm labor. <i>American Journal of Obstetrics and Gynecology</i> , 2013 , 208, S210-S211	6.4	2
30	Induction of Labor Versus Expectant Monitoring for Gestational Hypertension or Mild Preeclampsia After 36 Weeks Gestation (HYPITAT): A Multicentre, Open-Label Randomized Controlled Trial. <i>Obstetrical and Gynecological Survey</i> , 2009 , 64, 776-778	2.4	2
29	Morning versus evening induction of labour for improving outcomes. <i>Cochrane Database of Systematic Reviews</i> , 2009 ,		2
28	An unusual cause of hypertension in pregnancy. <i>Hypertension in Pregnancy</i> , 2004 , 23, 13-7	2	2
27	Aspirin and Aspirin Combined with Low-Molecular-Weight Heparin in Women with Unexplained Recurrent Miscarriage: a Randomized Controlled Multicenter Trial (ALIFE Study).. <i>Blood</i> , 2009 , 114, 488-488	2.2	2
26	'Blindness out of the womb', a historical account of the first report of posterior reversible encephalopathy syndrome (PRES): choose your title well or your findings will be neglected. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2020 , 127, 37-38	3.7	2
25	Ethnic differences in the impact of male fetal gender on the risk of spontaneous preterm birth. <i>Journal of Perinatology</i> , 2021 , 41, 2165-2172	3.1	2

24	CONDISOX- continued versus discontinued oxytocin stimulation of induced labour in a double-blind randomised controlled trial. <i>BMC Pregnancy and Childbirth</i> , 2019 , 19, 320	3.2	1
23	HELLP, een potentieel levensbedreigende aandoening. <i>Huisarts En Wetenschap</i> , 2013 , 56, 590-593	0.1	1
22	Association of three different techniques to measure blood pressure in the first trimester with the development of hypertensive disorders of pregnancy. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2013 , 92, 53-60	3.8	1
21	Does measurement of intrauterine pressure have predictive value during oxytocin-augmented labor?. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2016 , 29, 3239-42	2	1
20	Substandard care in maternal mortality due to hypertensive disease in pregnancy in the Netherlands. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2008 , 115, 1322-3; author reply 1323	3.7	1
19	Placental corticotrophin-releasing hormone mRNA and microparticles in maternal plasma are not measures of placental shedding of debris: a rebuttal. <i>Journal of Thrombosis and Haemostasis</i> , 2008 , 6, 1837-8; author reply 1838-9	15.4	1
18	Automated blood pressure measurement in pregnancy. <i>American Journal of Obstetrics and Gynecology</i> , 1994 , 170, 702-3	6.4	1
17	TBX2, a Novel Regulator of Labour. <i>Medicina (Lithuania)</i> , 2021 , 57,	3.1	1
16	Decreasing trend in preterm birth and perinatal mortality, do disparities also decline?. <i>BMC Public Health</i> , 2020 , 20, 783	4.1	0
15	Thyroid-stimulating hormone and free thyroxine fail to predict the severity and clinical course of hyperemesis gravidarum: A prospective cohort study. <i>Acta Obstetricia Et Gynecologica Scandinavica</i> , 2021 , 100, 1419-1429	3.8	0
14	Ethical review of COVID-19 research in the Netherlands; a mixed-method evaluation among medical research ethics committees and investigators. <i>PLoS ONE</i> , 2021 , 16, e0255040	3.7	0
13	The Development of a New Medical Device for Standardized Episiotomy: A Pre-Clinical Validation Study. <i>Gynecologic and Obstetric Investigation</i> , 2016 , 81, 207-14	2.5	
12	HYPITAT and the fallacy of pregnancy interruption [Authors' reply]. <i>Lancet, The</i> , 2010 , 375, 119-120	4.0	
11	A critical view on the use of ambulatory blood pressure monitoring in hypertensive disorders of pregnancy. <i>Journal of Hypertension</i> , 2010 , 28, 1111-1113	1.9	
10	Platelet response to vasopressin in preeclampsia. <i>American Journal of Obstetrics and Gynecology</i> , 1995 , 173, 1897-8	6.4	
9	Pregnancy Complications In Sickle Cell Disease Are More Prevalent In Women with Vaso-Occlusion Related Organ Damage Than Hemolysis Related Organ Damage. <i>Blood</i> , 2010 , 116, 2661-2661	2.2	
8	Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials 2020 , 17, e1003436		
7	Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials 2020 , 17, e1003436		

- 6 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436
- 5 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436
- 4 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436
- 3 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436
- 2 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436
- 1 Induction of labour at 41 weeks or expectant management until 42 weeks: A systematic review and an individual participant data meta-analysis of randomised trials **2020**, 17, e1003436