

Timm Denecke

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8766792/timm-denecke-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

148
papers

4,077
citations

32
h-index

59
g-index

157
ext. papers

4,699
ext. citations

4.2
avg. IF

4.82
L-index

#	Paper	IF	Citations
148	Low skeletal muscle mass defined by thoracic CT as a prognostic marker in acute pulmonary embolism.. <i>Nutrition</i> , 2022 , 98, 111622	4.8	
147	Diagnostic Benefit of High b-Value Computed Diffusion-Weighted Imaging in Patients with Hepatic Metastasis. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	2
146	Toward a Routine Assessment of Visceral Adipose Tissue Volume from Computed Tomographic Data. <i>Obesity</i> , 2021 , 29, 294-301	8	
145	Everolimus-induced pneumonitis in neuroendocrine neoplasms: correlation of CT findings and clinical signs. <i>Acta Radiologica</i> , 2021 , 62, 1006-1015	2	0
144	HBP-enhancing hepatocellular adenomas and how to discriminate them from FNH in Gd-EOB MRI. <i>BMC Medical Imaging</i> , 2021 , 21, 28	2.9	1
143	Postoperative single-sequence (PoSse) MRI: imaging work-up for CT-guided or endoscopic drainage indication of collections after hepatopancreaticobiliary surgery. <i>Abdominal Radiology</i> , 2021 , 46, 3418-3427	2.7	2
142	CT Texture analysis and CT scores for characterization of fluid collections. <i>BMC Medical Imaging</i> , 2021 , 21, 187	2.9	0
141	Diagnostik und Therapie des hepatozellulären Karzinoms. <i>Onkologie</i> , 2020 , 26, 1175-1192	0.1	
140	Portosystemic shunt surgery in the era of TIPS: imaging-based planning of the surgical approach. <i>Abdominal Radiology</i> , 2020 , 45, 2726-2735	3	1
139	Stent performance in palliative transhepatic treatment of malignant biliary obstruction: a randomized study comparing covered versus uncovered stents. <i>Acta Radiologica</i> , 2020 , 61, 1591-1599	2	1
138	Factors That Influence Conversion to Resectability and Survival After Resection of Metastases in RAS WT Metastatic Colorectal Cancer (mCRC): Analysis of FIRE-3- AIOKRK0306. <i>Annals of Surgical Oncology</i> , 2020 , 27, 2389-2401	3.1	7
137	Tumor Seeding along the Puncture Tract in CT-Guided Interstitial High-Dose-Rate Brachytherapy. <i>Journal of Vascular and Interventional Radiology</i> , 2020 , 31, 720-727	2.4	3
136	Hepatocellular adenomas: is there additional value in using Gd-EOB-enhanced MRI for subtype differentiation?. <i>European Radiology</i> , 2020 , 30, 3497-3506	8	10
135	Advantages of a T1-Weighted Gradient-Recalled Echo (GRE) Sequence With a Radial 3D Sampling Approach Versus 2D Turbo Spin-Echo and Cartesian 3D GRE Sequences in Head and Neck MRI. <i>American Journal of Roentgenology</i> , 2020 , 214, 747-753	5.4	5
134	Semi-automatic prostatic artery detection using cone-beam CT during prostatic arterial embolization. <i>Acta Radiologica</i> , 2020 , 61, 1116-1124	2	2
133	Complications of Computed Tomography-Guided High-Dose-Rate Brachytherapy (CT-HDRBT) and Risk Factors: Results from More than 10 Years of Experience. <i>CardioVascular and Interventional Radiology</i> , 2020 , 43, 284-294	2.7	6
132	Estimation of abdominal subcutaneous fat volume of obese adults from single-slice MRI data - Regression coefficients and agreement. <i>European Journal of Radiology</i> , 2020 , 130, 109184	4.7	2

131	Abdominal subcutaneous fat quantification in obese patients from limited field-of-view MRI data. <i>Scientific Reports</i> , 2020 , 10, 19039	4.9	1
130	Prognostic Significance of Somatostatin Receptor Heterogeneity in Progressive Neuroendocrine Tumor Treated with Lu-177 DOTATOC or Lu-177 DOTATATE. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2020 , 47, 881-894	8.8	16
129	Gd-EOB-DTPA-enhanced MRI T1 relaxometry as an imaging-based liver function test compared with C-methacetin breath test. <i>Acta Radiologica</i> , 2020 , 61, 291-301	2	5
128	Iterative CT reconstruction in abdominal low-dose CT used for hybrid SPECT-CT applications: effect on image quality, image noise, detectability, and reader's confidence. <i>Acta Radiologica Open</i> , 2019 , 8, 2058460119856266	1.2	4
127	Consistency of hepatocellular gadoxetic acid uptake in serial MRI examinations for evaluation of liver function. <i>Abdominal Radiology</i> , 2019 , 44, 2759-2768	3	7
126	Predicting liver failure after extended right hepatectomy following right portal vein embolization with gadoxetic acid-enhanced MRI. <i>European Radiology</i> , 2019 , 29, 5861-5872	8	11
125	Combined morphological and functional liver MRI using spin-lattice relaxation in the rotating frame (T1 ρ) in conjunction with Gadoxetic Acid-enhanced MRI. <i>Scientific Reports</i> , 2019 , 9, 2083	4.9	3
124	DCE-MR imaging of orbital lesions: diagnostic performance of the tumor flow residence time τ calculated by a multi-compartmental pharmacokinetic tumor model based on individual factors. <i>Acta Radiologica</i> , 2019 , 60, 643-652	2	6
123	Single-incision laparoscopic surgery portal vein embolisation before extended hepatectomy. <i>Journal of Minimal Access Surgery</i> , 2019 ,	1.2	1
122	Optimized imaging of the lower abdomen and pelvic region in hepatocyte-specific MRI: evaluation of a whole-abdomen first-pass shuttle protocol in patients with neuroendocrine neoplasms. <i>Acta Radiologica</i> , 2019 , 60, 1074-1083	2	2
121	Strengths, Weaknesses, Opportunities, and Threats of Centralized Pancreatic Surgery: a Single-Center Analysis of 3000 Consecutive Pancreatic Resections. <i>Journal of Gastrointestinal Surgery</i> , 2019 , 23, 492-502	3.3	1
120	The Predictive Value of the Maximal Liver Function Capacity Test for the Isolation of Primary Human Hepatocytes. <i>Tissue Engineering - Part C: Methods</i> , 2018 , 24, 179-186	2.9	
119	Performance survey on a new standardized formula for oral signal suppression in MRCP. <i>European Journal of Radiology Open</i> , 2018 , 5, 1-5	2.6	3
118	Comparison of ultrasound shear wave elastography with magnetic resonance elastography and renal microvascular flow in the assessment of chronic renal allograft dysfunction. <i>Acta Radiologica</i> , 2018 , 59, 1139-1145	2	14
117	Degenerative changes after posterior cruciate ligament reconstruction are irrespective of posterior knee stability: MRI-based long-term results. <i>Archives of Orthopaedic and Trauma Surgery</i> , 2018 , 138, 377-385	3.6	10
116	Cholangiocarcinoma: CT-guided High-Dose Rate Brachytherapy (CT-HDRBT) for Limited (4 cm) Tumors. <i>Anticancer Research</i> , 2018 , 38, 5843-5852	2.3	10
115	The role of hepatectomy for synchronous liver metastases from pancreatic adenocarcinoma. <i>Surgical Oncology</i> , 2018 , 27, 688-694	2.5	14
114	Multiple solid pancreatic lesions: Prevalence and features of non-malignancies on dynamic enhanced CT. <i>European Journal of Radiology</i> , 2018 , 105, 8-14	4.7	5

113	Routine portal vein resection for pancreatic adenocarcinoma shows no benefit in overall survival. <i>European Journal of Surgical Oncology</i> , 2018 , 44, 1094-1099	3.6	8
112	Is There Long-term Signal Intensity Increase in the Central Nervous System on T1-weighted Images after MR Imaging with the Hepatospecific Contrast Agent Gadoteric Acid? A Cross-sectional Study in 91 Patients. <i>Radiology</i> , 2017 , 282, 708-716	20.5	44
111	The Magnetic Field of Magnetic Resonance Imaging Systems Does Not Affect Cells Labeled with Micrometer-Sized Iron Oxide Particles. <i>Tissue Engineering - Part C: Methods</i> , 2017 , 23, 412-421	2.9	1
110	Gd-EOB-DTPA-enhanced MRI for monitoring future liver remnant function after portal vein embolization and extended hemihepatectomy: A prospective trial. <i>European Radiology</i> , 2017 , 27, 3080-3087	8.87	20
109	Efficacy of oral contrast agents for upper gastrointestinal signal suppression in MRCP: A systematic review of the literature. <i>Acta Radiologica Open</i> , 2017 , 6, 2058460117727315	1.2	15
108	Blood Transfusions and Tumor Biopsy May Increase HCC Recurrence Rates after Liver Transplantation. <i>Journal of Transplantation</i> , 2017 , 2017, 9731095	2.3	8
107	Segment-by-segment assessment of left ventricular myocardial affection in Anderson-Fabry disease by non-enhanced T1-mapping. <i>Acta Radiologica</i> , 2017 , 58, 914-921	2	7
106	Initial Experience with CT-Guided High-Dose-Rate Brachytherapy in the Multimodality Treatment of Neuroendocrine Tumor Liver Metastases. <i>Journal of Vascular and Interventional Radiology</i> , 2017 , 28, 672-682	2.4	5
105	Influence of fractional anisotropy thresholds on diffusion tensor imaging tractography of the periprostatic neurovascular bundle and selected pelvic tissues: do visualized tracts really represent nerves?. <i>Acta Radiologica</i> , 2017 , 58, 472-480	2	2
104	Impact of Doppler Ultrasound on Diagnosis and Therapy Control of Lialis Steal Syndrome After Liver Transplantation. <i>Annals of Transplantation</i> , 2017 , 22, 440-445	1.4	1
103	Diagnostic imaging of pancreatic neuroendocrine neoplasms (pNEN): tumor detection, staging, prognosis, and response to treatment. <i>Acta Radiologica</i> , 2016 , 57, 260-70	2	24
102	Clinical application of modern ultrasound techniques after liver transplantation. <i>Acta Radiologica</i> , 2016 , 57, 1161-70	2	12
101	Potential role of (68)Ga-DOTATOC PET/CT in screening for pancreatic neuroendocrine tumour in patients with von Hippel-Lindau disease. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2016 , 43, 2014-20	8.8	22
100	Significance of a Single-Time-Point Somatostatin Receptor SPECT/Multiphase CT Protocol in the Diagnostic Work-up of Gastroenteropancreatic Neuroendocrine Neoplasms. <i>Journal of Nuclear Medicine</i> , 2016 , 57, 180-5	8.9	4
99	Management of follow-up of neuroendocrine neoplasias. <i>Best Practice and Research in Clinical Endocrinology and Metabolism</i> , 2016 , 30, 129-40	6.5	6
98	Perioperative treatment options in resectable pancreatic cancer - how to improve long-term survival. <i>World Journal of Gastrointestinal Oncology</i> , 2016 , 8, 248-57	3.4	18
97	Maximizing Information From Routine Staging Computed Tomography in Functional Neuroendocrine Neoplasms: Are There Findings Indicating the Presence of Carcinoid Heart Disease?. <i>Journal of Computer Assisted Tomography</i> , 2016 , 40, 277-82	2.2	2
96	Cardiovascular Magnetic Resonance Imaging in Patients with an Implantable Loop Recorder. <i>Annals of Noninvasive Electrocardiology</i> , 2016 , 21, 319-324	1.5	8

95	Role of (68)Ga somatostatin receptor PET/CT in the detection of endogenous hyperinsulinaemic focus: an explorative study. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2016 , 43, 1593-600	8.8	31
94	Patient and tumour biology predict survival beyond the Milan criteria in liver transplantation for hepatocellular carcinoma. <i>Hpb</i> , 2015 , 17, 168-75	3.8	16
93	Imaging-based evaluation of liver function: comparison of ^{99m} Tc-mebrofenin hepatobiliary scintigraphy and Gd-EOB-DTPA-enhanced MRI. <i>European Radiology</i> , 2015 , 25, 1384-91	8	23
92	CT-guided high-dose-rate brachytherapy of unresectable hepatocellular carcinoma. <i>Strahlentherapie Und Onkologie</i> , 2015 , 191, 405-12	4.3	21
91	Portal vein embolization with plug/coils improves hepatectomy outcome. <i>Journal of Surgical Research</i> , 2015 , 194, 202-11	2.5	23
90	CT-guided Interstitial Brachytherapy of Hepatocellular Carcinoma before Liver Transplantation: an Equivalent Alternative to Transarterial Chemoembolization?. <i>European Radiology</i> , 2015 , 25, 2608-16	8	23
89	Streptozocin/5-fluorouracil chemotherapy is associated with durable response in patients with advanced pancreatic neuroendocrine tumours. <i>European Journal of Cancer</i> , 2015 , 51, 1253-62	7.5	69
88	Somatostatin receptor PET/CT in restaging of typical and atypical lung carcinoids. <i>EJNMMI Research</i> , 2015 , 5, 53	3.6	13
87	Efficacy of Prophylactic Low-Molecular Weight Heparin for Ambulatory Patients With Advanced Pancreatic Cancer: Outcomes From the CONKO-004 Trial. <i>Journal of Clinical Oncology</i> , 2015 , 33, 2028-34	2.2	172
86	Gd-EOB enhanced MRI T1-weighted 3D-GRE with and without elevated flip angle modulation for threshold-based liver volume segmentation. <i>Acta Radiologica</i> , 2015 , 56, 1419-27	2	5
85	Factors influencing hypertrophy of the left lateral liver lobe after portal vein embolization. <i>Langenbecks Archives of Surgery</i> , 2015 , 400, 237-46	3.4	12
84	Gadoxetic acid enhanced MRI for differentiation of FNH and HCA: a single centre experience. <i>European Radiology</i> , 2014 , 24, 1339-48	8	39
83	CT-guided high-dose-rate brachytherapy of liver tumours does not impair hepatic function and shows high overall safety and favourable survival rates. <i>Annals of Surgical Oncology</i> , 2014 , 21, 4284-92	3.1	12
82	Comparison of CT and MRI artefacts from coils and vascular plugs used for portal vein embolization. <i>European Journal of Radiology</i> , 2014 , 83, 692-5	4.7	5
81	An increased flip angle in late phase Gd-EOB-DTPA MRI shows improved performance in bile duct visualization compared to T2w-MRCP. <i>European Journal of Radiology</i> , 2014 , 83, 1723-7	4.7	8
80	Response to preoperative chemotherapy predicts survival in patients undergoing hepatectomy for liver metastases from gastric and esophageal cancer. <i>Journal of Gastrointestinal Surgery</i> , 2014 , 18, 1974-86	3.3	12
79	Optimized separation of left and right liver lobe in dynamic (99m)Tc-mebrofenin hepatobiliary scintigraphy using a hybrid SPECT-CT scanner. <i>Annals of Nuclear Medicine</i> , 2014 , 28, 897-902	2.5	1
78	Evaluation of gadolinium-EOB-DTPA uptake after portal vein embolization: value of an increased flip angle. <i>Acta Radiologica</i> , 2014 , 55, 149-54	2	5

77	Improved hypertrophy of future remnant liver after portal vein embolization with plugs, coils and particles. <i>CardioVascular and Interventional Radiology</i> , 2014 , 37, 1251-8	2.7	23
76	No infectious hepatic complications following radioembolization with 90Y microspheres in patients with biliodigestive anastomosis. <i>Anticancer Research</i> , 2014 , 34, 4315-21	2.3	10
75	Increase in left liver lobe function after preoperative right portal vein embolisation assessed with gadolinium-EOB-DTPA MRI. <i>European Radiology</i> , 2013 , 23, 2555-60	8	16
74	Treatment for liver metastasis from renal cell carcinoma with computed-tomography-guided high-dose-rate brachytherapy (CT-HDRBT): a case series. <i>World Journal of Urology</i> , 2013 , 31, 1525-30	4	13
73	Long-term results of liver resection for hepatocellular carcinoma in noncirrhotic liver. <i>Surgery</i> , 2013 , 153, 510-7	3.6	34
72	Size determination and response assessment of liver metastases with computed tomography--comparison of RECIST and volumetric algorithms. <i>European Journal of Radiology</i> , 2013 , 82, 1831-9	4.7	31
71	Computed-tomography-guided high-dose-rate brachytherapy (CT-HDRBT) ablation of metastases adjacent to the liver hilum. <i>European Journal of Radiology</i> , 2013 , 82, e509-14	4.7	35
70	Bilateral subtotal thyroidectomy versus hemithyroidectomy plus subtotal resection (Dunhill procedure) for benign goiter: long-term results of a prospective, randomized study. <i>World Journal of Surgery</i> , 2013 , 37, 84-90	3.3	22
69	Histopathologically confirmed focal nodular hyperplasia of the liver: gadoxetic acid-enhanced MRI characteristics. <i>Magnetic Resonance Imaging</i> , 2013 , 31, 755-60	3.3	28
68	Evaluation of radiological prognostic factors of hepatic metastases in patients with non-functional pancreatic neuroendocrine tumors. <i>European Journal of Radiology</i> , 2013 , 82, e550-5	4.7	20
67	Waitlist characteristics of patients at a single-center intestinal and multivisceral transplant program. <i>Transplant International</i> , 2013 , 26, 392-401	3	4
66	Computed tomography findings in septic patients with acute respiratory distress syndrome: correlation with survival and pulmonary versus extrapulmonary septic focus. <i>Journal of Computer Assisted Tomography</i> , 2013 , 37, 602-9	2.2	3
65	Quantification in (68)Ga-DOTA(0)-Phe(1)-Tyr(3)-octreotide positron emission tomography/computed tomography: can we be impartial about partial volume effects?. <i>Neuroendocrinology</i> , 2013 , 97, 369-74	5.6	11
64	Impact of neoadjuvant transarterial chemoembolization on tumor recurrence and patient survival after liver transplantation for hepatocellular carcinoma: a retrospective analysis. <i>Clinical Transplantation</i> , 2012 , 26, 764-74	3.8	31
63	Oncological superiority of hilar en bloc resection for the treatment of hilar cholangiocarcinoma. <i>Annals of Surgical Oncology</i> , 2012 , 19, 1602-8	3.1	179
62	Preliminary experience with CT-guided high-dose rate brachytherapy as an alternative treatment for hepatic recurrence of cholangiocarcinoma. <i>Hpb</i> , 2012 , 14, 791-7	3.8	3
61	Percutaneous computed tomography-guided high-dose-rate brachytherapy ablation of breast cancer liver metastases: initial experience with 80 lesions. <i>Journal of Vascular and Interventional Radiology</i> , 2012 , 23, 618-26	2.4	20
60	Computed tomography findings from patients with ARDS due to Influenza A (H1N1) virus-associated pneumonia. <i>European Journal of Radiology</i> , 2012 , 81, 389-94	4.7	23

59	An unusual cause of recurrent acute abdominal pain. <i>Gastroenterology</i> , 2012 , 142, e5-6	13.3	2
58	Magnetic resonance enteroclysis in patients with Crohn's disease: fat saturated T2-weighted sequences for evaluation of inflammatory activity. <i>Journal of Crohns and Colitis</i> , 2012 , 6, 294-301	1.5	9
57	Flip angle modulations in late phase Gd-EOB-DTPA MRI improve the identification of the biliary system. <i>European Journal of Radiology</i> , 2012 , 81, e991-5	4.7	16
56	ENETS Consensus Guidelines for the management of patients with neuroendocrine neoplasms from the jejunum-ileum and the appendix including goblet cell carcinomas. <i>Neuroendocrinology</i> , 2012 , 95, 135-56	5.6	316
55	Matched-pair comparison of radioembolization plus best supportive care versus best supportive care alone for chemotherapy refractory liver-dominant colorectal metastases. <i>CardioVascular and Interventional Radiology</i> , 2012 , 35, 1066-73	2.7	119
54	SUV-measurements and patient-specific corrections in pediatric Hodgkin-lymphoma: is there a benefit for PPV in early response assessment by FDG-PET?. <i>Pediatric Blood and Cancer</i> , 2012 , 59, 475-80 ³		13
53	Computed tomography-guided interstitial HDR brachytherapy (CT-HDRBT) of the liver in patients with irresectable intrahepatic cholangiocarcinoma. <i>CardioVascular and Interventional Radiology</i> , 2012 , 35, 581-7	2.7	27
52	Hepatocellular carcinoma: computed-tomography-guided high-dose-rate brachytherapy (CT-HDRBT) ablation of large (5-7 cm) and very large (>7 cm) tumours. <i>European Radiology</i> , 2012 , 22, 1101-9	8	50
51	Appearance of hepatocellular adenomas on gadoxetic acid-enhanced MRI. <i>European Radiology</i> , 2012 , 22, 1769-75	8	38
50	Formation of a chronic pain syndrome due to mesh shrinkage after laparoscopic intraperitoneal onlay mesh (I POM). <i>Surgical Laparoscopy, Endoscopy and Percutaneous Techniques</i> , 2012 , 22, e288-90	1.3	10
49	Treatment of hepatic metastases from gastric or gastroesophageal adenocarcinoma with computed tomography-guided high-dose-rate brachytherapy (CT-HDRBT). <i>Anticancer Research</i> , 2012 , 32, 5453-8	2.3	11
48	Repeated liver resection for recurrent hepatocellular carcinoma. <i>Journal of Gastroenterology and Hepatology (Australia)</i> , 2011 , 26, 1189-94	4	32
47	Assessment of the cerebellar arteries with multidetector computed tomography angiography benefits from submillimeter slice thickness. <i>Clinical Imaging</i> , 2011 , 35, 247-52	2.7	4
46	Distal pancreatectomy with en bloc resection of the celiac trunk for extended pancreatic tumor disease: an interdisciplinary approach. <i>CardioVascular and Interventional Radiology</i> , 2011 , 34, 1058-64	2.7	36
45	Safety of liver resection and effect on quality of life in patients with benign hepatic disease: single center experience. <i>BMC Surgery</i> , 2011 , 11, 16	2.3	19
44	Arterial versus portal venous embolization for induction of hepatic hypertrophy before extended right hemihepatectomy in hilar cholangiocarcinomas: a prospective randomized study. <i>Journal of Vascular and Interventional Radiology</i> , 2011 , 22, 1254-62	2.4	18
43	Laparoscopic radiofrequency ablation of liver tumors: comparison of MR guidance versus conventional laparoscopic ultrasound for needle positioning in a phantom model. <i>Minimally Invasive Therapy and Allied Technologies</i> , 2011 , 20, 212-7	2.1	4
42	⁶⁸ Ga-DOTATOC PET/CT of neuroendocrine tumors: spotlight on the CT phases of a triple-phase protocol. <i>Journal of Nuclear Medicine</i> , 2011 , 52, 697-704	8.9	74

41	CT-guided high-dose-rate brachytherapy of metachronous ovarian cancer metastasis to the liver: initial experience. <i>Anticancer Research</i> , 2011 , 31, 2597-602	2.3	11
40	Imaging of hypoxic-ischemic penumbra with (18)F-fluoromisonidazole PET/CT and measurement of related cerebral metabolism in aneurysmal subarachnoid hemorrhage. <i>Journal of Cerebral Blood Flow and Metabolism</i> , 2010 , 30, 36-45	7.3	33
39	Evolving experience with prevention and treatment of splenic artery syndrome after orthotopic liver transplantation. <i>Transplant International</i> , 2010 , 23, 831-41	3	39
38	Preoperative multidetector row computed tomography for evaluation and assessment of resection criteria in patients with pancreatic masses. <i>Acta Radiologica</i> , 2010 , 51, 1067-77	2	11
37	Impact of Multiphase 68Ga-DOTATOC-PET/CT on therapy management in patients with neuroendocrine tumors. <i>Neuroendocrinology</i> , 2010 , 91, 101-9	5.6	94
36	Multidetector computed tomography for preoperative assessment of hepatic vasculature and prediction of splenic artery steal syndrome in patients with liver cirrhosis before transplantation. <i>European Radiology</i> , 2010 , 20, 108-17	8	28
35	Potential impact of (68)Ga-DOTATOC PET/CT on stereotactic radiotherapy planning of meningiomas. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2010 , 37, 310-8	8.8	64
34	Assessment of histological response of paediatric bone sarcomas using FDG PET in comparison to morphological volume measurement and standardized MRI parameters. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2010 , 37, 1842-53	8.8	91
33	Modification of aminosilanized superparamagnetic nanoparticles: feasibility of multimodal detection using 3T MRI, small animal PET, and fluorescence imaging. <i>Molecular Imaging and Biology</i> , 2010 , 12, 25-34	3.8	65
32	Value of combined PET/CT for radiation planning in CT-guided percutaneous interstitial high-dose-rate single-fraction brachytherapy for colorectal liver metastases. <i>International Journal of Radiation Oncology Biology Physics</i> , 2010 , 77, 1178-85	4	13
31	Multislice computed tomography using a triple-phase contrast protocol for preoperative assessment of hepatic tumor load in patients with hepatocellular carcinoma before liver transplantation. <i>Transplant International</i> , 2009 , 22, 395-402	3	22
30	Extended bile duct resection and [corrected] liver and transplantation in patients with hilar cholangiocarcinoma: long-term results. <i>Liver Transplantation</i> , 2009 , 15, 1499-507	4.5	35
29	Early and late therapy response assessment with [18F]fluorodeoxyglucose positron emission tomography in pediatric Hodgkin's lymphoma: analysis of a prospective multicenter trial. <i>Journal of Clinical Oncology</i> , 2009 , 27, 4385-91	2.2	129
28	Directing Minimal Invasive Image Guided Therapy of Hepatic Colorectal Cancer Metastases □ Imaging Strategies for Patient Evaluation, Therapy Planning, Therapy Monitoring, and Follow-Up. <i>Current Medical Imaging</i> , 2009 , 5, 128-135	1.2	
27	Perfusion Scintigraphy with Integrated Single Photon Emission Computed Tomography/Computed Tomography in the Management of Transarterial Treatment of Hepatic Malignancies 2009 , 287-301		
26	Clinical Challenges and Images in GI: image 1. Mesenteric lipoma with pancreatic heterotopy. <i>Gastroenterology</i> , 2008 , 135, 741, 1018	13.3	
25	An orthotopic model of pancreatic somatostatin receptor (SSTR)-positive tumors allows bimodal imaging studies using 3T MRI and animal PET-based molecular imaging of SSTR expression. <i>Neuroendocrinology</i> , 2008 , 87, 233-42	5.6	8
24	Suspected chronic myocarditis at cardiac MR: diagnostic accuracy and association with immunohistologically detected inflammation and viral persistence. <i>Radiology</i> , 2008 , 246, 401-9	20.5	214

23	Use of positron emission tomography for staging, preoperative response assessment and posttherapeutic evaluation in children with Wilms tumour. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2008 , 35, 1642-50	8.8	41
22	Planning transarterial radioembolization of colorectal liver metastases with Yttrium 90 microspheres: evaluation of a sequential diagnostic approach using radiologic and nuclear medicine imaging techniques. <i>European Radiology</i> , 2008 , 18, 892-902	8	50
21	Brachytherapy of liver metastases. <i>Recent Results in Cancer Research</i> , 2008 , 177, 95-104	1.5	13
20	Transhepatic computed tomography-guided management of bleeding hepatic arterial pseudoaneurysm inaccessible for angiography in a liver transplant recipient. <i>Transplant International</i> , 2007 , 20, 728-30	3	3
19	PET/CT visualises inflammatory activity of pulmonary artery aneurysms in Behçt disease. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2007 , 34, 970	8.8	18
18	Positron emission tomography for staging of pediatric sarcoma patients: results of a prospective multicenter trial. <i>Journal of Clinical Oncology</i> , 2007 , 25, 5435-41	2.2	313
17	Influence of attenuation correction by integrated low-dose CT on somatostatin receptor SPECT. <i>Nuclear Medicine Communications</i> , 2007 , 28, 782-8	1.6	26
16	Multislice computed tomographic angiography for preinterventional planning of port placement for intra-arterial hepatic infusion chemotherapy. <i>Journal of Computer Assisted Tomography</i> , 2007 , 31, 66-71	2.2	5
15	Interventional Radiotherapy 2007 , 47-58		
14	Diagnostic value of 123I-IMT SPECT in the follow-up of head and neck cancer. <i>Oncology Research and Treatment</i> , 2006 , 29, 147-52	2.8	1
13	Long-lasting remission of pulmonary metastases of renal cell cancer under IFN-beta therapy in a patient with multiple sclerosis. <i>Oncology Research and Treatment</i> , 2006 , 29, 382-4	2.8	2
12	18F-FET PET for planning of thermotherapy using magnetic nanoparticles in recurrent glioblastoma. <i>International Journal of Hyperthermia</i> , 2006 , 22, 319-25	3.7	32
11	Relevance of image fusion with MRI for the interpretation of I-123 iodo-methyl-tyrosine scans in patients with suspected recurrent or residual brain tumor. <i>Clinical Nuclear Medicine</i> , 2006 , 31, 189-92	1.7	2
10	Correlative imaging strategies implementing CT, MRI, and PET for staging of childhood Hodgkin disease. <i>Journal of Pediatric Hematology/Oncology</i> , 2006 , 28, 501-12	1.2	50
9	Combined SPECT/CT imaging using 123I-IMT in the detection of recurrent or persistent head and neck cancer. <i>European Radiology</i> , 2006 , 16, 503-11	8	6
8	Value of axial and coronal maximum intensity projection (MIP) images in the detection of pulmonary nodules by multislice spiral CT: comparison with axial 1-mm and 5-mm slices. <i>European Radiology</i> , 2006 , 16, 325-32	8	65
7	Minimum intensity projections of the biliary system using 16-channel multidetector computed tomography in patients with biliary obstruction: comparison with MRCP. <i>European Radiology</i> , 2006 , 16, 1719-26	8	28
6	Comparison of CT, MRI and FDG-PET in response prediction of patients with locally advanced rectal cancer after multimodal preoperative therapy: is there a benefit in using functional imaging?. <i>European Radiology</i> , 2005 , 15, 1658-66	8	174

5	Influence of high magnetic field strengths and parallel acquisition strategies on image quality in cardiac 2D CINE magnetic resonance imaging: comparison of 1.5 T vs. 3.0 T. <i>European Radiology</i> , 2005 , 15, 1586-97	8	76
4	Preoperative evaluation of living kidney donors using multirow detector computed tomography: comparison with digital subtraction angiography and intraoperative findings. <i>Transplant International</i> , 2005 , 18, 1134-41	3	38
3	Fusion imaging using a hybrid SPECT-CT camera improves port perfusion scintigraphy for control of hepatic arterial infusion of chemotherapy in colorectal cancer patients. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2005 , 32, 1003-10	8.8	13
2	Response prediction by FDG-PET after neoadjuvant radiochemotherapy and combined regional hyperthermia of rectal cancer: correlation with endorectal ultrasound and histopathology. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2004 , 31, 811-9	8.8	97
1	Eight-channel multidetector computed tomography: unique potential for pediatric chest computed tomography angiography. <i>Journal of Thoracic Imaging</i> , 2002 , 17, 306-9	5.6	12