Motoki Iwasaki

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8769656/publications.pdf Version: 2024-02-01



MOTOKI IWASAKI

#	Article	IF	CITATIONS
1	Diabetes Mellitus and the Risk of Cancer. Archives of Internal Medicine, 2006, 166, 1871.	3.8	475
2	Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. Nature Genetics, 2017, 49, 1458-1467.	21.4	380
3	Green Tea Consumption and Prostate Cancer Risk in Japanese Men: A Prospective Study. American Journal of Epidemiology, 2007, 167, 71-77.	3.4	241
4	Plasma Isoflavone Level and Subsequent Risk of Breast Cancer Among Japanese Women: A Nested Case-Control Study From the Japan Public Health Center-Based Prospective Study Group. Journal of Clinical Oncology, 2008, 26, 1677-1683.	1.6	155
5	Daily Total Physical Activity Level and Premature Death in Men and Women: Results From a Large-Scale Population-Based Cohort Study in Japan (JPHC Study). Annals of Epidemiology, 2008, 18, 522-530.	1.9	147
6	Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. European Journal of Cancer Prevention, 2009, 18, 240-247.	1.3	131
7	Effect of Helicobacter pylori Infection Combined with CagA and Pepsinogen Status on Gastric Cancer Development among Japanese Men and Women: A Nested Case-Control Study. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 1341-1347.	2.5	123
8	Characterizing rare and low-frequency height-associated variants in the Japanese population. Nature Communications, 2019, 10, 4393.	12.8	123
9	Association of Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a Japanese Cohort. JAMA Internal Medicine, 2019, 179, 1509.	5.1	120
10	Dietary cadmium intake and breast cancer risk in Japanese women: A case–control study. International Journal of Hygiene and Environmental Health, 2014, 217, 70-77.	4.3	115
11	Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2010, 91, 456-464.	4.7	100
12	Effect of Coffee and Green Tea Consumption on the Risk of Liver Cancer: Cohort Analysis by Hepatitis Virus Infection Status. Cancer Epidemiology Biomarkers and Prevention, 2009, 18, 1746-1753.	2.5	98
13	Plasma C-Reactive Protein and Risk of Colorectal Cancer in a Nested Case-Control Study: Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2006, 15, 690-695.	2.5	94
14	Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study. Cancer Causes and Control, 2007, 18, 199-209.	1.8	88
15	Passive smoking and lung cancer in Japanese nonâ€smoking women: A prospective study. International Journal of Cancer, 2008, 122, 653-657.	5.1	81
16	Prediction of the 10â€year probability of gastric cancer occurrence in the <scp>J</scp> apanese population: the <scp>JPHC</scp> study cohort <scp>II</scp> . International Journal of Cancer, 2016, 138, 320-331.	5.1	78
17	Isoflavone intake and risk of lung cancer: a prospective cohort study in Japan. American Journal of Clinical Nutrition, 2010, 91, 722-728.	4.7	77
18	Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: A prospective cohort study in Japan. Cancer Detection and Prevention, 2007, 31, 3-11.	2.1	75

#	Article	IF	CITATIONS
19	Heterocyclic amines content of meat and fish cooked by Brazilian methods. Journal of Food Composition and Analysis, 2010, 23, 61-69.	3.9	74
20	Body Mass Index, Body Height, and Subsequent Risk of Colorectal Cancer in Middle-Aged and Elderly Japanese Men and Women: Japan Public Health Center-Based Prospective Study. Cancer Causes and Control, 2005, 16, 839-850.	1.8	72
21	Serum organochlorines and breast cancer risk in Japanese women: a case–control study. Cancer Causes and Control, 2009, 20, 567-580.	1.8	70
22	Genetic polymorphisms of ADH1B, ADH1C and ALDH2, alcohol consumption, and the risk of gastric cancer: the Japan Public Health Center-based prospective study. Carcinogenesis, 2015, 36, 223-231.	2.8	69
23	Social support and cancer incidence and mortality: the JPHC study cohort II. Cancer Causes and Control, 2013, 24, 847-860.	1.8	68
24	Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). Annals of Epidemiology, 2015, 25, 512-518.e3.	1.9	66
25	Cigarette Smoking and Completed Suicide among Middle-aged Men: A Population-based Cohort Study in Japan. Annals of Epidemiology, 2005, 15, 286-292.	1.9	61
26	Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. BMJ: British Medical Journal, 2018, 360, k671.	2.3	61
27	Plasma levels of carotenoids, retinol and tocopherol and the risk of gastric cancer in Japan: a nested case–control study. Carcinogenesis, 2008, 29, 1042-1048.	2.8	60
28	Increased Levels of Branched-Chain Amino Acid Associated With Increased Risk of Pancreatic Cancer in a Prospective Case–Control Study of a Large Cohort. Gastroenterology, 2018, 155, 1474-1482.e1.	1.3	59
29	Long-term Dietary Cadmium Intake and Cancer Incidence. Epidemiology, 2012, 23, 368-376.	2.7	58
30	Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2015, 101, 1029-1037.	4.7	58
31	10-Year risk of colorectal cancer: Development and validation of a prediction model in middle-aged Japanese men. Cancer Epidemiology, 2010, 34, 534-541.	1.9	56
32	Role and impact of menstrual and reproductive factors on breast cancer risk in Japan. European Journal of Cancer Prevention, 2007, 16, 116-123.	1.3	54
33	Risk factors for breast cancer: epidemiological evidence from Japanese studies. Cancer Science, 2011, 102, 1607-1614.	3.9	53
34	Green tea drinking and subsequent risk of breast cancer in a population to based cohort of Japanese women. Breast Cancer Research, 2010, 12, R88.	5.0	52
35	Alcohol and smoking and subsequent risk of prostate cancer in Japanese men: The Japan Public Health Centerâ€based prospective study. International Journal of Cancer, 2014, 134, 971-978.	5.1	52
36	GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. Carcinogenesis, 2018, 39, 652-660.	2.8	52

#	Article	IF	CITATIONS
37	Diabetes and cancer risk: A Mendelian randomization study. International Journal of Cancer, 2020, 146, 712-719.	5.1	52
38	Body Size and Risk for Breast Cancer in Relation to Estrogen and Progesterone Receptor Status in Japan. Annals of Epidemiology, 2007, 17, 304-312.	1.9	51
39	Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2017, 147, 1749-1756.	2.9	51
40	Red meat intake may increase the risk of colon cancer in Japanese, a population with relatively low red meat consumption. Asia Pacific Journal of Clinical Nutrition, 2011, 20, 603-12.	0.4	51
41	Background Characteristics of Basic Health Examination Participants: the JPHC Study Baseline Survey. Journal of Epidemiology, 2003, 13, 216-225.	2.4	49
42	Dietary fiber intake and subsequent risk of colorectal cancer: The Japan Public Health Centerâ€Based Prospective Study. International Journal of Cancer, 2006, 119, 1475-1480.	5.1	48
43	Metabolic factors and subsequent risk of hepatocellular carcinoma by hepatitis virus infection status: a large-scale population-based cohort study of Japanese men and women (JPHC Study Cohort II). Cancer Causes and Control, 2009, 20, 741-750.	1.8	48
44	Plasma tea polyphenol levels and subsequent risk of breast cancer among Japanese women: a nested case–control study. Breast Cancer Research and Treatment, 2010, 124, 827-834.	2.5	47
45	Plasma Tea Polyphenols and Gastric Cancer Risk: A Case-Control Study Nested in a Large Population-Based Prospective Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 343-351.	2.5	46
46	Validity of a Self-Administered Food Frequency Questionnaire for Middle-Aged Urban Cancer Screenees: Comparison With 4-Day Weighed Dietary Records. Journal of Epidemiology, 2011, 21, 447-458.	2.4	46
47	Serum triglycerides and colorectal adenoma in a case–control study among cancer screening examinees (Japan). Cancer Causes and Control, 2006, 17, 1245-1252.	1.8	45
48	Death by suicide and other externally caused injuries following a cancer diagnosis: the Japan Public Health Centerâ€based Prospective Study. Psycho-Oncology, 2014, 23, 1034-1041.	2.3	45
49	Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. BMJ, The, 2020, 368, m34.	6.0	45
50	Plasma insulin, <scp>C</scp> â€peptide and blood glucose and the risk of gastric cancer: The <scp>J</scp> apan <scp>P</scp> ublic <scp>H</scp> ealth <scp>C</scp> enterâ€based prospective study. International Journal of Cancer, 2015, 136, 1402-1410.	5.1	44
51	Serum aminotransferase level and the risk of hepatocellular carcinoma: a population-based cohort study in Japan. European Journal of Cancer Prevention, 2009, 18, 26-32.	1.3	43
52	Alcohol consumptionâ€associated breast cancer incidence and potential effect modifiers: the Japan Public Health Centerâ€based Prospective Study. International Journal of Cancer, 2010, 127, 685-695.	5.1	40
53	Dietary isoflavone intake and breast cancer risk in case–control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. Breast Cancer Research and Treatment, 2009, 116, 401-411.	2.5	39
54	Risk factors for epithelial ovarian cancer in Japan - results from the Japan Public Health Center-based Prospective Study cohort. International Journal of Oncology, 2011, 40, 21-30.	3.3	39

#	Article	IF	CITATIONS
55	Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. American Journal of Clinical Nutrition, 2020, 111, 1027-1035.	4.7	38
56	Leisure-time physical activity and breast cancer risk defined by estrogen and progesterone receptor status—The Japan Public Health Center-based Prospective Study. Preventive Medicine, 2011, 52, 227-233.	3.4	37
57	Rural-urban Differences in Sociodemographic, Social Network and Lifestyle Factors Related to Mortality of Middle-aged Japanese Men from the Komo-Ise Cohort Study Journal of Epidemiology, 2002, 12, 93-104.	2.4	36
58	Generalizability of Relative Risk Estimates from a Well-defined Population to a General Population. European Journal of Epidemiology, 2006, 21, 253-262.	5.7	36
59	Plasma levels of C-reactive protein and serum amyloid A and gastric cancer in a nested case-control study: Japan Public Health Center-based prospective study. Carcinogenesis, 2010, 31, 712-718.	2.8	36
60	Association Between Plasma 25-Hydroxyvitamin D and Colorectal Adenoma According to Dietary Calcium Intake and Vitamin D Receptor Polymorphism. American Journal of Epidemiology, 2012, 175, 236-244.	3.4	35
61	Validating the dietary inflammatory index using inflammatory biomarkers in a Japanese population: A cross-sectional study of the JPHC-FFQ validation study. Nutrition, 2020, 69, 110569.	2.4	35
62	Isoflavone, polymorphisms in estrogen receptor genes and breast cancer risk in case ontrol studies in Japanese, Japanese Brazilians and nonâ€Japanese Brazilians. Cancer Science, 2009, 100, 927-933.	3.9	34
63	Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of colorectal cancer: A hospital-based case-control study in Japan. Scandinavian Journal of Gastroenterology, 2009, 44, 952-959.	1.5	34
64	Cruciferous Vegetable Intake Is Inversely Associated with Lung Cancer Risk among Current Nonsmoking Men in the Japan Public Health Center (JPHC) Study. Journal of Nutrition, 2017, 147, 841-849.	2.9	34
65	Plasma testosterone and sex hormoneâ€binding globulin concentrations and the risk of prostate cancer among Japanese men: A nested caseâ€control study. Cancer Science, 2010, 101, 2652-2657.	3.9	31
66	The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. Journal of Epidemiology, 2020, 30, 46-54.	2.4	30
67	Screening of 214 Single Nucleotide Polymorphisms in 44 Candidate Cancer Susceptibility Genes: A Case–Control Study on Gastric and Colorectal Cancers in the Japanese Population. American Journal of Gastroenterology, 2008, 103, 1476-1487.	0.4	29
68	Secular trends in cancer mortality among Japanese immigrants in the state of São Paulo, Brazil, 1979–2001. European Journal of Cancer Prevention, 2008, 17, 1-8.	1.3	28
69	Death by Suicide and Other Externally Caused Injuries After Stroke in Japan (1990–2010). Psychosomatic Medicine, 2014, 76, 452-459.	2.0	28
70	Dietary consumption of antioxidant vitamins and subsequent lung cancer risk: The <scp>J</scp> apan <scp>P</scp> ublic <scp>H</scp> ealth <scp>C</scp> enterâ€based prospective study. International Journal of Cancer, 2018, 142, 2441-2460.	5.1	28
71	Low carbohydrate diet and all cause and cause-specific mortality. Clinical Nutrition, 2021, 40, 2016-2024.	5.0	28
72	Plasma folate and risk of colorectal cancer in a nested case-control study: the Japan Public Health Center-based prospective study. Cancer Causes and Control, 2008, 19, 67-74.	1.8	27

#	Article	IF	CITATIONS
73	Reproductive factors, exogenous female hormone use and colorectal cancer risk: the Japan Public Health Center-based Prospective Study. European Journal of Cancer Prevention, 2008, 17, 515-524.	1.3	27
74	Methionine Synthase A2756G Polymorphism Interacts with Alcohol and Folate Intake to Influence the Risk of Colorectal Adenoma. Cancer Epidemiology Biomarkers and Prevention, 2009, 18, 267-274.	2.5	27
75	Impact of viral load of hepatitis C on the incidence of hepatocellular carcinoma: A population-based cohort study (JPHC Study). Cancer Letters, 2011, 300, 173-179.	7.2	26
76	Determination of Sub-ppb Cadmium in Urine by Solid-Phase Extraction and Inductively Coupled Plasma-Mass Spectrometry. Analytical Sciences, 2008, 24, 1049-1052.	1.6	25
77	Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of stomach cancer: a hospital-based case-control study in Japan. Gastric Cancer, 2009, 12, 198-205.	5.3	25
78	Dietary Heterocyclic Amine Intake, <i>NAT2</i> Genetic Polymorphism, and Colorectal Adenoma Risk: The Colorectal Adenoma Study in Tokyo. Cancer Epidemiology Biomarkers and Prevention, 2015, 24, 613-620.	2.5	25
79	Body mass index and colorectal cancer risk: A Mendelian randomization study. Cancer Science, 2021, 112, 1579-1588.	3.9	25
80	Dietary patterns and prostate cancer risk in Japanese: the Japan Public Health Center-based Prospective Study (JPHC Study). Cancer Causes and Control, 2018, 29, 589-600.	1.8	23
81	Dietary Inflammatory Index Is Associated With Inflammation in Japanese Men. Frontiers in Nutrition, 2021, 8, 604296.	3.7	23
82	Leisure-time physical activity and breast cancer risk by hormone receptor status: effective life periods and exercise intensity. Cancer Causes and Control, 2010, 21, 1787-1798.	1.8	22
83	Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. Cancer Prevention Research, 2017, 10, 535-541.	1.5	21
84	<i>CYP1A1</i> , <i>GSTM1</i> and <i>GSTT1</i> genetic polymorphisms and gastric cancer risk among Japanese: A nested case–control study within a largeâ€scale populationâ€based prospective study. International Journal of Cancer, 2016, 139, 759-768.	5.1	20
85	Association between NAT2, CYP1A1, and CYP1A2 genotypes, heterocyclic aromatic amines, and prostate cancer risk: a case control study in Japan. Environmental Health and Preventive Medicine, 2017, 22, 72.	3.4	20
86	Dietary patterns and colorectal cancer risk in middle-aged adults: AÂlarge population-based prospective cohort study. Clinical Nutrition, 2018, 37, 1019-1026.	5.0	20
87	Reproductive history and risk of cognitive impairment in Japanese women. Maturitas, 2019, 128, 22-28.	2.4	20
88	Dietary Isoflavone Intake, Polymorphisms in the CYP17, CYP19, 17β-HSD1, and SHBG Genes, and Risk of Breast Cancer in Case-Control Studies in Japanese, Japanese Brazilians, and Non-Japanese Brazilians. Nutrition and Cancer, 2010, 62, 466-475.	2.0	19
89	High-Negative Anti– <i>Helicobacter pylori</i> IgG Antibody Titers and Long-Term Risk of Gastric Cancer: Results from a Large-Scale Population-Based Cohort Study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2020, 29, 420-426.	2.5	19
90	Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. European Journal of Clinical Nutrition, 2021, 75, 954-968.	2.9	19

#	Article	IF	CITATIONS
91	Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. European Journal of Cancer Prevention, 2018, 27, 171-179.	1.3	19
92	Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. Atherosclerosis, 2018, 272, 226-232.	0.8	18
93	Lowâ€carbohydrate diet and risk of cancer incidence: The Japan Public Health Centerâ€based prospective study. Cancer Science, 2022, 113, 744-755.	3.9	17
94	Development of a quantitative food frequency questionnaire for assessing food, nutrient, and heterocyclic aromatic amines intake in Japanese Brazilians for a colorectal adenoma case–control study. International Journal of Food Sciences and Nutrition, 2009, 60, 128-139.	2.8	16
95	Female reproductive factors and risk of all-cause and cause-specific mortality among women: The Japan Public Health Center–based Prospective Study (JPHC study). Annals of Epidemiology, 2018, 28, 597-604.e6.	1.9	16
96	Plasma Isoflavone Concentrations Are Not Associated with Gastric Cancer Risk among Japanese Men and Women1,2. Journal of Nutrition, 2013, 143, 1293-1298.	2.9	15
97	The association between complete and partial non-response to psychosocial questions and suicide: the JPHC Study. European Journal of Public Health, 2015, 25, 424-430.	0.3	14
98	Alcohol consumption, genetic variants in the alcohol- and folate metabolic pathways and colorectal cancer risk: the JPHC Study. Scientific Reports, 2016, 6, 36607.	3.3	14
99	Body mass index change during adulthood and risk of oesophageal squamous-cell carcinoma in a Japanese population: the Japan Public Health (JPHC)-based prospective study. British Journal of Cancer, 2017, 117, 1715-1722.	6.4	14
100	Epidemiology of nonmelanoma skin cancer in Japan: Occupational type, lifestyle, and family history of cancer. Cancer Science, 2020, 111, 4257-4265.	3.9	14
101	Absence of specific symptoms in chronic hepatitis C. Journal of Gastroenterology, 2002, 37, 709-716.	5.1	13
102	Validity of a self-administered food frequency questionnaire in the estimation of heterocyclic aromatic amines. Cancer Causes and Control, 2014, 25, 1015-1028.	1.8	13
103	Association of <i>Escherichia coli</i> containing polyketide synthase in the gut microbiota with colorectal neoplasia in Japan. Cancer Science, 2022, 113, 277-286.	3.9	13
104	Glycemic index and glycemic load and risk of colorectal cancer: a population-based cohort study (JPHC Study). Cancer Causes and Control, 2016, 27, 583-593.	1.8	12
105	Validity and Reproducibility of a Self-Administered Food Frequency Questionnaire for the Assessment of Sugar Intake in Middle-Aged Japanese Adults. Nutrients, 2019, 11, 554.	4.1	12
106	Occupational sitting time and subsequent risk of cancer: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2020, 111, 974-984.	3.9	11
107	Impact of five modifiable lifestyle habits on the probability of cancer occurrence in a Japanese population-based cohort: Results from the JPHC study. Preventive Medicine, 2013, 57, 685-689.	3.4	10
108	Comparison between the impact of fermented and unfermented soy intake on the risk of liver cancer: the JPHC Study. European Journal of Nutrition, 2021, 60, 1389-1401.	3.9	10

#	Article	IF	CITATIONS
109	Reproductive Factors and Lung Cancer Risk among Never-Smoking Japanese Women with 21 Years of Follow-Up: A Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1185-1192.	2.5	10
110	Association between meat and saturated fatty acid intake and lung cancer risk: The Japan Public Health Centerâ€based prospective study. International Journal of Cancer, 2020, 147, 3019-3028.	5.1	10
111	Longâ€ŧerm antihypertensive drug use and risk of cancer: The Japan Public Health Centerâ€based prospective study. Cancer Science, 2021, 112, 1997-2005.	3.9	9
112	Higher Dietary Non-enzymatic Antioxidant Capacity Is Associated with Decreased Risk of All-Cause and Cardiovascular Disease Mortality in Japanese Adults. Journal of Nutrition, 2019, 149, 1967-1976.	2.9	8
113	Relationship between dietary non-enzymatic antioxidant capacity and type 2 diabetes risk in the Japan Public Health Center-based Prospective Study. Nutrition, 2019, 66, 62-69.	2.4	8
114	Doneness preferences, meat and meat-derived heterocyclic amines intake, and N-acetyltransferase 2 polymorphisms: association with colorectal adenoma in Japanese Brazilians. European Journal of Cancer Prevention, 2020, 29, 7-14.	1.3	8
115	Inclusion of a geneâ€environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. Cancer Science, 2020, 111, 3835-3844.	3.9	8
116	Dietary fiber intake and risk of gastric cancer: The <scp>Japan Public Health Center</scp> â€based prospective study. International Journal of Cancer, 2021, 148, 2664-2673.	5.1	8
117	Sugary drink consumption and risk of kidney and bladder cancer in Japanese adults. Scientific Reports, 2021, 11, 21701.	3.3	8
118	Association between dietary sugar intake and colorectal adenoma among cancer screening examinees in Japan. Cancer Science, 2020, 111, 3862-3872.	3.9	7
119	Sugary Drink Consumption and Subsequent Colorectal Cancer Risk: The Japan Public Health Center–Based Prospective Cohort Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 782-788.	2.5	7
120	Alcohol consumption, tobacco smoking, and subsequent risk of renal cell carcinoma: The JPHC study. Cancer Science, 2021, 112, 5068-5077.	3.9	7
121	Low Back Pain and Smoking in a Community Sample in Japan. Journal of Occupational Health, 2002, 44, 207-213.	2.1	6
122	Cruciferous vegetable intake and colorectal cancer risk: Japan public health center-based prospective study. European Journal of Cancer Prevention, 2019, 28, 420-427.	1.3	6
123	Dietary heterocyclic aromatic amine intake and cancer risk: epidemiological evidence from Japanese studies. Genes and Environment, 2021, 43, 33.	2.1	6
124	Meat consumption and gastric cancer risk: The Japan Public Health Center-based Prospective Study. American Journal of Clinical Nutrition, 2021, , .	4.7	6
125	Inverse Association between Fruit and Vegetable Intake and All-Cause Mortality: Japan Public Health Center-Based Prospective Study. Journal of Nutrition, 2022, 152, 2245-2254.	2.9	6
126	Prediagnostic plasma polyphenol concentrations and colon cancer risk: The JPHC nested case–control study. Clinical Nutrition, 2022, 41, 1950-1960.	5.0	6

#	Article	IF	CITATIONS
127	Association between polymorphisms in glutathione S-transferase Mu3 and IgG titer levels in serum against Helicobacter pylori. Journal of Human Genetics, 2009, 54, 557-563.	2.3	5
128	Plasma Câ€peptide and glycated albumin and subsequent risk of cancer: From a large prospective caseâ€cohort study in Japan. International Journal of Cancer, 2019, 144, 718-729.	5.1	5
129	Dietary glycemic index, glycemic load, and endometrial cancer risk: The Japan Public Health Centerâ€based Prospective Study. Cancer Science, 2021, 112, 3682-3690.	3.9	5
130	Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. Preventive Medicine, 2021, 148, 106561.	3.4	5
131	The Validity and Reproducibility of Dietary Non-enzymatic Antioxidant Capacity Estimated by Self-administered Food Frequency Questionnaires. Journal of Epidemiology, 2018, 28, 428-436.	2.4	4
132	Validation Study of Diabetes Definitions Using Japanese Diagnosis Procedure Combination Data Among Hospitalized Patients. Journal of Epidemiology, 2023, 33, 165-169.	2.4	4
133	Risk Stratification Score Improves Sensitivity for Advanced Colorectal Neoplasia in Colorectal Cancer Screening: The Oshima Study Workgroup. Clinical and Translational Gastroenterology, 2021, 12, e00319.	2.5	4
134	Body Mass Index, Height, Weight Change, and Subsequent Lung Cancer Risk: The Japan Public Health Center–Based Prospective Study. Cancer Epidemiology Biomarkers and Prevention, 2021, 30, 1708-1716.	2.5	4
135	The association between plasma C-peptide concentration and the risk of prostate cancer: a nested case–control study within a Japanese population-based prospective study. European Journal of Cancer Prevention, 2018, 27, 461-467.	1.3	3
136	Female reproductive factors and risk of external causes of death among women: The Japan Public Health Center-based Prospective Study (JPHC Study). Scientific Reports, 2019, 9, 14329.	3.3	3
137	Exploratory Research on Determinants of Place of Death in a Large-scale Cohort Study: The JPHC Study. Journal of Epidemiology, 2023, 33, 120-126.	2.4	3
138	Association between coffee consumption and risk of prostate cancer in Japanese men: a population-based cohort study in Japan. Cancer Epidemiology Biomarkers and Prevention, 2021, , cebp.0484.2021.	2.5	3
139	Association of B Vitamins and Methionine Intake with the Risk of Gastric Cancer: The Japan Public Health Center–based Prospective Study. Cancer Prevention Research, 2022, 15, 101-110.	1.5	3
140	Association between Meat, Fish, and Fatty Acid Intake and Non-Hodgkin Lymphoma Incidence: The Japan Public Health Center–Based Prospective Study. Journal of Nutrition, 2022, 152, 1895-1906.	2.9	3
141	Dietary fibre intake is associated with reduced risk of lung cancer: a Japan public health centre-based prospective study (JPHC). International Journal of Epidemiology, 2022, 51, 1142-1152.	1.9	2
142	Commentary: Factors Associated With Non-participation in Cohort Studies Emphasize the Need to Generalize the Results With Care. Journal of Epidemiology, 2015, 25, 89-90.	2.4	1
143	Total, animal, and plant protein intake and pneumonia mortality in the Japan Public Health Center–based Prospective Study. American Journal of Clinical Nutrition, 2022, 115, 781-789.	4.7	1
144	Association of Plasma Iron Status with Subsequent Risk of Total and Site-Specific Cancer: A Large Case–Cohort Study within JPHC Study. Cancer Prevention Research, 2022, 15, 669-678.	1.5	1

Motok	NUACAR/

#	Article	IF	CITATIONS
145	Public access to summary statistics for genome-wide association studies of body mass index, weight, and height among healthy Japanese individuals: the Japanese Consortium of Genetic Epidemiology studies. Journal of Epidemiology, 2021, , .	2.4	0
146	Title is missing!. , 2020, 15, e0244007.		0
147	Title is missing!. , 2020, 15, e0244007.		0
148	Title is missing!. , 2020, 15, e0244007.		0
149	Title is missing!. , 2020, 15, e0244007.		0