

Motoki Iwasaki

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8769656/publications.pdf>

Version: 2024-02-01

149
papers

6,061
citations

66250

44
h-index

100535

70
g-index

150
all docs

150
docs citations

150
times ranked

10034
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Validation Study of Diabetes Definitions Using Japanese Diagnosis Procedure Combination Data Among Hospitalized Patients. <i>Journal of Epidemiology</i> , 2023, 33, 165-169. | 1.1 | 4 |
| 2 | Exploratory Research on Determinants of Place of Death in a Large-scale Cohort Study: The JPHC Study. <i>Journal of Epidemiology</i> , 2023, 33, 120-126. | 1.1 | 3 |
| 3 | Low-carbohydrate diet and risk of cancer incidence: The Japan Public Health Center-based prospective study. <i>Cancer Science</i> , 2022, 113, 744-755. | 1.7 | 17 |
| 4 | Association of B Vitamins and Methionine Intake with the Risk of Gastric Cancer: The Japan Public Health Center-based Prospective Study. <i>Cancer Prevention Research</i> , 2022, 15, 101-110. | 0.7 | 3 |
| 5 | Association of <i>Escherichia coli</i> containing polyketide synthase in the gut microbiota with colorectal neoplasia in Japan. <i>Cancer Science</i> , 2022, 113, 277-286. | 1.7 | 13 |
| 6 | Dietary fibre intake is associated with reduced risk of lung cancer: a Japan public health centre-based prospective study (JPHC). <i>International Journal of Epidemiology</i> , 2022, 51, 1142-1152. | 0.9 | 2 |
| 7 | Total, animal, and plant protein intake and pneumonia mortality in the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2022, 115, 781-789. | 2.2 | 1 |
| 8 | Association between Meat, Fish, and Fatty Acid Intake and Non-Hodgkin Lymphoma Incidence: The Japan Public Health Center-Based Prospective Study. <i>Journal of Nutrition</i> , 2022, 152, 1895-1906. | 1.3 | 3 |
| 9 | Inverse Association between Fruit and Vegetable Intake and All-Cause Mortality: Japan Public Health Center-Based Prospective Study. <i>Journal of Nutrition</i> , 2022, 152, 2245-2254. | 1.3 | 6 |
| 10 | Prediagnostic plasma polyphenol concentrations and colon cancer risk: The JPHC nested case-control study. <i>Clinical Nutrition</i> , 2022, 41, 1950-1960. | 2.3 | 6 |
| 11 | Association of Plasma Iron Status with Subsequent Risk of Total and Site-Specific Cancer: A Large Case-Cohort Study within JPHC Study. <i>Cancer Prevention Research</i> , 2022, 15, 669-678. | 0.7 | 1 |
| 12 | Comparison between the impact of fermented and unfermented soy intake on the risk of liver cancer: the JPHC Study. <i>European Journal of Nutrition</i> , 2021, 60, 1389-1401. | 1.8 | 10 |
| 13 | Low carbohydrate diet and all cause and cause-specific mortality. <i>Clinical Nutrition</i> , 2021, 40, 2016-2024. | 2.3 | 28 |
| 14 | Dietary fiber intake and risk of gastric cancer: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2021, 148, 2664-2673. | 2.3 | 8 |
| 15 | Fermented soy products intake and risk of cardiovascular disease and total cancer incidence: The Japan Public Health Center-based Prospective study. <i>European Journal of Clinical Nutrition</i> , 2021, 75, 954-968. | 1.3 | 19 |
| 16 | Sugary Drink Consumption and Subsequent Colorectal Cancer Risk: The Japan Public Health Center-Based Prospective Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, 30, 782-788. | 1.1 | 7 |
| 17 | Body mass index and colorectal cancer risk: A Mendelian randomization study. <i>Cancer Science</i> , 2021, 112, 1579-1588. | 1.7 | 25 |
| 18 | Risk Stratification Score Improves Sensitivity for Advanced Colorectal Neoplasia in Colorectal Cancer Screening: The Oshima Study Workgroup. <i>Clinical and Translational Gastroenterology</i> , 2021, 12, e00319. | 1.3 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Long-term antihypertensive drug use and risk of cancer: The Japan Public Health Center-based prospective study. <i>Cancer Science</i> , 2021, 112, 1997-2005. | 1.7 | 9 |
| 20 | Reproductive Factors and Lung Cancer Risk among Never-Smoking Japanese Women with 21 Years of Follow-Up: A Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, 30, 1185-1192. | 1.1 | 10 |
| 21 | Dietary Inflammatory Index Is Associated With Inflammation in Japanese Men. <i>Frontiers in Nutrition</i> , 2021, 8, 604296. | 1.6 | 23 |
| 22 | Body Mass Index, Height, Weight Change, and Subsequent Lung Cancer Risk: The Japan Public Health Center-based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, 30, 1708-1716. | 1.1 | 4 |
| 23 | Dietary glycemic index, glycemic load, and endometrial cancer risk: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2021, 112, 3682-3690. | 1.7 | 5 |
| 24 | Association of sugary drink consumption with all-cause and cause-specific mortality: the Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2021, 148, 106561. | 1.6 | 5 |
| 25 | Dietary heterocyclic aromatic amine intake and cancer risk: epidemiological evidence from Japanese studies. <i>Genes and Environment</i> , 2021, 43, 33. | 0.9 | 6 |
| 26 | Alcohol consumption, tobacco smoking, and subsequent risk of renal cell carcinoma: The JPHC study. <i>Cancer Science</i> , 2021, 112, 5068-5077. | 1.7 | 7 |
| 27 | Sugary drink consumption and risk of kidney and bladder cancer in Japanese adults. <i>Scientific Reports</i> , 2021, 11, 21701. | 1.6 | 8 |
| 28 | Association between coffee consumption and risk of prostate cancer in Japanese men: a population-based cohort study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, , cebp.0484.2021. | 1.1 | 3 |
| 29 | Meat consumption and gastric cancer risk: The Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2021, , . | 2.2 | 6 |
| 30 | Public access to summary statistics for genome-wide association studies of body mass index, weight, and height among healthy Japanese individuals: the Japanese Consortium of Genetic Epidemiology studies. <i>Journal of Epidemiology</i> , 2021, , . | 1.1 | 0 |
| 31 | The Japan Public Health Center-based Prospective Study for the Next Generation (JPHC-NEXT): Study Design and Participants. <i>Journal of Epidemiology</i> , 2020, 30, 46-54. | 1.1 | 30 |
| 32 | Diabetes and cancer risk: A Mendelian randomization study. <i>International Journal of Cancer</i> , 2020, 146, 712-719. | 2.3 | 52 |
| 33 | Validating the dietary inflammatory index using inflammatory biomarkers in a Japanese population: A cross-sectional study of the JPHC-FFQ validation study. <i>Nutrition</i> , 2020, 69, 110569. | 1.1 | 35 |
| 34 | High-Negative Anti- <i>Helicobacter pylori</i> IgG Antibody Titers and Long-Term Risk of Gastric Cancer: Results from a Large-Scale Population-Based Cohort Study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020, 29, 420-426. | 1.1 | 19 |
| 35 | Doneness preferences, meat and meat-derived heterocyclic amines intake, and N-acetyltransferase 2 polymorphisms: association with colorectal adenoma in Japanese Brazilians. <i>European Journal of Cancer Prevention</i> , 2020, 29, 7-14. | 0.6 | 8 |
| 36 | Epidemiology of nonmelanoma skin cancer in Japan: Occupational type, lifestyle, and family history of cancer. <i>Cancer Science</i> , 2020, 111, 4257-4265. | 1.7 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Association between dietary sugar intake and colorectal adenoma among cancer screening examinees in Japan. <i>Cancer Science</i> , 2020, 111, 3862-3872. | 1.7 | 7 |
| 38 | Inclusion of a gene-environment interaction between alcohol consumption and the aldehyde dehydrogenase 2 genotype in a risk prediction model for upper aerodigestive tract cancer in Japanese men. <i>Cancer Science</i> , 2020, 111, 3835-3844. | 1.7 | 8 |
| 39 | Occupational sitting time and subsequent risk of cancer: The Japan Public Health Center-based Prospective Study. <i>Cancer Science</i> , 2020, 111, 974-984. | 1.7 | 11 |
| 40 | Association of soy and fermented soy product intake with total and cause specific mortality: prospective cohort study. <i>BMJ, The</i> , 2020, 368, m34. | 3.0 | 45 |
| 41 | Dietary fiber intake and total and cause-specific mortality: the Japan Public Health Center-based prospective study. <i>American Journal of Clinical Nutrition</i> , 2020, 111, 1027-1035. | 2.2 | 38 |
| 42 | Association between meat and saturated fatty acid intake and lung cancer risk: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2020, 147, 3019-3028. | 2.3 | 10 |
| 43 | Title is missing!. , 2020, 15, e0244007. | | 0 |
| 44 | Title is missing!. , 2020, 15, e0244007. | | 0 |
| 45 | Title is missing!. , 2020, 15, e0244007. | | 0 |
| 46 | Title is missing!. , 2020, 15, e0244007. | | 0 |
| 47 | Reproductive history and risk of cognitive impairment in Japanese women. <i>Maturitas</i> , 2019, 128, 22-28. | 1.0 | 20 |
| 48 | Association of Animal and Plant Protein Intake With All-Cause and Cause-Specific Mortality in a Japanese Cohort. <i>JAMA Internal Medicine</i> , 2019, 179, 1509. | 2.6 | 120 |
| 49 | Higher Dietary Non-enzymatic Antioxidant Capacity Is Associated with Decreased Risk of All-Cause and Cardiovascular Disease Mortality in Japanese Adults. <i>Journal of Nutrition</i> , 2019, 149, 1967-1976. | 1.3 | 8 |
| 50 | Characterizing rare and low-frequency height-associated variants in the Japanese population. <i>Nature Communications</i> , 2019, 10, 4393. | 5.8 | 123 |
| 51 | Female reproductive factors and risk of external causes of death among women: The Japan Public Health Center-based Prospective Study (JPHC Study). <i>Scientific Reports</i> , 2019, 9, 14329. | 1.6 | 3 |
| 52 | Validity and Reproducibility of a Self-Administered Food Frequency Questionnaire for the Assessment of Sugar Intake in Middle-Aged Japanese Adults. <i>Nutrients</i> , 2019, 11, 554. | 1.7 | 12 |
| 53 | Relationship between dietary non-enzymatic antioxidant capacity and type 2 diabetes risk in the Japan Public Health Center-based Prospective Study. <i>Nutrition</i> , 2019, 66, 62-69. | 1.1 | 8 |
| 54 | Cruciferous vegetable intake and colorectal cancer risk: Japan public health center-based prospective study. <i>European Journal of Cancer Prevention</i> , 2019, 28, 420-427. | 0.6 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Plasma C-peptide and glycated albumin and subsequent risk of cancer: From a large prospective case-cohort study in Japan. <i>International Journal of Cancer</i> , 2019, 144, 718-729. | 2.3 | 5 |
| 56 | GWAS identifies two novel colorectal cancer loci at 16q24.1 and 20q13.12. <i>Carcinogenesis</i> , 2018, 39, 652-660. | 1.3 | 52 |
| 57 | Dietary consumption of antioxidant vitamins and subsequent lung cancer risk: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2018, 142, 2441-2460. | 2.3 | 28 |
| 58 | Plasma levels of n-3 fatty acids and risk of coronary heart disease among Japanese: The Japan Public Health Center-based (JPHC) study. <i>Atherosclerosis</i> , 2018, 272, 226-232. | 0.4 | 18 |
| 59 | Plasma 25-hydroxyvitamin D concentration and subsequent risk of total and site specific cancers in Japanese population: large case-cohort study within Japan Public Health Center-based Prospective Study cohort. <i>BMJ: British Medical Journal</i> , 2018, 360, k671. | 2.4 | 61 |
| 60 | Dietary patterns and prostate cancer risk in Japanese: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Cancer Causes and Control</i> , 2018, 29, 589-600. | 0.8 | 23 |
| 61 | Dietary patterns and colorectal cancer risk in middle-aged adults: A large population-based prospective cohort study. <i>Clinical Nutrition</i> , 2018, 37, 1019-1026. | 2.3 | 20 |
| 62 | The association between plasma C-peptide concentration and the risk of prostate cancer: a nested case-control study within a Japanese population-based prospective study. <i>European Journal of Cancer Prevention</i> , 2018, 27, 461-467. | 0.6 | 3 |
| 63 | Increased Levels of Branched-Chain Amino Acid Associated With Increased Risk of Pancreatic Cancer in a Prospective Case-Control Study of a Large Cohort. <i>Gastroenterology</i> , 2018, 155, 1474-1482.e1. | 0.6 | 59 |
| 64 | The Validity and Reproducibility of Dietary Non-enzymatic Antioxidant Capacity Estimated by Self-administered Food Frequency Questionnaires. <i>Journal of Epidemiology</i> , 2018, 28, 428-436. | 1.1 | 4 |
| 65 | Female reproductive factors and risk of all-cause and cause-specific mortality among women: The Japan Public Health Center-based Prospective Study (JPHC study). <i>Annals of Epidemiology</i> , 2018, 28, 597-604.e6. | 0.9 | 16 |
| 66 | Cigarette smoking, alcohol drinking, and oral cavity and pharyngeal cancer in the Japanese: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2018, 27, 171-179. | 0.6 | 19 |
| 67 | Cruciferous Vegetable Intake Is Inversely Associated with Lung Cancer Risk among Current Nonsmoking Men in the Japan Public Health Center (JPHC) Study. <i>Journal of Nutrition</i> , 2017, 147, 841-849. | 1.3 | 34 |
| 68 | Body mass index change during adulthood and risk of oesophageal squamous-cell carcinoma in a Japanese population: the Japan Public Health (JPHC)-based prospective study. <i>British Journal of Cancer</i> , 2017, 117, 1715-1722. | 2.9 | 14 |
| 69 | Genome-wide association study identifies 112 new loci for body mass index in the Japanese population. <i>Nature Genetics</i> , 2017, 49, 1458-1467. | 9.4 | 380 |
| 70 | Inclusion of a Genetic Risk Score into a Validated Risk Prediction Model for Colorectal Cancer in Japanese Men Improves Performance. <i>Cancer Prevention Research</i> , 2017, 10, 535-541. | 0.7 | 21 |
| 71 | Fermented Soy Product Intake Is Inversely Associated with the Development of High Blood Pressure: The Japan Public Health Center-Based Prospective Study. <i>Journal of Nutrition</i> , 2017, 147, 1749-1756. | 1.3 | 51 |
| 72 | Association between NAT2, CYP1A1, and CYP1A2 genotypes, heterocyclic aromatic amines, and prostate cancer risk: a case control study in Japan. <i>Environmental Health and Preventive Medicine</i> , 2017, 22, 72. | 1.4 | 20 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Prediction of the 10-year probability of gastric cancer occurrence in the Japanese population: the JPHC study cohort II. <i>International Journal of Cancer</i> , 2016, 138, 320-331. | 2.3 | 78 |
| 74 | <i>CYP1A1</i> , <i>GSTM1</i> and <i>GSTT1</i> genetic polymorphisms and gastric cancer risk among Japanese: A nested case-control study within a large-scale population-based prospective study. <i>International Journal of Cancer</i> , 2016, 139, 759-768. | 2.3 | 20 |
| 75 | Alcohol consumption, genetic variants in the alcohol- and folate metabolic pathways and colorectal cancer risk: the JPHC Study. <i>Scientific Reports</i> , 2016, 6, 36607. | 1.6 | 14 |
| 76 | Glycemic index and glycemic load and risk of colorectal cancer: a population-based cohort study (JPHC Study). <i>Cancer Causes and Control</i> , 2016, 27, 583-593. | 0.8 | 12 |
| 77 | Commentary: Factors Associated With Non-participation in Cohort Studies Emphasize the Need to Generalize the Results With Care. <i>Journal of Epidemiology</i> , 2015, 25, 89-90. | 1.1 | 1 |
| 78 | Plasma insulin, C-peptide and blood glucose and the risk of gastric cancer: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2015, 136, 1402-1410. | 2.3 | 44 |
| 79 | The association between complete and partial non-response to psychosocial questions and suicide: the JPHC Study. <i>European Journal of Public Health</i> , 2015, 25, 424-430. | 0.1 | 14 |
| 80 | Association of coffee intake with total and cause-specific mortality in a Japanese population: the Japan Public Health Center-based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2015, 101, 1029-1037. | 2.2 | 58 |
| 81 | Dietary Heterocyclic Amine Intake, <i>NAT2</i> Genetic Polymorphism, and Colorectal Adenoma Risk: The Colorectal Adenoma Study in Tokyo. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2015, 24, 613-620. | 1.1 | 25 |
| 82 | Association of green tea consumption with mortality due to all causes and major causes of death in a Japanese population: the Japan Public Health Center-based Prospective Study (JPHC Study). <i>Annals of Epidemiology</i> , 2015, 25, 512-518.e3. | 0.9 | 66 |
| 83 | Genetic polymorphisms of <i>ADH1B</i> , <i>ADH1C</i> and <i>ALDH2</i> , alcohol consumption, and the risk of gastric cancer: the Japan Public Health Center-based prospective study. <i>Carcinogenesis</i> , 2015, 36, 223-231. | 1.3 | 69 |
| 84 | Death by suicide and other externally caused injuries following a cancer diagnosis: the Japan Public Health Center-based Prospective Study. <i>Psycho-Oncology</i> , 2014, 23, 1034-1041. | 1.0 | 45 |
| 85 | Death by Suicide and Other Externally Caused Injuries After Stroke in Japan (1990-2010). <i>Psychosomatic Medicine</i> , 2014, 76, 452-459. | 1.3 | 28 |
| 86 | Alcohol and smoking and subsequent risk of prostate cancer in Japanese men: The Japan Public Health Center-based prospective study. <i>International Journal of Cancer</i> , 2014, 134, 971-978. | 2.3 | 52 |
| 87 | Validity of a self-administered food frequency questionnaire in the estimation of heterocyclic aromatic amines. <i>Cancer Causes and Control</i> , 2014, 25, 1015-1028. | 0.8 | 13 |
| 88 | Dietary cadmium intake and breast cancer risk in Japanese women: A case-control study. <i>International Journal of Hygiene and Environmental Health</i> , 2014, 217, 70-77. | 2.1 | 115 |
| 89 | Impact of five modifiable lifestyle habits on the probability of cancer occurrence in a Japanese population-based cohort: Results from the JPHC study. <i>Preventive Medicine</i> , 2013, 57, 685-689. | 1.6 | 10 |
| 90 | Social support and cancer incidence and mortality: the JPHC study cohort II. <i>Cancer Causes and Control</i> , 2013, 24, 847-860. | 0.8 | 68 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Plasma Isoflavone Concentrations Are Not Associated with Gastric Cancer Risk among Japanese Men and Women ^{1,2} . <i>Journal of Nutrition</i> , 2013, 143, 1293-1298. | 1.3 | 15 |
| 92 | Association Between Plasma 25-Hydroxyvitamin D and Colorectal Adenoma According to Dietary Calcium Intake and Vitamin D Receptor Polymorphism. <i>American Journal of Epidemiology</i> , 2012, 175, 236-244. | 1.6 | 35 |
| 93 | Long-term Dietary Cadmium Intake and Cancer Incidence. <i>Epidemiology</i> , 2012, 23, 368-376. | 1.2 | 58 |
| 94 | Risk factors for epithelial ovarian cancer in Japan - results from the Japan Public Health Center-based Prospective Study cohort. <i>International Journal of Oncology</i> , 2011, 40, 21-30. | 1.4 | 39 |
| 95 | Impact of viral load of hepatitis C on the incidence of hepatocellular carcinoma: A population-based cohort study (JPHC Study). <i>Cancer Letters</i> , 2011, 300, 173-179. | 3.2 | 26 |
| 96 | Validity of a Self-Administered Food Frequency Questionnaire for Middle-Aged Urban Cancer Screenings: Comparison With 4-Day Weighed Dietary Records. <i>Journal of Epidemiology</i> , 2011, 21, 447-458. | 1.1 | 46 |
| 97 | Risk factors for breast cancer: epidemiological evidence from Japanese studies. <i>Cancer Science</i> , 2011, 102, 1607-1614. | 1.7 | 53 |
| 98 | Leisure-time physical activity and breast cancer risk defined by estrogen and progesterone receptor status ^{â€”} The Japan Public Health Center-based Prospective Study. <i>Preventive Medicine</i> , 2011, 52, 227-233. | 1.6 | 37 |
| 99 | Red meat intake may increase the risk of colon cancer in Japanese, a population with relatively low red meat consumption. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2011, 20, 603-12. | 0.3 | 51 |
| 100 | 10-Year risk of colorectal cancer: Development and validation of a prediction model in middle-aged Japanese men. <i>Cancer Epidemiology</i> , 2010, 34, 534-541. | 0.8 | 56 |
| 101 | Plasma tea polyphenol levels and subsequent risk of breast cancer among Japanese women: a nested case ^{â€”} control study. <i>Breast Cancer Research and Treatment</i> , 2010, 124, 827-834. | 1.1 | 47 |
| 102 | Leisure-time physical activity and breast cancer risk by hormone receptor status: effective life periods and exercise intensity. <i>Cancer Causes and Control</i> , 2010, 21, 1787-1798. | 0.8 | 22 |
| 103 | Alcohol consumption ^{â€”} associated breast cancer incidence and potential effect modifiers: the Japan Public Health Center ^{â€”} based Prospective Study. <i>International Journal of Cancer</i> , 2010, 127, 685-695. | 2.3 | 40 |
| 104 | Heterocyclic amines content of meat and fish cooked by Brazilian methods. <i>Journal of Food Composition and Analysis</i> , 2010, 23, 61-69. | 1.9 | 74 |
| 105 | Plasma testosterone and sex hormone ^{â€”} binding globulin concentrations and the risk of prostate cancer among Japanese men: A nested case ^{â€”} control study. <i>Cancer Science</i> , 2010, 101, 2652-2657. | 1.7 | 31 |
| 106 | Consumption of sodium and salted foods in relation to cancer and cardiovascular disease: the Japan Public Health Center ^{â€”} based Prospective Study. <i>American Journal of Clinical Nutrition</i> , 2010, 91, 456-464. | 2.2 | 100 |
| 107 | Isoflavone intake and risk of lung cancer: a prospective cohort study in Japan. <i>American Journal of Clinical Nutrition</i> , 2010, 91, 722-728. | 2.2 | 77 |
| 108 | Plasma levels of C-reactive protein and serum amyloid A and gastric cancer in a nested case-control study: Japan Public Health Center-based prospective study. <i>Carcinogenesis</i> , 2010, 31, 712-718. | 1.3 | 36 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Dietary Isoflavone Intake, Polymorphisms in the CYP17, CYP19, 17 β -HSD1, and SHBG Genes, and Risk of Breast Cancer in Case-Control Studies in Japanese, Japanese Brazilians, and Non-Japanese Brazilians. <i>Nutrition and Cancer</i> , 2010, 62, 466-475. | 0.9 | 19 |
| 110 | Green tea drinking and subsequent risk of breast cancer in a population to based cohort of Japanese women. <i>Breast Cancer Research</i> , 2010, 12, R88. | 2.2 | 52 |
| 111 | Development of a quantitative food frequency questionnaire for assessing food, nutrient, and heterocyclic aromatic amines intake in Japanese Brazilians for a colorectal adenoma caseâ€“control study. <i>International Journal of Food Sciences and Nutrition</i> , 2009, 60, 128-139. | 1.3 | 16 |
| 112 | Association between polymorphisms in glutathione S-transferase Mu3 and IgG titer levels in serum against <i>Helicobacter pylori</i> . <i>Journal of Human Genetics</i> , 2009, 54, 557-563. | 1.1 | 5 |
| 113 | Methionine Synthase A2756G Polymorphism Interacts with Alcohol and Folate Intake to Influence the Risk of Colorectal Adenoma. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 267-274. | 1.1 | 27 |
| 114 | Effect of Coffee and Green Tea Consumption on the Risk of Liver Cancer: Cohort Analysis by Hepatitis Virus Infection Status. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2009, 18, 1746-1753. | 1.1 | 98 |
| 115 | Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of stomach cancer: a hospital-based case-control study in Japan. <i>Gastric Cancer</i> , 2009, 12, 198-205. | 2.7 | 25 |
| 116 | Dietary isoflavone intake and breast cancer risk in caseâ€“control studies in Japanese, Japanese Brazilians, and non-Japanese Brazilians. <i>Breast Cancer Research and Treatment</i> , 2009, 116, 401-411. | 1.1 | 39 |
| 117 | Serum organochlorines and breast cancer risk in Japanese women: a caseâ€“control study. <i>Cancer Causes and Control</i> , 2009, 20, 567-580. | 0.8 | 70 |
| 118 | Metabolic factors and subsequent risk of hepatocellular carcinoma by hepatitis virus infection status: a large-scale population-based cohort study of Japanese men and women (JPHC Study Cohort II). <i>Cancer Causes and Control</i> , 2009, 20, 741-750. | 0.8 | 48 |
| 119 | Isoflavone, polymorphisms in estrogen receptor genes and breast cancer risk in caseâ€“control studies in Japanese, Japanese Brazilians and nonâ€“Japanese Brazilians. <i>Cancer Science</i> , 2009, 100, 927-933. | 1.7 | 34 |
| 120 | Association between dietary heterocyclic amine levels, genetic polymorphisms of NAT2, CYP1A1, and CYP1A2 and risk of colorectal cancer: A hospital-based case-control study in Japan. <i>Scandinavian Journal of Gastroenterology</i> , 2009, 44, 952-959. | 0.6 | 34 |
| 121 | Impact of metabolic factors on subsequent cancer risk: results from a large-scale population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009, 18, 240-247. | 0.6 | 131 |
| 122 | Serum aminotransferase level and the risk of hepatocellular carcinoma: a population-based cohort study in Japan. <i>European Journal of Cancer Prevention</i> , 2009, 18, 26-32. | 0.6 | 43 |
| 123 | Plasma folate and risk of colorectal cancer in a nested case-control study: the Japan Public Health Center-based prospective study. <i>Cancer Causes and Control</i> , 2008, 19, 67-74. | 0.8 | 27 |
| 124 | Passive smoking and lung cancer in Japanese nonâ€“smoking women: A prospective study. <i>International Journal of Cancer</i> , 2008, 122, 653-657. | 2.3 | 81 |
| 125 | Daily Total Physical Activity Level and Premature Death in Men and Women: Results From a Large-Scale Population-Based Cohort Study in Japan (JPHC Study). <i>Annals of Epidemiology</i> , 2008, 18, 522-530. | 0.9 | 147 |
| 126 | Screening of 214 Single Nucleotide Polymorphisms in 44 Candidate Cancer Susceptibility Genes: A Caseâ€“Control Study on Gastric and Colorectal Cancers in the Japanese Population. <i>American Journal of Gastroenterology</i> , 2008, 103, 1476-1487. | 0.2 | 29 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Plasma Tea Polyphenols and Gastric Cancer Risk: A Case-Control Study Nested in a Large Population-Based Prospective Study in Japan. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2008, 17, 343-351. | 1.1 | 46 |
| 128 | Plasma levels of carotenoids, retinol and tocopherol and the risk of gastric cancer in Japan: a nested case-control study. <i>Carcinogenesis</i> , 2008, 29, 1042-1048. | 1.3 | 60 |
| 129 | Plasma Isoflavone Level and Subsequent Risk of Breast Cancer Among Japanese Women: A Nested Case-Control Study From the Japan Public Health Center-Based Prospective Study Group. <i>Journal of Clinical Oncology</i> , 2008, 26, 1677-1683. | 0.8 | 155 |
| 130 | Determination of Sub-ppb Cadmium in Urine by Solid-Phase Extraction and Inductively Coupled Plasma-Mass Spectrometry. <i>Analytical Sciences</i> , 2008, 24, 1049-1052. | 0.8 | 25 |
| 131 | Secular trends in cancer mortality among Japanese immigrants in the state of São Paulo, Brazil, 1979-2001. <i>European Journal of Cancer Prevention</i> , 2008, 17, 1-8. | 0.6 | 28 |
| 132 | Reproductive factors, exogenous female hormone use and colorectal cancer risk: the Japan Public Health Center-based Prospective Study. <i>European Journal of Cancer Prevention</i> , 2008, 17, 515-524. | 0.6 | 27 |
| 133 | Green Tea Consumption and Prostate Cancer Risk in Japanese Men: A Prospective Study. <i>American Journal of Epidemiology</i> , 2007, 167, 71-77. | 1.6 | 241 |
| 134 | Role and impact of menstrual and reproductive factors on breast cancer risk in Japan. <i>European Journal of Cancer Prevention</i> , 2007, 16, 116-123. | 0.6 | 54 |
| 135 | Body Size and Risk for Breast Cancer in Relation to Estrogen and Progesterone Receptor Status in Japan. <i>Annals of Epidemiology</i> , 2007, 17, 304-312. | 0.9 | 51 |
| 136 | Colorectal cancer screening using fecal occult blood test and subsequent risk of colorectal cancer: A prospective cohort study in Japan. <i>Cancer Detection and Prevention</i> , 2007, 31, 3-11. | 2.1 | 75 |
| 137 | Physical activity and risk of colorectal cancer in Japanese men and women: the Japan Public Health Center-based prospective Study. <i>Cancer Causes and Control</i> , 2007, 18, 199-209. | 0.8 | 88 |
| 138 | Generalizability of Relative Risk Estimates from a Well-defined Population to a General Population. <i>European Journal of Epidemiology</i> , 2006, 21, 253-262. | 2.5 | 36 |
| 139 | Serum triglycerides and colorectal adenoma in a case-control study among cancer screening examinees (Japan). <i>Cancer Causes and Control</i> , 2006, 17, 1245-1252. | 0.8 | 45 |
| 140 | Dietary fiber intake and subsequent risk of colorectal cancer: The Japan Public Health Center-Based Prospective Study. <i>International Journal of Cancer</i> , 2006, 119, 1475-1480. | 2.3 | 48 |
| 141 | Plasma C-Reactive Protein and Risk of Colorectal Cancer in a Nested Case-Control Study: Japan Public Health Center-Based Prospective Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006, 15, 690-695. | 1.1 | 94 |
| 142 | Diabetes Mellitus and the Risk of Cancer. <i>Archives of Internal Medicine</i> , 2006, 166, 1871. | 4.3 | 475 |
| 143 | Effect of Helicobacter pylori Infection Combined with CagA and Pepsinogen Status on Gastric Cancer Development among Japanese Men and Women: A Nested Case-Control Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2006, 15, 1341-1347. | 1.1 | 123 |
| 144 | Body Mass Index, Body Height, and Subsequent Risk of Colorectal Cancer in Middle-Aged and Elderly Japanese Men and Women: Japan Public Health Center-Based Prospective Study. <i>Cancer Causes and Control</i> , 2005, 16, 839-850. | 0.8 | 72 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Cigarette Smoking and Completed Suicide among Middle-aged Men: A Population-based Cohort Study in Japan. <i>Annals of Epidemiology</i> , 2005, 15, 286-292. | 0.9 | 61 |
| 146 | Background Characteristics of Basic Health Examination Participants: the JPHC Study Baseline Survey. <i>Journal of Epidemiology</i> , 2003, 13, 216-225. | 1.1 | 49 |
| 147 | Rural-urban Differences in Sociodemographic, Social Network and Lifestyle Factors Related to Mortality of Middle-aged Japanese Men from the Komo-Ise Cohort Study.. <i>Journal of Epidemiology</i> , 2002, 12, 93-104. | 1.1 | 36 |
| 148 | Low Back Pain and Smoking in a Community Sample in Japan. <i>Journal of Occupational Health</i> , 2002, 44, 207-213. | 1.0 | 6 |
| 149 | Absence of specific symptoms in chronic hepatitis C. <i>Journal of Gastroenterology</i> , 2002, 37, 709-716. | 2.3 | 13 |