Emmanuelle Peters

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8773730/publications.pdf

Version: 2024-02-01

135

all docs

132 6,978 43
papers citations h-index

135

docs citations

h-index g-index

135 5235
times ranked citing authors

66879

78

#	Article	IF	CITATIONS
1	Cognitive bias modification for paranoia (CBM-pa): a randomised controlled feasibility study in patients with distressing paranoid beliefs. Psychological Medicine, 2023, 53, 4614-4626.	2.7	6
2	Comparing psychotic experiences in low-and-middle-income-countries and high-income-countries with a focus on measurement invariance. Psychological Medicine, 2022, 52, 1509-1516.	2.7	16
3	Associations between responses to voices, distress and appraisals during daily life: an ecological validation of the cognitive behavioural model. Psychological Medicine, 2022, 52, 538-547.	2.7	5
4	Social isolation and psychosis: an investigation of social interactions and paranoia in daily life. European Archives of Psychiatry and Clinical Neuroscience, 2022, 272, 119-127.	1.8	22
5	The digital divide: factors impacting on uptake of remote therapy in a South London psychological therapy service for people with psychosis. Journal of Mental Health, 2022, 31, 825-832.	1.0	18
6	Multisite randomised controlled trial of trauma-focused cognitive behaviour therapy for psychosis to reduce post-traumatic stress symptoms in people with co-morbid post-traumatic stress disorder and psychosis, compared to treatment as usual: study protocol for the STAR (Study of Trauma And) Tj ETQq0 0 0 0	rg <mark>B7</mark> /Ove	rl8ck 10 Tf 5
7	Need for care, adversity exposure and perceived stress in clinical and healthy voice-hearers. Psychological Medicine, 2021, 51, 1944-1950.	2.7	5
8	Exploring the Development, Validity, and Utility of the Short-Form Version of the CHoice of Outcome In Cbt for PsychosEs: A Patient-Reported Outcome Measure of Psychological Recovery. Schizophrenia Bulletin, 2021, 47, 653-661.	2.3	8
9	The Early Youth Engagement in first episode psychosis (EYE-2) study: pragmatic cluster randomised controlled trial of implementation, effectiveness and cost-effectiveness of a team-based motivational engagement intervention to improve engagement. Trials, 2021, 22, 272.	0.7	5
10	Optimising AVATAR therapy for people who hear distressing voices: study protocol for the AVATAR2 multi-centre randomised controlled trial. Trials, 2021, 22, 366.	0.7	21
11	Psychophysiological stress-reactivity in clinical and non-clinical voice-hearers. Schizophrenia Research, 2021, 235, 52-59.	1.1	1
12	Networks underpinning emotion: A systematic review and synthesis of functional and effective connectivity. Neurolmage, 2021, 243, 118486.	2.1	24
13	You read my mind: fMRI markers of threatening appraisals in people with persistent psychotic experiences. NPJ Schizophrenia, 2021, 7, 49.	2.0	2
14	Trauma and Social Pathways to Psychosis, and Where the Two Paths Meet. Frontiers in Psychiatry, 2021, 12, 804971.	1.3	6
15	Changes in care costs associated with cognitive behavioural therapy for psychosis delivered in routine clinical practice. Journal of Mental Health, 2020, 29, 605-610.	1.0	6
16	Clinical and theoretical relevance of responses to analogues of psychotic experiences in people with psychotic experiences with and without a need-for-care: an experimental study. Psychological Medicine, 2020, 50, 761-770.	2.7	5
17	Linguistic alignment and theory of mind impairments in schizophrenia patients' dialogic interactions. Psychological Medicine, 2020, 50, 2194-2202.	2.7	16
18	Insight-related beliefs and controllability appraisals contribute little to hallucinated voices: a transdiagnostic network analysis study. European Archives of Psychiatry and Clinical Neuroscience, 2020, 271, 1525-1535.	1.8	2

#	Article	IF	Citations
19	Mindfulness-based crisis interventions (MBCI) for psychosis within acute inpatient psychiatric settings; a feasibility randomised controlled trial. BMC Psychiatry, 2020, 20, 193.	1.1	18
20	The effects of voice content on stress reactivity: A simulation paradigm of auditory verbal hallucinations. Schizophrenia Research, 2019, , .	1.1	3
21	Improving Access to psychological therapies for people with severe mental illness (IAPT-SMI): Lessons from the South London and Maudsley psychosis demonstration site. Behaviour Research and Therapy, 2019, 116, 104-110.	1.6	15
22	Beyond Trauma: A Multiple Pathways Approach to Auditory Hallucinations in Clinical and Nonclinical Populations. Schizophrenia Bulletin, 2019, 45, S24-S31.	2.3	51
23	Overgeneral autobiographical memory bias in clinical and non-clinical voice hearers. Psychological Medicine, 2019, 49, 113-120.	2.7	6
24	A systematic review and discussion of symptom specific cognitive behavioural approaches to delusions and hallucinations. Schizophrenia Research, 2019, 203, 66-79.	1.1	73
25	OUP accepted manuscript. Schizophrenia Bulletin, 2018, 44, 126-136.	2.3	12
26	The COMMAND trial of cognitive therapy to prevent harmful compliance with command hallucinations: predictors of outcome and mediators of change. Psychological Medicine, 2018, 48, 1966-1974.	2.7	23
27	The U&I study: study protocol for a feasibility randomised controlled trial of a pre-cognitive behavioural therapy digital â€`informed choice' intervention to improve attitudes towards uptake and implementation of CBT for psychosis. Trials, 2018, 19, 644.	0.7	10
28	Using fMRI and machine learning to predict symptom improvement following cognitive behavioural therapy for psychosis. NeuroImage: Clinical, 2018, 20, 1053-1061.	1.4	12
29	The COMMAND trial of cognitive therapy for harmful compliance with command hallucinations (CTCH): a qualitative study of acceptability and tolerability in the UK. BMJ Open, 2018, 8, e021657.	0.8	8
30	A systematic scoping review of psychological therapies for psychosis within acute psychiatric in-patient settings. British Journal of Psychiatry, 2018, 213, 490-497.	1.7	31
31	Psychotic Experiences and Related Distress: A Cross-national Comparison and Network Analysis Based on 7141 Participants From 13 Countries. Schizophrenia Bulletin, 2018, 44, 1185-1194.	2.3	54
32	Moment-to-moment associations between negative affect, aberrant salience, and paranoia. Cognitive Neuropsychiatry, 2018, 23, 299-306.	0.7	30
33	The role of shame in people with a diagnosis of schizophrenia. British Journal of Clinical Psychology, 2017, 56, 115-129.	1.7	18
34	Experiences of outcome monitoring in service users with psychosis: Findings from an Improving Access to Psychological Therapies for people with Severe Mental Illness (<scp>IAPT</scp> â€ <scp>SMI</scp>) demonstration site. British Journal of Clinical Psychology, 2017, 56, 253-272.	1.7	26
35	Service user satisfaction with cognitive behavioural therapy for psychosis: Associations with therapy outcomes and perceptions of the therapist. British Journal of Clinical Psychology, 2017, 56, 84-102.	1.7	23
36	Clinical relevance of appraisals of persistent psychotic experiences in people with and without a need for care: an experimental study. Lancet Psychiatry, the, 2017, 4, 927-936.	3.7	42

#	Article	IF	Citations
37	Moment-to-moment associations between emotional disturbances, aberrant salience and persecutory delusions. European Psychiatry, 2017, 41, s838-s838.	0.1	3
38	Cognitive Bias Modification for paranoia (CBM-pa): study protocol for a randomised controlled trial. Trials, 2017, 18, 298.	0.7	15
39	Auditory verbal hallucinations and continuum models of psychosis: A systematic review of the healthy voice-hearer literature. Clinical Psychology Review, 2017, 51, 125-141.	6.0	161
40	Integrated Trauma-Focused Cognitive-Behavioural Therapy for Post-traumatic Stress and Psychotic Symptoms: A Case-Series Study Using Imaginal Reprocessing Strategies. Frontiers in Psychiatry, 2017, 8, 92.	1.3	29
41	Appraisals of psychotic experiences: an experimental investigation of symptomatic, remitted and non-need-for-care individuals. Psychological Medicine, 2016, 46, 1249-1263.	2.7	12
42	Cognitive and neural models of threat appraisal in psychosis: A theoretical integration. Psychiatry Research, 2016, 239, 131-138.	1.7	34
43	Clinical, socioâ€demographic and psychological characteristics in individuals with persistent psychotic experiences with and without a "need for careâ€. World Psychiatry, 2016, 15, 41-52.	4.8	119
44	Mindfulness-Based Crisis Interventions for patients with psychotic symptoms on acute psychiatric wards (amBITION study): protocol for a feasibility randomised controlled trial. Pilot and Feasibility Studies, 2016, 2, .	0.5	11
45	A brief CBT intervention for depersonalisation/derealisation in psychosis: study protocol for a feasibility randomised controlled trial. Pilot and Feasibility Studies, 2016, 2, 47.	0.5	11
46	Functional connectivity predictors and mechanisms of cognitive behavioural therapies: A systematic review with recommendations. Australian and New Zealand Journal of Psychiatry, 2016, 50, 311-321.	1.3	15
47	Mapping Depression in Schizophrenia: A Functional Magnetic Resonance Imaging Study. Schizophrenia Bulletin, 2016, 42, 802-813.	2.3	17
48	Cognitive Behavioral Therapy Normalizes Functional Connectivity for Social Threat in Psychosis. Schizophrenia Bulletin, 2016, 42, 684-692.	2.3	30
49	Effects of environmental noise on cognitive (dys)functions in schizophrenia: A pilot within-subjects experimental study. Schizophrenia Research, 2016, 173, 101-108.	1.1	16
50	Psychological therapies for psychosis: a view from the hills. Lancet Psychiatry, the, 2016, 3, 9-10.	3.7	3
51	The long-term effectiveness of cognitive behavior therapy for psychosis within a routine psychological therapies service. Frontiers in Psychology, 2015, 6, 1658.	1.1	26
52	Orbitofrontal cortex, emotional decision-making and response to cognitive behavioural therapy for psychosis. Psychiatry Research - Neuroimaging, 2015, 231, 298-307.	0.9	21
53	Which psychotic experiences are associated with a need for clinical care?. European Psychiatry, 2015, 30, 648-654.	0.1	17
54	Changes in delusional dimensions and emotions over eight weeks of antipsychotic treatment in acute patients. Psychiatry Research, 2015, 228, 393-398.	1.7	4

#	Article	IF	CITATIONS
55	Opportunities and challenges in Improving Access to Psychological Therapies for people with Severe Mental Illness (IAPT-SMI): Evaluating the first operational year of the South London and Maudsley (SLaM) demonstration site for psychosis. Behaviour Research and Therapy, 2015, 64, 24-30.	1.6	67
56	Psychological flexibility and nonjudgemental acceptance in voice hearers: relationships with omnipotence and distress. Australian and New Zealand Journal of Psychiatry, 2014, 48, 1150-1162.	1.3	37
57	Understanding noise stress-induced cognitive impairment in healthy adults and its implications for schizophrenia. Noise and Health, 2014, 16, 166.	0.4	48
58	Auditory Verbal Hallucinations in Persons With and Without a Need for Care. Schizophrenia Bulletin, 2014, 40, S255-S264.	2.3	236
59	Appraisals and Responses to Experimental Symptom Analogues in Clinical and Nonclinical Individuals With Psychotic Experiences. Schizophrenia Bulletin, 2014, 40, 845-855.	2.3	33
60	Predictors of distress associated with psychoticâ€like anomalous experiences in clinical and nonâ€clinical populations. British Journal of Clinical Psychology, 2014, 53, 213-227.	1.7	69
61	An oversimplification of psychosis, its treatment, and its outcomes?. British Journal of Psychiatry, 2014, 205, 159-160.	1.7	9
62	Cognitive Biases Questionnaire for Psychosis. Schizophrenia Bulletin, 2014, 40, 300-313.	2.3	66
63	Psychological Therapies for Auditory Hallucinations (Voices): Current Status and Key Directions for Future Research. Schizophrenia Bulletin, 2014, 40, S202-S212.	2.3	153
64	Higher-order semantic processing in formal thought disorder in schizophrenia. Psychiatry Research, 2014, 216, 168-176.	1.7	9
65	Jumping to perceptions and to conclusions: Specificity to hallucinations and delusions. Schizophrenia Research, 2014, 154, 68-72.	1.1	23
66	Cognitive behaviour therapy to prevent harmful compliance with command hallucinations (COMMAND): a randomised controlled trial. Lancet Psychiatry, the, 2014, 1, 23-33.	3.7	120
67	Verbatim recall in formal thought disorder in schizophrenia: a study of contextual influences. Cognitive Neuropsychiatry, 2014, 19, 337-358.	0.7	9
68	Changes in delusions in the early phase of antipsychotic treatment – An experience sampling study. Psychiatry Research, 2014, 215, 568-573.	1.7	42
69	Symptom Dimensions of the Psychotic Symptom Rating Scales in Psychosis: A Multisite Study. Schizophrenia Bulletin, 2014, 40, S265-S274.	2.3	92
70	Cognitive Behavioural Therapy for Psychosis. , 2014, , 139-164.		0
71	Detecting improvements in acute psychotic symptoms using experience sampling methodology. Psychiatry Research, 2013, 210, 82-88.	1.7	36
72	The role of safety-seeking behaviours in maintaining threat appraisals in psychosis. Behaviour Research and Therapy, 2013, 51, 75-81.	1.6	32

#	Article	IF	CITATIONS
7 3	Cognitive biases and auditory verbal hallucinations in healthy and clinical individuals. Psychological Medicine, 2013, 43, 2339-2347.	2.7	28
74	Intrusions in trauma and psychosis: information processing and phenomenology. Psychological Medicine, 2012, 42, 2313-2323.	2.7	14
7 5	Appraisals, psychotic symptoms and affect in daily life. Psychological Medicine, 2012, 42, 1013-1023.	2.7	41
76	It's not what you hear, it's the way you think about it: appraisals as determinants of affect and behaviour in voice hearers. Psychological Medicine, 2012, 42, 1507-1514.	2.7	75
77	A qualitative comparison of psychoticâ€like phenomena in clinical and nonâ€clinical populations. British Journal of Clinical Psychology, 2012, 51, 37-53.	1.7	59
78	The impact of negative schemas, mood and psychological flexibility on delusional ideation – mediating and moderating effects. Psychosis, 2012, 4, 6-18.	0.4	34
79	Sensorimotor gating and clinical outcome following cognitive behaviour therapy for psychosis. Schizophrenia Research, 2012, 134, 232-238.	1.1	29
80	Ethnicity, violent offending, and vulnerability to schizophrenia: A pilot study. Psychology and Psychotherapy: Theory, Research and Practice, 2012, 85, 143-149.	1.3	2
81	Predicting changes in delusional ideation: The role of mindfulness and negative schemas. Psychology and Psychotherapy: Theory, Research and Practice, 2012, 85, 243-259.	1.3	13
82	Neural changes following cognitive behaviour therapy for psychosis: a longitudinal study. Brain, 2011, 134, 2396-2407.	3.7	61
83	Coping styles predict responsiveness to cognitive behaviour therapy in psychosis. Psychiatry Research, 2011, 187, 354-362.	1.7	35
84	The structural brain basis of responsiveness to cognitive behaviour therapy for psychosis. International Clinical Psychopharmacology, 2011, 26, e114-e115.	0.9	0
85	Clinical presentation and early care relationships in †poorâ€me' and †badâ€me' paranoia. British Journ Clinical Psychology, 2011, 50, 211-216.	nal of 1.7	6
86	A multi-centre, randomised controlled trial of cognitive therapy to prevent harmful compliance with command hallucinations. BMC Psychiatry, 2011, 11, 155.	1.1	18
87	Psychotic-Like Experiences, Appraisals, and Trauma. Journal of Nervous and Mental Disease, 2010, 198, 813-819.	0.5	63
88	Do Antipsychotics Improve Reasoning Biases? A Review. Psychosomatic Medicine, 2010, 72, 681-693.	1.3	57
89	Delusions and belief flexibility in psychosis. Psychology and Psychotherapy: Theory, Research and Practice, 2010, 83, 45-57.	1.3	33
90	A randomised controlled trial of cognitive behaviour therapy for psychosis in a routine clinical service. Acta Psychiatrica Scandinavica, 2010, 122, 302-318.	2.2	71

#	Article	IF	Citations
91	CHoice of Outcome In Cbt for psychosEs (CHOICE): The Development of a New Service User–Led Outcome Measure of CBT for Psychosis. Schizophrenia Bulletin, 2010, 36, 126-135.	2.3	107
92	Cognitive insight predicts favorable outcome in cognitive behavioral therapy for psychosis. Psychosis, 2010, 2, 23-33.	0.4	57
93	Jumping to conclusions and perceptions in early psychosis: Relationship with delusional beliefs. Cognitive Neuropsychiatry, 2010, 15, 422-440.	0.7	23
94	Cognitive insight in psychosis: The relationship between self-certainty and self-reflection dimensions and neuropsychological measures. Psychiatry Research, 2010, 178, 284-289.	1.7	65
95	The role of metacognitive beliefs in determining the impact of anomalous experiences: a comparison of help-seeking and non-help-seeking groups of people experiencing psychotic-like anomalies. Psychological Medicine, 2009, 39, 939-950.	2.7	70
96	The Therapeutic Relationship in CBT for Psychosis: Client, Therapist and Therapy Factors. Behavioural and Cognitive Psychotherapy, 2009, 37, 527-540.	0.9	56
97	Schizotypy, delusional ideation and wellâ€being in an American new religious movement population. Clinical Psychology and Psychotherapy, 2009, 16, 479-484.	1.4	16
98	Dorsolateral Prefrontal Cortex Activity Predicts Responsiveness to Cognitive–Behavioral Therapy in Schizophrenia. Biological Psychiatry, 2009, 66, 594-602.	0.7	105
99	Structural magnetic resonance imaging predictors of responsiveness to cognitive behaviour therapy in psychosis. Schizophrenia Research, 2009, 115, 146-155.	1.1	31
100	Specificity of the jumpâ€ŧo onclusions bias in deluded patients. British Journal of Clinical Psychology, 2008, 47, 239-244.	1.7	35
101	Responding mindfully to unpleasant thoughts and images: Reliability and validity of the Southampton mindfulness questionnaire (SMQ). British Journal of Clinical Psychology, 2008, 47, 451-455.	1.7	329
102	Uncontrollable voices and their relationship to gating deficits in schizophrenia. Schizophrenia Research, 2008, 101, 185-194.	1.1	39
103	Neurological basis of poor insight in psychosis: A voxel-based MRI study. Schizophrenia Research, 2008, 103, 40-51.	1.1	85
104	Misattribution bias of threat-related facial expressions is related to a longer duration of illness and poor executive function in schizophrenia and schizoaffective disorder. European Psychiatry, 2008, 23, 14-19.	0.1	130
105	Service-User Satisfaction with CBT for Psychosis. Behavioural and Cognitive Psychotherapy, 2007, 35, 109.	0.9	30
106	Identity and Cause of Problems: The Perceptions of Patients with a Diagnosis of Schizophrenia. Behavioural and Cognitive Psychotherapy, 2007, 35, 237.	0.9	2
107	Insight in psychosis: influence of cognitive ability and self-esteem. British Journal of Psychiatry, 2007, 191, 234-237.	1.7	56
108	The psychological processes underlying symptoms of thought interference in psychosis. Behaviour Research and Therapy, 2007, 45, 2726-2741.	1.6	21

#	Article	IF	CITATIONS
109	Insight, distress and coping styles in schizophrenia. Schizophrenia Research, 2007, 94, 12-22.	1.1	89
110	Cognitive functioning in delusions: A longitudinal analysis. Behaviour Research and Therapy, 2006, 44, 481-514.	1.6	164
111	Need for closure and anxiety in delusions: A longitudinal investigation in early psychosis. Behaviour Research and Therapy, 2006, 44, 1385-1396.	1.6	40
112	Psychological processes underlying delusional thinking in late-onset psychosis: a preliminary investigation. International Journal of Geriatric Psychiatry, 2006, 21, 768-777.	1.3	11
113	Disease, deficit or denial? Models of poor insight in psychosis. Acta Psychiatrica Scandinavica, 2005, 112, 4-17.	2.2	121
114	Measuring Delusional Ideation: The 21-Item Peters et al. Delusions Inventory (PDI). Schizophrenia Bulletin, 2004, 30, 1005-1022.	2.3	491
115	Reasoning in Believers in the Paranormal. Journal of Nervous and Mental Disease, 2004, 192, 727-733.	0.5	37
116	Self-discrepancies, attentional bias and persecutory delusions. British Journal of Clinical Psychology, 2003, 42, 1-12.	1.7	52
117	Trainee clinical psychologists' adaptation and professional functioning: a longitudinal study. Clinical Psychology and Psychotherapy, 2003, 10, 41-54.	1.4	29
118	A quantitative genetic analysis of schizotypal personality traits. Psychological Medicine, 2003, 33, 803-816.	2.7	159
119	NEED FOR CLOSURE AND JUMPING-TO-CONCLUSIONS IN DELUSION-PRONE INDIVIDUALS. Journal of Nervous and Mental Disease, 2002, 190, 27-31.	0.5	169
120	Perceptual organization deficits in psychotic patients. Psychiatry Research, 2002, 110, 125-135.	1.7	18
121	Schizotypy and patterns of lateral asymmetry on hemisphere-specific language tasks. Psychiatry Research, 2001, 103, 179-192.	1.7	44
122	Neuroimaging correlates of negative priming. NeuroReport, 2001, 12, 3619-3624.	0.6	48
123	The Incidence of Schizotypy among Cannabis and Alcohol Users. Journal of Nervous and Mental Disease, 2001, 189, 741-748.	0.5	110
124	The relationship between cognitive inhibition and psychotic symptoms Journal of Abnormal Psychology, 2000, 109, 386-395.	2.0	100
125	A longitudinal study of the psychological adaptation of trainee clinical psychologists. Clinical Psychology and Psychotherapy, 2000, 7, 394-400.	1.4	12

Measurement of Delusional Ideation in the Normal Population: Introducing the PDI (Peters et al.) Tj ETQq0 0 0 rgBT $_{2.3}$ Qverlock $_{660}$ Tf 50 6

8

126

#	ARTICLE	IF	CITATION
127	Cognitive Asymmetry Patterns in Schizophrenia: Active and Withdrawn Syndromes and Sex Differences as Moderators. Schizophrenia Bulletin, 1999, 25, 349-362.	2.3	36
128	Delusional ideation in religious and psychotic populations. British Journal of Clinical Psychology, 1999, 38, 83-96.	1.7	207
129	The incidence of schizotypy in new religious movements. Personality and Individual Differences, 1999, 27, 55-67.	1.6	51
130	The psychological adaptation of psychologists in clinical training: the role of cognition, coping and social support., 1998, 5, 238-252.		13
131	Reasoning biases in delusion-prone individuals. British Journal of Clinical Psychology, 1998, 37, 285-302.	1.7	80
132	â€~Cognitive inhibition' and positive symptomatology in schizotypy. British Journal of Clinical Psychology, 1994, 33, 33-48.	1.7	84