

Graham Fent

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8781433/publications.pdf>

Version: 2024-02-01

13
papers

135
citations

1464605

7
h-index

1526636

10
g-index

13
all docs

13
docs citations

13
times ranked

196
citing authors

#	ARTICLE	IF	CITATIONS
1	Cardiovascular medication in patients with raised NT-proBNP, but no heart failure in the SHEAF registry. <i>Open Heart</i> , 2022, 9, e001974.	0.9	0
2	Predictors of subclinical systemic sclerosis primary heart involvement characterised by microvasculopathy and myocardial fibrosis. <i>Rheumatology</i> , 2021, 60, 2934-2945.	0.9	18
3	Using cardiovascular magnetic resonance to define mechanisms of comorbidity and to measure the effect of biological therapy: the CADERA observational study. <i>Efficacy and Mechanism Evaluation</i> , 2021, 8, 1-42.	0.9	0
4	Characterisation of the patients with suspected heart failure: experience from the SHEAF registry. <i>Open Heart</i> , 2021, 8, e001448.	0.9	10
5	Cardiovascular outcomes in systemic sclerosis with abnormal cardiovascular MRI and serum cardiac biomarkers. <i>RMD Open</i> , 2021, 7, e001689.	1.8	11
6	Abnormal electrophysiological testing associates with future incidental significant arrhythmia in scleroderma. <i>Rheumatology</i> , 2020, 59, 899-900.	0.9	4
7	Cardiovascular effects of biological versus conventional synthetic disease-modifying antirheumatic drug therapy in treatment-naïve, early rheumatoid arthritis. <i>Annals of the Rheumatic Diseases</i> , 2020, 79, 1414-1422.	0.5	32
8	Clinical predictors of all-cause mortality in patients presenting to specialist heart failure clinic with raised NT-proBNP and no heart failure. <i>ESC Heart Failure</i> , 2020, 7, 1791-1800.	1.4	7
9	Incidental significant arrhythmia in scleroderma associates with cardiac magnetic resonance measure of fibrosis and hs-TnI and NT-proBNP. <i>Rheumatology</i> , 2019, 58, 1221-1226.	0.9	31
10	First cardiovascular MRI study in individuals at risk of rheumatoid arthritis detects abnormal aortic stiffness suggesting an anti-citrullinated peptide antibody-mediated role for accelerated atherosclerosis. <i>Annals of the Rheumatic Diseases</i> , 2019, 78, 1138-1140.	0.5	15
11	Implantable loop recorder in systemic sclerosis over three years confirms incidental significant arrhythmia and suggests CMR and cardiac biomarker association. <i>Rheumatology</i> , 2018, 57, .	0.9	0
12	Carotid artery volumetric measures associate with clinical ten-year cardiovascular (CV) risk scores and individual traditional CV risk factors in rheumatoid arthritis; a carotid-MRI feasibility study. <i>Arthritis Research and Therapy</i> , 2018, 20, 266.	1.6	4
13	New Cardiac Magnetic Resonance Reference Ranges for Right Ventricular Volumes and Systolic Function. <i>Circulation: Cardiovascular Imaging</i> , 2016, 9, e004589.	1.3	3