

Anna Karjalainen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8783859/publications.pdf>

Version: 2024-02-01

10
papers

266
citations

1163117

8
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

497
citing authors

#	ARTICLE	IF	CITATIONS
1	Infliximab for 6 months added on combination therapy in early rheumatoid arthritis: 2-year results from an investigator-initiated, randomised, double-blind, placebo-controlled study (the NEO-RACo) Tj ETQq1 1 0.784314 rgBT76 Overlook	1.0	76
2	Rheumatoid Atlantoaxial Subluxation Can Be Prevented by Intensive Use of Traditional Disease Modifying Antirheumatic Drugs. <i>Journal of Rheumatology</i> , 2009, 36, 273-278.	2.0	50
3	Targeted treatment with a combination of traditional DMARDs produces excellent clinical and radiographic long-term outcomes in early rheumatoid arthritis regardless of initial infliximab. The 5-year follow-up results of a randomised clinical trial, the NEO-RACo trial. <i>Annals of the Rheumatic Diseases</i> , 2014, 73, 1954-1961.	0.9	40
4	Effectiveness and drug survival of TNF-inhibitors in the treatment of psoriatic arthritis: A prospective cohort study. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 46, 732-739.	3.4	37
5	1,25(OH) 2 D 3 and calcipotriol, its hypocalcemic analog, exert a long-lasting anti-inflammatory and anti-proliferative effect in synoviocytes cultured from patients with rheumatoid arthritis and osteoarthritis. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2017, 173, 13-22.	2.5	25
6	Cervical Spine Involvement among Patients with Rheumatoid Arthritis Treated Actively with Treat-to-target Strategy: 10-year Results of the NEO-RACo Study. <i>Journal of Rheumatology</i> , 2020, 47, 1160-1164.	2.0	12
7	Patient-reported outcomes as predictors of remission in early rheumatoid arthritis patients treated with tight control treat-to-target approach. <i>Rheumatology International</i> , 2017, 37, 825-830.	3.0	9
8	High burden of adverse events is associated with reduced remission rates in early rheumatoid arthritis. <i>Clinical Rheumatology</i> , 2018, 37, 1689-1694.	2.2	8
9	Failure in Longterm Treatment is Rare in Actively Treated Patients with Rheumatoid Arthritis, But May Be Predicted by High Health Assessment Score at Baseline and by Residual Disease Activity at 3 and 6 Months: The 5-year Followup Results of the Randomized Clinical NEO-RACo Trial. <i>Journal of Rheumatology</i> , 2014, 41, 2379-2385.	2.0	6
10	Cytokine data obtained from synovial stromal cells of patients with rheumatoid arthritis or osteoarthritis. <i>Data in Brief</i> , 2017, 12, 593-602.	1.0	3