

Stephane Arques

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8784522/stephane-arques-publications-by-citations.pdf>

Version: 2024-04-29

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

48
papers

1,564
citations

16
h-index

39
g-index

50
ext. papers

1,790
ext. citations

3.5
avg, IF

4.87
L-index

#	Paper	IF	Citations
48	Adjusted clopidogrel loading doses according to vasodilator-stimulated phosphoprotein phosphorylation index decrease rate of major adverse cardiovascular events in patients with clopidogrel resistance: a multicenter randomized prospective study. <i>Journal of the American College of Cardiology</i> , 2008 , 51, 1404-11	15.1	467
47	Tailored clopidogrel loading dose according to platelet reactivity monitoring to prevent acute and subacute stent thrombosis. <i>American Journal of Cardiology</i> , 2009 , 103, 5-10	3	237
46	Human serum albumin in the clinical syndrome of heart failure. <i>Journal of Cardiac Failure</i> , 2011 , 17, 451-8,3		133
45	Human serum albumin in cardiovascular diseases. <i>European Journal of Internal Medicine</i> , 2018 , 52, 8-12	3.9	110
44	Clopidogrel loading dose adjustment according to platelet reactivity monitoring in patients carrying the 2C19*2 loss of function polymorphism. <i>Journal of the American College of Cardiology</i> , 2010 , 56, 1630-6	15.1	98
43	Current clinical applications of spectral tissue Doppler echocardiography (E/EVratio) as a noninvasive surrogate for left ventricular diastolic pressures in the diagnosis of heart failure with preserved left ventricular systolic function. <i>Cardiovascular Ultrasound</i> , 2007 , 5, 16	2.4	74
42	Hypoalbuminemia in elderly patients with acute diastolic heart failure. <i>Journal of the American College of Cardiology</i> , 2003 , 42, 712-6	15.1	66
41	Usefulness of serum albumin concentration for in-hospital risk stratification in frail, elderly patients with acute heart failure. Insights from a prospective, monocenter study. <i>International Journal of Cardiology</i> , 2008 , 125, 265-7	3.2	48
40	Usefulness of serum albumin and serum total cholesterol in the prediction of hospital death in older patients with severe, acute heart failure. <i>Archives of Cardiovascular Diseases</i> , 2011 , 104, 502-8	2.7	34
39	Usefulness of bedside tissue Doppler echocardiography and B-type natriuretic peptide (BNP) in differentiating congestive heart failure from noncardiac cause of acute dyspnea in elderly patients with a normal left ventricular ejection fraction and permanent, nonvalvular atrial fibrillation: insights from a prospective, monocenter study. <i>Echocardiography</i> , 2007 , 24, 499-507	1.5	34
38	Intra-individual variability in clopidogrel responsiveness in coronary artery disease patients under long term therapy. <i>Platelets</i> , 2010 , 21, 503-7	3.6	27
37	Usefulness of N-terminal-pro-brain natriuretic peptide levels in predicting survival in heart transplant recipients. <i>American Journal of Cardiology</i> , 2004 , 94, 1585-7	3	27
36	Accuracy of tissue Doppler echocardiography in the emergency diagnosis of decompensated heart failure with preserved left ventricular systolic function: comparison with B-type natriuretic peptide measurement. <i>Echocardiography</i> , 2005 , 22, 657-64	1.5	21
35	Impact of P2Y12-ADP receptor polymorphism on the efficacy of clopidogrel dose-adjustment according to platelet reactivity monitoring in coronary artery disease patients. <i>Thrombosis Research</i> , 2010 , 125, e167-70	8.2	18
34	Accuracy of tissue Doppler echocardiography in the diagnosis of new-onset congestive heart failure in patients with levels of B-type natriuretic peptide in the midrange and normal left ventricular ejection fraction. <i>Echocardiography</i> , 2006 , 23, 627-34	1.5	17
33	Prevalence of angiographic coronary artery disease in patients hospitalized for acute diastolic heart failure without clinical and electrocardiographic evidence of myocardial ischemia on admission. <i>American Journal of Cardiology</i> , 2004 , 94, 133-5	3	16
32	B-type natriuretic peptide and tissue Doppler study findings in elderly patients hospitalized for acute diastolic heart failure. <i>American Journal of Cardiology</i> , 2005 , 96, 104-7	3	16

31	Comparative accuracy of color M-mode and tissue Doppler echocardiography in the emergency diagnosis of congestive heart failure in chronic hypertensive patients with normal left ventricular ejection fraction. <i>American Journal of Cardiology</i> , 2005 , 96, 1456-9	3	15
30	Potentials and limitations of color M-Mode and tissue Doppler indexes in identifying pseudonormal mitral filling pattern in patients with acute symptoms of heart failure and preserved left ventricular systolic function. <i>American Journal of Cardiology</i> , 2004 , 93, 1057-60	3	14
29	Prevalence and characteristics of coronary artery disease in heart failure with preserved and mid-range ejection fractions: A systematic angiography approach. <i>Archives of Cardiovascular Diseases</i> , 2018 , 111, 109-118	2.7	13
28	Predictors of B-type natriuretic peptide and left atrial volume index in patients with preserved left ventricular systolic function: an echocardiographic-catheterization study. <i>Archives of Cardiovascular Diseases</i> , 2010 , 103, 3-9	2.7	13
27	Usefulness of combined Doppler indexes in predicting left ventricular filling pressures in patients with acute heart failure. <i>American Journal of Cardiology</i> , 2003 , 92, 649-50	3	8
26	Tissue Doppler echocardiography for the diagnosis of new-onset heart failure with normal ejection fraction: influence of serum protein concentration on clinical interpretation in elderly patients. <i>Archives of Cardiovascular Diseases</i> , 2008 , 101, 343-50	2.7	6
25	Clinical Relevance of the Spectral Tissue Doppler E/eV Ratio in the Management of Patients with Atrial Fibrillation: a Comprehensive Review of the Literature. <i>Journal of Atrial Fibrillation</i> , 2018 , 11, 2038 ^{0.8}		6
24	Accuracy of the isovolumic relaxation time in the emergency diagnosis of new-onset congestive heart failure with preserved left ventricular systolic function in the setting of B-type natriuretic peptide levels in the mid-range. <i>International Journal of Cardiology</i> , 2008 , 124, 400-3	3.2	5
23	Comparison of B-type natriuretic peptide with left atrial enlargement by echocardiography for the diagnosis of new-onset congestive heart failure with a preserved left ventricular systolic function in the setting of longstanding hypertension. <i>International Journal of Cardiology</i> , 2008 , 128, 123-5	3.2	5
22	Accuracy of bedside tissue Doppler echocardiography for the prediction of in-hospital mortality in elderly patients with acute heart failure with preserved left ventricular systolic function. comparison with B-type natriuretic peptide measurement. <i>International Journal of Cardiology</i> , 2007 , 123, 69-72	3.2	4
21	Clinical relevance of spectral tissue Doppler-derived E/eMn the diagnosis of heart failure with preserved ejection fraction. <i>European Journal of Heart Failure</i> , 2018 , 20, 941	12.3	4
20	Usefulness of basal B-type natriuretic peptide levels for the diagnosis of diastolic heart failure in young patients: an echocardiographic-catheterization study. <i>International Journal of Cardiology</i> , 2010 , 145, 51-2	3.2	3
19	Rationale for a widespread use of the spectral tissue Doppler-derived E/EV despite its failure to closely predict invasively measured left ventricular diastolic pressures. <i>International Journal of Cardiology</i> , 2009 , 135, 139-40	3.2	3
18	Angiographic coronary artery disease associated with hypertensive heart failure and normal ejection fraction. Insights from a prospective monocenter study. <i>International Journal of Cardiology</i> , 2008 , 130, 75-7	3.2	3
17	Does Hypoalbuminemia Play a Role in the Worsening of Heart Failure?. <i>American Journal of Cardiology</i> , 2020 , 125, 1755	3	2
16	Correlation of NT-proBNP with diastolic left ventricular function in elderly patients with ischemic stroke. <i>International Journal of Cardiology</i> , 2010 , 140, 126-8	3.2	2
15	Usefulness of left atrial volume for the diagnosis of diastolic heart failure: an echocardiographic-catheterization study. <i>International Journal of Cardiology</i> , 2010 , 144, 317-9	3.2	2
14	Shortened mitral A wave deceleration time: an under-recognized but promising Doppler-derived index of left ventricular myocardial dysfunction in patients with normal left ventricular ejection fraction. <i>International Journal of Cardiology</i> , 2013 , 167, 3076-7	3.2	1

13	Hypoalbuminemia as a contributor to the progression of cardiovascular diseases. <i>Annals of Thoracic Surgery</i> , 2011 , 92, 2305; author reply 2305-6	2.7	1
12	Hypoalbuminemia as a risk factor of incident heart failure in the elderly. <i>American Heart Journal</i> , 2010 , 160, e39; author reply e41	4.9	1
11	Identification and quantification of prosthetic mitral regurgitation by flow convergence method using transthoracic approach. <i>Cardiovascular Ultrasound</i> , 2009 , 7, 7	2.4	1
10	Complementary role of bedside Doppler echocardiography to natriuretic peptides levels in the middle range for the diagnosis of acute heart failure. <i>European Journal of Heart Failure</i> , 2008 , 10, 1268	12.3	1
9	Pressions de remplissage en échocardiographie-doppler: Intérêt des nouveaux indices en routine. <i>Archives Des Maladies Du Coeur Et Des Vaisseaux - Pratique</i> , 2004 , 2004, 13-15	0	1
8	Pulmonary venous flow by Doppler echocardiography: usefulness of diastolic wave deceleration time in predicting filling pressures. <i>Journal of the American College of Cardiology</i> , 2004 , 43, 925-6; author reply 926	15.1	1
7	Serum albumin and cardiovascular disease: Does low serum albumin contribute to the emergence and worsening of some cardiovascular diseases?. <i>European Journal of Internal Medicine</i> , 2020 , 80, 122-123	3.9	1
6	Deterioration in Left Ventricular Systolic Function in Heart Failure With Preserved and Midrange Ejection Fraction: Don't Miss Coronary Artery Disease. <i>American Journal of Cardiology</i> , 2019 , 124, 1806	3	1
5	Myocardial oedema and congestive heart failure: one piece of the puzzle?. <i>European Journal of Heart Failure</i> , 2018 , 20, 827	12.3	
4	Accuracy of Doppler-derived pulmonary artery hypertension to predict heart failure with normal ejection fraction. <i>International Journal of Cardiology</i> , 2010 , 145, 370-372	3.2	
3	Predicting noninvasively left ventricular filling pressures. <i>American Journal of Cardiology</i> , 2008 , 102, 371-372		
2	Tissue Doppler echocardiography in predicting exercise intolerance. <i>American Journal of Cardiology</i> , 2006 , 98, 1298	3	
1	Clinical relevance of tissue doppler-derived e/e _v in patients with cardiovascular disease. <i>Echocardiography</i> , 2021 , 38, 375-376	1.5	