

Thais Steemburgo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8785297/publications.pdf>

Version: 2024-02-01

29
papers

987
citations

566801

15
h-index

552369

26
g-index

38
all docs

38
docs citations

38
times ranked

1573
citing authors

#	ARTICLE	IF	CITATIONS
1	Fiber intake and glycemic control in patients with type 2 diabetes mellitus: a systematic review with meta-analysis of randomized controlled trials. <i>Nutrition Reviews</i> , 2013, 71, 790-801.	2.6	162
2	Leptin and TNF-alpha promoter methylation levels measured by MSP could predict the response to a low-calorie diet. <i>Journal of Physiology and Biochemistry</i> , 2011, 67, 463-470.	1.3	149
3	Effect of Antihyperglycemic Agents Added to Metformin and a Sulfonylurea on Glycemic Control and Weight Gain in Type 2 Diabetes: A Network Meta-analysis. <i>Annals of Internal Medicine</i> , 2011, 154, 672.	2.0	125
4	Improvement of the metabolic syndrome profile by soluble fibre "guar gum" in patients with type 2 diabetes: a randomised clinical trial. <i>British Journal of Nutrition</i> , 2013, 110, 1601-1610.	1.2	95
5	Intake of soluble fibers has a protective role for the presence of metabolic syndrome in patients with type 2 diabetes. <i>European Journal of Clinical Nutrition</i> , 2009, 63, 127-133.	1.3	56
6	The role of Dietary Approaches to Stop Hypertension (DASH) diet food groups in blood pressure in type 2 diabetes. <i>British Journal of Nutrition</i> , 2012, 108, 155-162.	1.2	42
7	Sources of Protein and Polyunsaturated Fatty Acids of the Diet and Microalbuminuria in Type 2 Diabetes Mellitus. <i>Journal of the American College of Nutrition</i> , 2008, 27, 528-537.	1.1	35
8	High Dietary Glycemic Index and Low Fiber Content Are Associated with Metabolic Syndrome in Patients with Type 2 Diabetes. <i>Journal of the American College of Nutrition</i> , 2011, 30, 141-148.	1.1	35
9	The rs9939609 Polymorphism in the <i>FTO</i> Gene Is Associated with Fat and Fiber Intakes in Patients with Type 2 Diabetes. <i>Journal of Nutrigenetics and Nutrigenomics</i> , 2013, 6, 97-106.	1.8	35
10	Higher fiber intake is associated with lower blood pressure levels in patients with type 1 diabetes. <i>Archives of Endocrinology and Metabolism</i> , 2018, 62, 47-54.	0.3	35
11	Nutritional risk assessment in critically ill cancer patients: systematic review. <i>Revista Brasileira De Terapia Intensiva</i> , 2015, 27, 274-83.	0.1	34
12	NUTRIC Score: Isolated and Combined Use With the NRS ₂₀₀₂ to Predict Hospital Mortality in Critically Ill Patients. <i>Journal of Parenteral and Enteral Nutrition</i> , 2020, 44, 1250-1256.	1.3	23
13	Endothelial dysfunction and serum fatty acid composition in patients with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2008, 57, 1167-1172.	1.5	22
14	Influence of thickening agents on rheological properties and sensory attributes of dysphagic diet. <i>Journal of Texture Studies</i> , 2021, 52, 587-602.	1.1	16
15	The rs7204609 Polymorphism in the Fat Mass and Obesity-Associated Gene is Positively Associated With Central Obesity and Microalbuminuria in Patients With Type 2 Diabetes From Southern Brazil. , 2012, 22, 228-236.		14
16	Hand Grip Strength and nutritional status in hospitalized oncological patients. <i>Revista De Nutricao</i> , 2018, 31, 489-499.	0.4	11
17	Methods of nutritional assessment and functional capacity in the identification of unfavorable clinical outcomes in hospitalized patients with cancer: a systematic review. <i>Nutrition Reviews</i> , 2022, 80, 786-811.	2.6	10
18	Basal metabolic rate in Brazilian patients with type 2 diabetes: comparison between measured and estimated values. <i>Archives of Endocrinology and Metabolism</i> , 2019, 63, 53-61.	0.3	6

#	ARTICLE	IF	CITATIONS
19	Efficacy of single-dose cholecalciferol in the blood pressure of patients with type 2 diabetes, hypertension and hypovitaminoses D. <i>Scientific Reports</i> , 2020, 10, 19611.	1.6	5
20	Nutritional status and dietary factors in cystic fibrosis patients with delta F508 mutation. <i>Revista De Nutricao</i> , 2015, 28, 359-369.	0.4	4
21	Relationship between the <i>FTO</i> Genotype and Early Chronic Kidney Disease in Type 2 Diabetes: The Mediating Role of Central Obesity, Hypertension, and High Albuminuria. <i>Lifestyle Genomics</i> , 2021, 14, 73-80.	0.6	4
22	Predictive equations for evaluation for resting energy expenditure in Brazilian patients with type 2 diabetes: what can we use?. <i>BMC Nutrition</i> , 2020, 6, 56.	0.6	2
23	Under-reporting of the energy intake in patients with type 2 diabetes. <i>Journal of Human Nutrition and Dietetics</i> , 2021, 34, 73-80.	1.3	2
24	Metabolic Syndrome in Hypertensive Patients: Correlation Between Anthropometric Data and Laboratory Findings: Response to Bulhoes and Araujo. <i>Diabetes Care</i> , 2007, 30, e140-e140.	4.3	0
25	Higher fibre intake is associated with lower blood pressure levels in patients with type 1 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2015, 7, .	1.2	0
26	Validity of Predictive Equations for Metabolic Basal Rate in Brazilian Patients with Type 2 Diabetes (P12-039-19). <i>Current Developments in Nutrition</i> , 2019, 3, nzz035.P12-039-19.	0.1	0
27	Relationship Between Risk, Nutritional Status and Functional Capacity and Clinical Outcomes in Cancer Patients: A Systematic Review. <i>Current Developments in Nutrition</i> , 2021, 5, 277.	0.1	0
28	Validade de Equações Preditivas para Gasto Energético de Repouso em Pacientes Brasileiros com Diabetes Tipo 2. <i>International Journal of Nutrology</i> , 2018, 11, .	0.0	0
29	Two variants of the Nutritional Risk in the Critically Ill Score as predictors of mortality in Intensive Care Unit patients at a Brazilian University Hospital. <i>Revista De Nutricao</i> , 0, 33, .	0.4	0