

Thais Steemburgo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8785297/thais-steemburgo-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

26

papers

966

citations

16

h-index

31

g-index

38

ext. papers

1,157

ext. citations

3.8

avg, IF

4.8

L-index

#	Paper	IF	Citations
26	Fiber intake and glycemic control in patients with type 2 diabetes mellitus: a systematic review with meta-analysis of randomized controlled trials. <i>Nutrition Reviews</i> , 2013 , 71, 790-801	6.4	131
25	Leptin and TNF-alpha promoter methylation levels measured by MSP could predict the response to a low-calorie diet. <i>Journal of Physiology and Biochemistry</i> , 2011 , 67, 463-70	5	122
24	Effect of antihyperglycemic agents added to metformin and a sulfonylurea on glycemic control and weight gain in type 2 diabetes: a network meta-analysis. <i>Annals of Internal Medicine</i> , 2011 , 154, 672-9	8	104
23	Validity of Predictive Equations for Metabolic Basal Rate in Brazilian Patients with Type 2 Diabetes (P12-039-19). <i>Current Developments in Nutrition</i> , 2019 , 3,	0.4	78
22	Higher fibre intake is associated with lower blood pressure levels in patients with type 1 diabetes. <i>Diabetology and Metabolic Syndrome</i> , 2015 , 7,	5.6	78
21	Relationship Between Risk, Nutritional Status and Functional Capacity and Clinical Outcomes in Cancer Patients: A Systematic Review. <i>Current Developments in Nutrition</i> , 2021 , 5, 277-277	0.4	78
20	Improvement of the metabolic syndrome profile by soluble fibre - guar gum - in patients with type 2 diabetes: a randomised clinical trial. <i>British Journal of Nutrition</i> , 2013 , 110, 1601-10	3.6	72
19	Intake of soluble fibers has a protective role for the presence of metabolic syndrome in patients with type 2 diabetes. <i>European Journal of Clinical Nutrition</i> , 2009 , 63, 127-33	5.2	40
18	The role of Dietary Approaches to Stop Hypertension (DASH) diet food groups in blood pressure in type 2 diabetes. <i>British Journal of Nutrition</i> , 2012 , 108, 155-62	3.6	33
17	Sources of protein and polyunsaturated fatty acids of the diet and microalbuminuria in type 2 diabetes mellitus. <i>Journal of the American College of Nutrition</i> , 2008 , 27, 528-37	3.5	31
16	High dietary glycemic index and low fiber content are associated with metabolic syndrome in patients with type 2 diabetes. <i>Journal of the American College of Nutrition</i> , 2011 , 30, 141-8	3.5	30
15	Nutritional risk assessment in critically ill cancer patients: systematic review. <i>Revista Brasileira De Terapia Intensiva</i> , 2015 , 27, 274-83	1.2	26
14	The rs9939609 polymorphism in the FTO gene is associated with fat and fiber intakes in patients with type 2 diabetes. <i>Journal of Nutrigenetics and Nutrigenomics</i> , 2013 , 6, 97-106		26
13	Higher fiber intake is associated with lower blood pressure levels in patients with type 1 diabetes. <i>Archives of Endocrinology and Metabolism</i> , 2018 , 62, 47-54	2.2	19
12	Endothelial dysfunction and serum fatty acid composition in patients with type 2 diabetes mellitus. <i>Metabolism: Clinical and Experimental</i> , 2008 , 57, 1167-72	12.7	17
11	The rs7204609 polymorphism in the fat mass and obesity-associated gene is positively associated with central obesity and microalbuminuria in patients with type 2 diabetes from Southern Brazil. <i>Journal of Renal Nutrition</i> , 2012 , 22, 228-236	3	11
10	NUTRIC Score: Isolated and Combined Use With the NRS-2002 to Predict Hospital Mortality in Critically Ill Patients. <i>Journal of Parenteral and Enteral Nutrition</i> , 2020 , 44, 1250-1256	4.2	10

9	Hand Grip Strength and nutritional status in hospitalized oncological patients. <i>Revista De Nutricao</i> , 2018 , 31, 489-499	1.8	4
8	Nutritional status and dietary factors in cystic fibrosis patients with delta F508 mutation. <i>Revista De Nutricao</i> , 2015 , 28, 359-369	1.8	3
7	Efficacy of single-dose cholecalciferol in the blood pressure of patients with type 2 diabetes, hypertension and hypovitaminoses D. <i>Scientific Reports</i> , 2020 , 10, 19611	4.9	3
6	Influence of thickening agents on rheological properties and sensory attributes of dysphagic diet. <i>Journal of Texture Studies</i> , 2021 ,	3.6	3
5	Basal metabolic rate in Brazilian patients with type 2 diabetes: comparison between measured and estimated values. <i>Archives of Endocrinology and Metabolism</i> , 2019 , 63, 53-61	2.2	1
4	Predictive equations for evaluation for resting energy expenditure in Brazilian patients with type 2 diabetes: what can we use?. <i>BMC Nutrition</i> , 2020 , 6, 56	2.5	1
3	Under-reporting of the energy intake in patients with type 2 diabetes. <i>Journal of Human Nutrition and Dietetics</i> , 2021 , 34, 73-80	3.1	1
2	Relationship between the FTO Genotype and Early Chronic Kidney Disease in Type 2 Diabetes: The Mediating Role of Central Obesity, Hypertension, and High Albuminuria. <i>Lifestyle Genomics</i> , 2021 , 14, 73-80	2	0
1	Metabolic syndrome in hypertensive patients: correlation between anthropometric data and laboratory findings: response to Bulhões and Araújo. <i>Diabetes Care</i> , 2007 , 30, e140	14.6	