
## Lawrence M Leeman

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/880022/publications.pdf Version: 2024-02-01



| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Hypertensive disorders of pregnancy. American Family Physician, 2008, 78, 93-100.   | 0.1 | 95        |
| 2  | Validity of Self-Reported Drug Use Information Among Pregnant Women. Maternal and Child Health<br>Journal, 2016, 20, 41-47.   | 1.5 | 85        |
| 3  | The Validity of Phosphatidylethanol in Dried Blood Spots of Newborns for the Identification of<br>Prenatal Alcohol Exposure. Alcoholism: Clinical and Experimental Research, 2014, 38, 1078-1085.                           | 2.4 | 67        |
| 4  | Patient-centered Care to Address Barriers for Pregnant Women with Opioid Dependence. Obstetrics and Gynecology Clinics of North America, 2017, 44, 95-107.  | 1.9 | 66        |
| 5  | The Effect of Perineal Lacerations on Pelvic Floor Function and Anatomy at 6 Months Postpartum in a<br>Prospective Cohort of Nulliparous Women. Birth, 2016, 43, 293-302.   | 2.2 | 58        |
| 6  | Neonatal Opioid Withdrawal Syndrome. Obstetrics and Gynecology Clinics of North America, 2014, 41, 317-334.   | 1.9 | 57        |
| 7  | Postpartum Perineal Pain in a Low Episiotomy Setting: Association with Severity of Genital Trauma,<br>Labor Care, and Birth Variables. Birth, 2009, 36, 283-288.  | 2.2 | 52        |
| 8  | Hypertensive Disorders of Pregnancy. American Family Physician, 2016, 93, 121-7.  | 0.1 | 50        |
| 9  | Association between prenatal opioid exposure, neonatal opioid withdrawal syndrome, and<br>neurodevelopmental and behavioral outcomes at 5–8 months of age. Early Human Development, 2019,<br>128, 69-76.                    | 1.8 | 37        |
| 10 | Do Unsutured Second-Degree Perineal Lacerations Affect Postpartum Functional Outcomes?. Journal of the American Board of Family Medicine, 2007, 20, 451-457.  | 1.5 | 35        |
| 11 | Perspectives on risk: Assessment of risk profiles and outcomes among women planning community birth in the United States. Birth, 2017, 44, 209-221.   | 2.2 | 34        |
| 12 | Misoprostol for intrauterine device insertion in nulliparous women: a randomized controlled trial.<br>American Journal of Obstetrics and Gynecology, 2014, 210, 208.e1-208.e5.  | 1.3 | 33        |
| 13 | Planned Home <scp>VBAC</scp> in the United States, 2004–2009: Outcomes, Maternity Care Practices, and Implications for Shared Decision Making. Birth, 2015, 42, 299-308.  | 2.2 | 33        |
| 14 | Immediate postpartum levonorgestrel intrauterine device insertion and breast-feeding outcomes: a<br>noninferiority randomized controlled trial. American Journal of Obstetrics and Gynecology, 2017, 217,<br>665.e1-665.e8. | 1.3 | 33        |
| 15 | Medical Barriers to Effective Contraception. Obstetrics and Gynecology Clinics of North America, 2007, 34, 19-29.   | 1.9 | 32        |
| 16 | Pelvic floor symptoms and quality of life changes during first pregnancy: a prospective cohort study.<br>International Urogynecology Journal, 2017, 28, 1701-1707.  | 1.4 | 32        |
| 17 | Patient-Choice Vaginal Delivery?. Annals of Family Medicine, 2006, 4, 265-268.  | 1.9 | 30        |
| 18 | Diminishing Availability of Trial of Labor After Cesarean Delivery in New Mexico Hospitals. Obstetrics<br>and Gynecology, 2013, 122, 242-247.   | 2.4 | 30        |

| #  | Article  | lF  | CITATIONS |
|----|--|-----|-----------|
| 19 | An ecological momentary assessment study examining posttraumatic stress disorder symptoms,<br>prenatal bonding, and substance use among pregnant women. Drug and Alcohol Dependence, 2019, 195,<br>33-39.                          | 3.2 | 30        |
| 20 | Effect of ibuprofen vs acetaminophen on postpartum hypertension in preeclampsia with severe<br>features: aÂdouble-masked, randomized controlled trial. American Journal of Obstetrics and<br>Gynecology, 2018, 218, 616.e1-616.e8. | 1.3 | 29        |
| 21 | The impact of Advanced Life Support in Obstetrics (ALSO) training in low-resource countries.<br>International Journal of Gynecology and Obstetrics, 2015, 131, 209-215.  | 2.3 | 28        |
| 22 | The nature and management of labor pain: part I. Nonpharmacologic pain relief. American Family<br>Physician, 2003, 68, 1109-12.  | 0.1 | 26        |
| 23 | The nature and management of labor pain: part II. Pharmacologic pain relief. American Family Physician, 2003, 68, 1115-20.   | 0.1 | 25        |
| 24 | Advanced Gestational Age Increases Serum Carbohydrate-Deficient Transferrin Levels in Abstinent<br>Pregnant Women. Alcohol and Alcoholism, 2012, 47, 683-687.  | 1.6 | 24        |
| 25 | Prevalence of alcohol use in pregnant women with substance use disorder. Drug and Alcohol<br>Dependence, 2018, 187, 305-310.   | 3.2 | 23        |
| 26 | The effect of prenatal substance use and maternal contingent responsiveness on infant affect. Early<br>Human Development, 2017, 115, 51-59.  | 1.8 | 21        |
| 27 | A Native American Community with a 7% Cesarean Delivery Rate: Does Case Mix, Ethnicity, or Labor<br>Management Explain the Low Rate?. Annals of Family Medicine, 2003, 1, 36-43.   | 1.9 | 19        |
| 28 | Perineal body stretch during labor does not predict perineal laceration, postpartum incontinence, or postpartum sexual function: a cohort study. International Urogynecology Journal, 2016, 27, 1193-1200.                         | 1.4 | 19        |
| 29 | "You can't do that 'round here†a case study of the introduction of medical abortion care at a<br>University Medical Center. Contraception, 2005, 71, 84-88.  | 1.5 | 18        |
| 30 | Ethylglucuronide in maternal hair as a biomarker of prenatal alcohol exposure. Alcohol, 2015, 49,<br>617-623.  | 1.7 | 18        |
| 31 | Hepatitis C Cascade of Care Among Pregnant Women on Opioid Agonist Pharmacotherapy Attending a<br>Comprehensive Prenatal Program. Maternal and Child Health Journal, 2017, 21, 1778-1783.  | 1.5 | 18        |
| 32 | Characteristics and outcomes of women using emergency medical services for third-trimester pregnancy-related problems in India: a prospective observational study. BMJ Open, 2016, 6, e011459.                                     | 1.9 | 16        |
| 33 | Group Prenatal Care for Women Receiving Medicationâ€Assisted Treatment for Opioid Use Disorder in<br>Pregnancy: An Interprofessional Approach. Journal of Midwifery and Women's Health, 2019, 64, 217-224.                         | 1.3 | 16        |
| 34 | Community Versus Outâ€ofâ€Hospital Birth: What's in a Name?. Journal of Midwifery and Women's Health,<br>2019, 64, 9-11.   | 1.3 | 15        |
| 35 | Breastfeeding Intention Compared With Breastfeeding Postpartum Among Women Receiving<br>Medication-Assisted Treatment. Journal of Human Lactation, 2019, 35, 71-79.  | 1.6 | 15        |
| 36 | Prenatal maternal posttraumatic stress disorder as a risk factor for adverse birth weight and<br>gestational age outcomes: A systematic review and meta-analysis. Journal of Affective Disorders, 2021,<br>295, 530-540.           | 4.1 | 15        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | A mother's bond: An ecological momentary assessment study of posttraumatic stress disorder<br>symptoms and substance craving during pregnancy Psychology of Addictive Behaviors, 2020, 34,<br>269-280.    | 2.1 | 15        |
| 38 | A randomized controlled trial of nitrous oxide for intrauterine device insertion in nulliparous women. International Journal of Gynecology and Obstetrics, 2016, 135, 145-148.                            | 2.3 | 14        |
| 39 | Hypersynchrony in MEG spectral amplitude in prospectively-identified 6-month-old infants prenatally exposed to alcohol. NeuroImage: Clinical, 2018, 17, 826-834.  | 2.7 | 14        |
| 40 | Repair of obstetric perineal lacerations. American Family Physician, 2003, 68, 1585-90.   | 0.1 | 14        |
| 41 | Association between intrapartum fetal heart rate patterns and neonatal abstinence syndrome in<br>methadone exposed neonates. Journal of Maternal-Fetal and Neonatal Medicine, 2011, 24, 955-959.          | 1.5 | 13        |
| 42 | Emergency in the clinic: a simulation curriculum to improve outpatient safety. American Journal of Obstetrics and Gynecology, 2017, 217, 699.e1-699.e13.  | 1.3 | 12        |
| 43 | Post-discharge healthcare utilization in infants with neonatal opioid withdrawal syndrome.<br>Neurotoxicology and Teratology, 2021, 86, 106975.   | 2.4 | 12        |
| 44 | Can mifepristone medication abortion be successfully integrated into medical practices that do not offer surgical abortion?. Contraception, 2007, 76, 96-100.   | 1.5 | 11        |
| 45 | Finding the breech: Influence of breech presentation on mode of delivery based on timing of diagnosis, attempt at external cephalic version, and provider success with version. Birth, 2017, 44, 222-229. | 2.2 | 10        |
| 46 | Nitrous oxide versus oral sedation for pain management of first-trimester surgical abortion — a randomized study. Contraception, 2017, 96, 118-123.   | 1.5 | 10        |
| 47 | State of the breech in 2020: Guidelines support maternal choice, but skills are lost… Birth, 2020, 47,<br>165-168.  | 2.2 | 8         |
| 48 | Prevalence of marijuana use in pregnant women with concurrent opioid use disorder or alcohol use<br>in pregnancy. Addiction Science & Clinical Practice, 2022, 17, 3.                                     | 2.6 | 8         |
| 49 | Isolated oligohydramnios at term: is induction indicated?. Journal of Family Practice, 2005, 54, 25-32.   | 0.2 | 8         |
| 50 | Postpartum Weight Retention in Primiparous Women and Weight Outcomes in Their Offspring.<br>Journal of Midwifery and Women's Health, 2019, 64, 427-434.   | 1.3 | 7         |
| 51 | You Didn't Drink During Pregnancy, Did You?. Alcoholism: Clinical and Experimental Research, 2021, 45, 543-547.   | 2.4 | 7         |
| 52 | Clinical Practice Guideline Executive Summary: Labor After Cesarean/Planned Vaginal Birth After<br>Cesarean. Annals of Family Medicine, 2015, 13, 80-81.  | 1.9 | 6         |
| 53 | Disparities in breastfeeding outcomes among women with opioid use disorder. Acta Paediatrica,<br>International Journal of Paediatrics, 2020, 109, 1064-1066.  | 1.5 | 6         |
| 54 | Prenatal opioid-exposed infant extracellular miRNA signature obtained at birth predicts severity of neonatal opioid withdrawal syndrome. Scientific Reports, 2022, 12, 5941.                              | 3.3 | 6         |

LAWRENCE M LEEMAN

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Prenatal Counseling Regarding Cesarean Delivery. Obstetrics and Gynecology Clinics of North America, 2008, 35, 473-495.   | 1.9 | 5         |
| 56 | The effect of prenatal alcohol co-exposure on neonatal abstinence syndrome in infants born to<br>mothers in opioid maintenance treatment. Journal of Maternal-Fetal and Neonatal Medicine, 2016, 29,<br>783-788.                            | 1.5 | 5         |
| 57 | Best Practices for Engaging Pregnant and Postpartum Women at Risk of Substance Use in Longitudinal<br>Research Studies: a Qualitative Examination of Participant Preferences. Adversity and Resilience<br>Science, 2020, 1, 235-246.        | 2.6 | 5         |
| 58 | Maternal verbal scaffolding: association with higher language skills for 20-month-old children with prenatal polysubstance exposure. Early Human Development, 2021, 160, 105423.  | 1.8 | 1         |
| 59 | Cesarean delivery. clinical procedures and techniques: video, monograph and interactive CME. Journal of Midwifery and Women's Health, 2004, 49, e4-e4.  | 1.3 | 0         |
| 60 | Induction of labour at 37–38â€weeks in women with large fetuses decreases the likelihood of shoulder<br>dystocia; however, overall benefit of early-term delivery has not been demonstrated. Evidence-Based<br>Medicine, 2015, 20, 198-198. | 0.6 | 0         |
| 61 | Patient-choice cesarean delivery. American Family Physician, 2005, 72, 697, 700, 705.   | 0.1 | 0         |
| 62 | Promoting Safety in Community-Based Birth Settings. American Family Physician, 2021, 103, 650-652.  | 0.1 | 0         |