

# Simone N Santos

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8826667/publications.pdf>

Version: 2024-02-01

21  
papers

627  
citations

759055

12  
h-index

677027

22  
g-index

22  
all docs

22  
docs citations

22  
times ranked

1223  
citing authors

#	ARTICLE	IF	CITATIONS
1	Early Morning Attenuation of Endothelial Function in Healthy Humans. <i>Circulation</i> , 2004, 109, 2507-2510.	1.6	234
2	Luminal Esophageal Temperature Monitoring With a Deflectable Esophageal Temperature Probe and Intracardiac Echocardiography May Reduce Esophageal Injury During Atrial Fibrillation Ablation Procedures. <i>Circulation: Arrhythmia and Electrophysiology</i> , 2011, 4, 149-156.	2.1	63
3	Sarcopenia, but not excess weight or increased caloric intake, is associated with coronary subclinical atherosclerosis in the very elderly. <i>Atherosclerosis</i> , 2017, 258, 138-144.	0.4	48
4	Association of systemic inflammatory activity with coronary and carotid atherosclerosis in the very elderly. <i>Atherosclerosis</i> , 2011, 216, 212-216.	0.4	38
5	Timing and Dose of Statin Therapy Define Its Impact on Inflammatory and Endothelial Responses During Myocardial Infarction. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 1240-1246.	1.1	36
6	Relation of markers of inflammation (C-reactive protein, white blood cell count, and Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 547 Td (lipop 171-176.	0.8	28
7	Low zinc levels is associated with increased inflammatory activity but not with atherosclerosis, arteriosclerosis or endothelial dysfunction among the very elderly. <i>BBA Clinical</i> , 2014, 2, 1-6.	4.1	28
8	HDL levels and oxidizability during myocardial infarction are associated with reduced endothelial-mediated vasodilation and nitric oxide bioavailability. <i>Atherosclerosis</i> , 2014, 237, 840-846.	0.4	25
9	Elevated CETP activity during acute phase of myocardial infarction is independently associated with endothelial dysfunction and adverse clinical outcome. <i>Atherosclerosis</i> , 2014, 237, 777-783.	0.4	22
10	Association of Cardiovascular Risk Factors with Microvascular and Conduit Artery Function in Hypertensive Subjects. <i>American Journal of Hypertension</i> , 2007, 20, 735-742.	1.0	20
11	Most of the patients presenting myocardial infarction would not be eligible for intensive lipid-lowering based on clinical algorithms or plasma C-reactive protein. <i>Atherosclerosis</i> , 2011, 214, 148-150.	0.4	16
12	Enhanced parathyroid hormone levels are associated with left ventricle hypertrophy in very elderly men and women. <i>Journal of the American Society of Hypertension</i> , 2015, 9, 697-704.	2.3	16
13	Low HDL cholesterol but not high LDL cholesterol is independently associated with subclinical coronary atherosclerosis in healthy octogenarians. <i>Aging Clinical and Experimental Research</i> , 2015, 27, 61-67.	1.4	14
14	Omega-3 intake is associated with attenuated inflammatory response and cardiac remodeling after myocardial infarction. <i>Nutrition Journal</i> , 2019, 18, 29.	1.5	10
15	Atheroprotective Properties of Serum IGF-1 in the Carotid and Coronary Territories and Beneficial Role on the Physical Fitness of the Oldest Old. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2016, 71, 1281-1288.	1.7	8
16	Catecholaminergic polymorphic ventricular tachycardia: a current overview. <i>Future Cardiology</i> , 2009, 5, 191-199.	0.5	4
17	Osteopontin in bone mineral density of very old Brazilians. <i>Journal of Bone and Mineral Metabolism</i> , 2013, 31, 449-454.	1.3	4
18	Apolipoprotein E genotype is associated with apolipoprotein B plasma levels but not with coronary calcium score in very elderly individuals in primary care setting. <i>Gene</i> , 2014, 539, 275-278.	1.0	3

#	ARTICLE	IF	CITATIONS
19	TCF7L2 polymorphism is associated with low nitric oxide release, endothelial dysfunction and enhanced inflammatory response after myocardial infarction. <i>BBA Clinical</i> , 2016, 5, 159-165.	4.1	2
20	<i>To the Editor:</i> . <i>Journal of Cardiovascular Electrophysiology</i> , 2009, 20, E68; author reply E69.	0.8	1
21	Adverse outcome has a U-shaped relation with acute phase change in insulin sensitivity after ST-Elevation Myocardial Infarction. <i>International Journal of Cardiology</i> , 2018, 254, 16-22.	0.8	1