

Jacek Dabert

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8832189/publications.pdf>

Version: 2024-02-01

52
papers

1,075
citations

566801

15
h-index

433756

31
g-index

53
all docs

53
docs citations

53
times ranked

782
citing authors

#	ARTICLE	IF	CITATIONS
1	Molecular phylogeny of acariform mites (Acari, Arachnida): Strong conflict between phylogenetic signal and long-branch attraction artifacts. <i>Molecular Phylogenetics and Evolution</i> , 2010, 56, 222-241.	1.2	272
2	A new feather mite species of the genus <i>Proctophyllodes</i> Robin, 1877 (Astigmata: Proctophyllodidae) from the Long-tailed Tit <i>Aegithalos caudatus</i> (Passeriformes: Aegithalidae) – morphological description with DNA barcode data. <i>Zootaxa</i> , 2012, 3253, 54.	0.2	138
3	Origin and Evolution of Feather Mites (Astigmata). <i>Experimental and Applied Acarology</i> , 1999, 23, 437-454.	0.7	104
4	Molecular evidence for species separation in the water mite <i>Hygrobates nigromaculatus</i> Lebert, 1879 (Acari, Hydrachnidia): evolutionary consequences of the loss of larval parasitism. <i>Aquatic Sciences</i> , 2010, 72, 347-360.	0.6	68
5	Higher-level molecular phylogeny of the water mites (Acariformes: Prostigmata: Parasitengonina: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50 147 T	1.2	54
6	Phylogeny of Feather Mite Subfamily Avenzoariinae (Acari: Analgoidea: Avenzoariidae) Inferred from Combined Analyses of Molecular and Morphological Data. <i>Molecular Phylogenetics and Evolution</i> , 2001, 20, 124-135.	1.2	36
7	Differences in speciation progress in feather mites (Analgoidea) inhabiting the same host: the case of <i>Zachvatkinia</i> and <i>Alloptes</i> living on arctic and long-tailed skuas. <i>Experimental and Applied Acarology</i> , 2015, 65, 163-179.	0.7	27
8	Phylogenetic position of the enigmatic clawless eutardigrade genus <i>Apodibius</i> Dastych, 1983 (Tardigrada), based on 18S and 28S rRNA sequence data from its type species <i>A. confusus</i> . <i>Molecular Phylogenetics and Evolution</i> , 2014, 70, 70-75.	1.2	25
9	<i>Bifidochaetus</i> , a new Arctic genus of freshwater Chaetonotida (Gastrotricha) from Spitsbergen revealed by an integrative taxonomic approach. <i>Invertebrate Systematics</i> , 2016, 30, 398.	0.5	22
10	Morphological adaptation for precopulatory guarding in astigmatic mites (Acari: Acaridida). <i>International Journal of Acarology</i> , 1992, 18, 49-54.	0.3	20
11	Six new feather mite species (Acari: Astigmata) from the carolina parakeet <i>Conuropsis carolinensis</i> (Psittaciformes: Psittacidae), an extinct parrot of North America. <i>Journal of Natural History</i> , 2005, 39, 2257-2278.	0.2	20
12	Multidisciplinary analysis of <i>Knemidocoptes jamaicensis</i> parasitising the Common Chaffinch, <i>Fringilla coelebs</i> : proofs for a multispecies complex?. <i>Parasitology Research</i> , 2013, 112, 2373-2380.	0.6	20
13	Title is missing!. , 1999, 23, 525-549.		17
14	<i>Picobia dziabaszezewskii</i> sp. nov. (Acari, Syringophilidae) – combined description (morphology with DNA) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 T	0.2	17
15	DNA-barcoding contradicts morphology in quill mite species <i>Torotrogla merulae</i> and <i>T. rubeculi</i> (Prostigmata: Syringophilidae). <i>Folia Parasitologica</i> , 2013, 60, 51-60.	0.7	17
16	DNA barcodes reveal female dimorphism in syringophilid mites (Actinotrichida: Prostigmata: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 147 T	0.7	17
17	RAPD technique for taxonomic studies of <i>Pellia epiphylla</i> -complex (Hepaticae, Metzgeriales). <i>Genetica</i> , 1998, 104, 179-187.	0.5	16
18	The first report of <i>Knemidocoptes intermedius</i> Fain et Macfarlane, 1967 (Acari: Astigmata) in naturally infected European birds. <i>Parasitology Research</i> , 2011, 109, 237-240.	0.6	14

#	ARTICLE	IF	CITATIONS
19	A new quill mite <i>Syringophiloidus pseudonigritae</i> sp. nov. (Prostigmata, Syringophilidae) parasitizing <i>Pseudonigrita arnaudi</i> (Passeriformes, Ploceidae) – a combined description using morphology and DNA barcode data. <i>Zootaxa</i> , 2012, 3532, 64.	0.2	14
20	New feather mites of the family Pterolichidae (Acari: Pterolichoidea) from parrots (Aves: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50,702 Td (P	1.1	12
21	A New Species of the Genus <i>Proctophyllodes</i> (Analgoidea: Proctophyllodidae) from Cetti's Warbler <i>Cettia cetti</i> (Passeriformes: Sylviidae) with DNA Barcode Data. <i>Annales Zoologici</i> , 2008, 58, 397-402.	0.1	12
22	The explosive radiation, intense host-shifts and long-term failure to speciate in the evolutionary history of the feather mite genus <i>Analges</i> (Acariformes: Analgidae) from European passerines. <i>Zoological Journal of the Linnean Society</i> , 2022, 195, 673-694.	1.0	8
23	A new species of the feather mite genus <i>Titanolichus</i> Gaud & Atyeo, 1996 (Acari: Astigmata: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50,702 Td (P	1.1	7
24	Sweet or salty? The origin of freshwater gastrotrichs (Gastrotricha, Chaetonotida) revealed by molecular phylogenetic analysis. <i>Cladistics</i> , 2020, 36, 458-480.	1.5	7
25	<i>Picobia polonica</i> sp. n. (Acari: Prostigmata: Syringophilidae), a new species of quill mite from the domestic hen, <i>Gallus gallus domesticus</i> (Aves: Phasianidae). <i>Folia Parasitologica</i> , 2001, 48, 154-158.	0.7	7
26	New feather mite taxa of the <i>Rhytidelasma</i> Gaud, 1966 generic group (Astigmata: Pterolichidae) from the red-flanked lorikeet <i>Charmosyna placentis placentis</i> (Psittacidae). <i>Systematic Parasitology</i> , 2004, 58, 91-104.	0.5	6
27	Three new feather mite genera of the <i>Protolichus</i> generic group (Astigmata, Pterolichidae) from parrots (Aves, Psittaciformes) of the Old World. <i>Acta Parasitologica</i> , 2007, 52, 386.	0.4	6
28	Systematic revision of the feather mite genera <i>Apexolichus</i> Gaud et Atyeo and <i>Titanolichus</i> Gaud et Atyeo (Astigmata, Pterolichidae), parasites of parrots of the Old World (Psittaciformes, Psittacidae). <i>Acta Parasitologica</i> , 2008, 53, .	0.4	6
29	A new species of the genus <i>Coraciacarus</i> (Gabuciniidae, Pterolichoidea) from the huia <i>Heteralocha acutirostris</i> (Callaeatidae, Passeriformes), an extinct bird species from New Zealand. <i>Journal of Natural History</i> , 2008, 42, 2763-2776.	0.2	6
30	Two new species of the feather mite genus <i>Analges</i> Nitzsch, 1818 (Analgoidea: Analgidae) from accentors (Passeriformes: Prunellidae) – morphological descriptions with DNA barcode data. <i>Systematic and Applied Acarology</i> , 2018, 23, 2288.	0.5	6
31	Observations on the quill mites (Acari: Syringophilidae) from charadriiform birds. <i>Zootaxa</i> , 2006, 1156, 51.	0.2	6
32	New species of the feather mite genus <i>Protolichus</i> Trouessart, 1884 (Astigmata, Pterolichidae) from lories and lorikeets (Aves: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50,702 Td (Psittaciforme	1.1	5
33	Ultrastructural characterization of <i>Acarispora calculifera</i> n.gen., n.sp., a new microsporidium (Opisthokonta: Chytridiopsida) from the feather mite <i>Falculifer rostratus</i> (Astigmata: Pterolichoidea). <i>Acta Parasitologica</i> , 2015, 60, 200-10.	0.4	5
34	Unusual way of feeding by the deutonymph of <i>Neottialges evansi</i> (Actinotrichida, Astigmata,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50,702 Td (P of Morphology, 2016, 277, 1368-1389.	0.6	5
35	Molecular phylogeny of Eupodidae reveals that the family Cocceupodidae (Actinotrichida;) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50,702 Td (P 43-57.	0.7	5
36	Long-term stasis in acariform mites provides evidence for morphologically stable evolution: Molecular vs. morphological differentiation in Linopodes (Acariformes; Prostigmata). <i>Molecular Phylogenetics and Evolution</i> , 2021, 163, 107237.	1.2	5

#	ARTICLE	IF	CITATIONS
37	A revised diagnosis of the feather mite genus <i>Magimelia</i> Gaud, 1961 (Pterolichoidea: Pterolichidae.) Tj ETQq1 1 0.784314 rgBT /Overlock 10	0.5	4
38	A New Species of the Feather Mite Genus <i>Pterotrogus</i> Gaud (Analgoidea: Pteronyssidae) from the Ivory-billed Woodpecker <i>Campephilus principalis</i> L. (Aves: Piciformes). <i>Annals of the Entomological Society of America</i> , 2005, 98, 13-17.	1.3	4
39	New feather mite species (Acari, Astigmata) from the Sulphur-crested Cockatoo <i>Cacatua galerita</i> and Yellow-crested Cockatoo <i>C. sulphurea</i> (Psittaciformes, Cacatuidae). <i>Acta Parasitologica</i> , 2007, 52, 250.	0.4	4
40	<i>Anhemialges bakeri</i> sp. nov. (Analgoidea, Analgidae) – a new species of feather mite from the Common Chiffchaff <i>Phylloscopus collybita</i> (Passeriformes, Sylviidae) from England. <i>Acta Parasitologica</i> , 2010, 55, .	0.4	4
41	Systematic revision of the feather mite genus <i>Protolichus</i> Trouessart, 1884 (Astigmata, Pterolichidae). <i>Zootaxa</i> , 2010, 2526, .	0.2	4
42	A new feather mite species of the genus <i>Neumannella</i> Trouessart, 1916 (Analgoidea, Dermoglyphidae) from the Red-winged Tinamou <i>Rhynchotus rufescens</i> (Temminck, 1815) (Aves, Tinamiformes) with remarks to the evolution of host-parasite associations of the genus. <i>Acta Parasitologica</i> , 2014, 59, 197-205.	0.4	4
43	Feather mites of the genera <i>Dubininia</i> and <i>Cacatualges</i> (Acari: Xolalgidae) associated with parrots (Aves: Psittaciformes) of the Old World. <i>Zootaxa</i> , 2017, 4272, 451-490.	0.2	3
44	Delimiting species of water mites of the genus <i>Hydrodroma</i> (Acari: Hydrachnidiae: Hydrodromidae) from North America and Europe: Integrative evidence of species status from COI sequences and morphology. <i>Zoologischer Anzeiger</i> , 2020, 284, 16-29.	0.4	3
45	<i>Xoloptes blaszaki</i> sp. nov. (Pterolichoidea: Pterolichidae) – A New Species of Feather Mite from <i>Alectoris rufa</i> (Galliformes: Phasianidae) from Europe with DNA Barcode Data. <i>Annales Zoologici</i> , 2008, 58, 391-396.	0.1	2
46	A new species of the feather mite genus <i>Dubininia</i> Vassilev, 1958 (Acari: Xolalgidae) from the Black-thighed Falconet <i>Microhierax fringillarius</i> (Falconiformes: Falconidae). <i>Acta Parasitologica</i> , 2015, 60, 248-53.	0.4	2
47	The genetic structure of hypoderatid mites (Actinotrichida: Astigmata) parasitizing great cormorant (<i>Phalacrocorax carbo</i>) during host post-breeding dispersal in Milicz, SW Poland. <i>Acta Parasitologica</i> , 2017, 62, 76-89.	0.4	2
48	Two New Species of Mites (Acari: Myobiidae and Myocoptidae) from Skin and Fur of Small Mammals in Poland. <i>Annales Zoologici</i> , 2008, 58, 383-390.	0.1	1
49	<i>Szeptyckiana</i> , a new genus of the feather mite family Pterolichidae (Acari, Astigmata,) Tj ETQq1 1 0.784314 rgBT /Overlock 10	0.2	1
50	A NEW SPECIES OF THE FEATHER MITE GENUS <i>CALCEALGES</i> GAUD, 1952 (ACARIFORMES: TROUESSARTIIDAE) FROM THE LOWLAND TINY GREENBUL <i>PHYLLASTREPHUS DEBILIS</i> (PASSERIFORMES: PYCNONOTIDAE): MORPHOLOGICAL DESCRIPTIONS WITH DNA BARCODE DATA. <i>Acarina</i> , 2019, 27, 165-174.	0.1	1
51	<i>Syringoplutarchusia nordmanni</i> , a new genus and species of the feather mite family Syringobiidae (Acari, Astigmata, Pterolichoidea) from the Black-winged Pratincole <i>Glareola nordmanni</i> (Charadriiformes, Glareolidae). <i>Acta Parasitologica</i> , 2007, 52, 62.	0.4	0
52	Preliminary assessment of mating duration and prolonged post-copulatory associations in <i>Arrenurus</i> water mites (Actinotrichida: Parasitengonina: Hydrachnidiae). <i>Acarologia</i> , 2022, 62, 84-93.	0.2	0