

# Jacek Dabert

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/8832189/jacek-dabert-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

51  
papers

807  
citations

13  
h-index

27  
g-index

53  
ext. papers

926  
ext. citations

1.6  
avg, IF

4.04  
L-index

| #  | Paper   | IF  | Citations      |
|----|---|-----|----------------|
| 51 | Preliminary assessment of mating duration and prolonged post-copulatory associations in <i>Arrhenurus</i> water mites (Actinotrichida: Parasitengonina: Hydrachnidiae). <i>Acarologia</i> , <b>2022</b> , 62, 84-93   | 0.7 |                |
| 50 | Long-term stasis in acariform mites provides evidence for morphologically stable evolution: Molecular vs. morphological differentiation in Linopodes (Acariformes; Prostigmata). <i>Molecular Phylogenetics and Evolution</i> , <b>2021</b> , 163, 107237   | 4.1 | 2              |
| 49 | Sweet or salty? The origin of freshwater gastrotrichs (Gastrotricha, Chaetonotida) revealed by molecular phylogenetic analysis. <i>Cladistics</i> , <b>2020</b> , 36, 458-480   | 3.5 | 1              |
| 48 | Molecular phylogeny of Eupodidae reveals that the family Cocceupodidae (Actinotrichida; Eupodoidea) and its genus <i>Filieupodes</i> are valid taxa. <i>Experimental and Applied Acarology</i> , <b>2020</b> , 80, 43-57 <sup>2,1</sup>   |     | 4              |
| 47 | Delimiting species of water mites of the genus <i>Hydrodroma</i> (Acari: Hydrachnidiae: Hydrodromidae) from North America and Europe: Integrative evidence of species status from COI sequences and morphology. <i>Zoologischer Anzeiger</i> , <b>2020</b> , 284, 16-29   | 1.1 | 3              |
| 46 | A NEW SPECIES OF THE FEATHER MITE GENUS <i>CALCEALGES</i> GAUD, 1952 (ACARIFORMES: TROUESSARTIIDAE) FROM THE LOWLAND TINY GREENBUL <i>PHYLLASTREPHUS</i> DEBILIS (PASSERIFORMES: PYCNONOTIDAE): MORPHOLOGICAL DESCRIPTIONS WITH DNA BARCODE DATA. <i>Acarina</i> , <b>2019</b> , 27, 165-174  | 0.5 |                |
| 45 | Two new species of the feather mite genus <i>Analges</i> Nitzsch, 1818 (Analgoidea: Analgidae) from accentors (Passeriformes: Prunellidae) [morphological descriptions with DNA barcode data. <i>Systematic and Applied Acarology</i> , <b>2018</b> , 23, 2288  | 0.8 | 3              |
| 44 | The genetic structure of hypoderatid mites (Actinotrichida: Astigmata) parasitizing great cormorant ( <i>Phalacrocorax carbo</i> ) during host post-breeding dispersal in Milicz, SW Poland. <i>Acta Parasitologica</i> , <b>2017</b> , 62, 76-89   | 1.7 | 2              |
| 43 | Feather mites of the genera <i>Dubininia</i> and <i>Cacatualges</i> (Acari: Xolalgidae) associated with parrots (Aves: Psittaciformes) of the Old World. <i>Zootaxa</i> , <b>2017</b> , 4272, 451-490   | 0.5 | 1              |
| 42 | Unusual way of feeding by the deutonymph of <i>Neottialges evansi</i> (Actinotrichida, Astigmata, Hypoderatidae), a subcutaneous parasite of cormorants, revealed by fine structural analyses. <i>Journal of Morphology</i> , <b>2016</b> , 277, 1368-89  | 1.6 | 4              |
| 41 | <i>Bifidochaetus</i> , a new Arctic genus of freshwater Chaetonotida (Gastrotricha) from Spitsbergen revealed by an integrative taxonomic approach. <i>Invertebrate Systematics</i> , <b>2016</b> , 30, 398   | 1.2 | 18             |
| 40 | Higher-level molecular phylogeny of the water mites (Acariformes: Prostigmata: Parasitengonina: Hydrachnidiae). <i>Molecular Phylogenetics and Evolution</i> , <b>2016</b> , 101, 75-90   | 4.1 | 3 <sup>1</sup> |
| 39 | A new species of the feather mite genus <i>Dubininia</i> Vassilev, 1958 (Acari: Xolalgidae) from the Black-thighed Falconet <i>Microhierax fringillarius</i> (Falconiformes: Falconidae). <i>Acta Parasitologica</i> , <b>2015</b> , 60, 248-53   | 1.7 | 1              |
| 38 | Ultrastructural characterization of <i>Acarispora falculifera</i> n.gen., n.sp., a new microsporidium (Opisthokonta: Chytridiopsida) from the feather mite <i>Falculifer rostratus</i> (Astigmata: Pterolichoidea). <i>Acta Parasitologica</i> , <b>2015</b> , 60, 200-10   | 1.7 | 5              |
| 37 | Differences in speciation progress in feather mites (Analgoidea) inhabiting the same host: the case of <i>Zachvatkinia</i> and <i>Alloptes</i> living on arctic and long-tailed skuas. <i>Experimental and Applied Acarology</i> , <b>2015</b> , 65, 163-79   | 2.1 | 20             |
| 36 | Phylogenetic position of the enigmatic clawless eutardigrade genus <i>Apodibius</i> Dastych, 1983 (Tardigrada), based on 18S and 28S rRNA sequence data from its type species <i>A. confusus</i> . <i>Molecular Phylogenetics and Evolution</i> , <b>2014</b> , 70, 70-5  | 4.1 | 21             |
| 35 | A new feather mite species of the genus <i>Neumannella</i> Trouessart, 1916 (Analgoidea, Dermoglyphidae) from the Red-winged Tinamou <i>Rhynchotus rufescens</i> (Temminck, 1815) (Aves, Tinamiformes) with remarks to the evolution of host-parasite associations of the genus. <i>Acta Parasitologica</i> , <b>2014</b> , 59, 197-205 | 1.7 | 3              |

|    |   |     |     |
|----|---|-----|-----|
| 34 | New species of the feather mite genus <i>Protolichus</i> Trouessart, 1884 (Astigmata, Pterolichidae) from lorries and lorikeets (Aves: Psittaciformes). <i>Zootaxa</i> , <b>2014</b> , 3774, 131-51   | 0.5 | 5   |
| 33 | DNA barcodes reveal female dimorphism in syringophilid mites (Actinotrichida: Prostigmata: Cheyletoidea): <i>Stibarokris phoeniconaias</i> and <i>Ciconichenophilus phoeniconaias</i> are conspecific. <i>Folia Parasitologica</i> , <b>2014</b> , 61, 272-6                            | 1.8 | 12  |
| 32 | Multidisciplinary analysis of <i>Knemidocoptes jamaicensis</i> parasitising the Common Chaffinch, <i>Fringilla coelebs</i> : proofs for a multispecies complex?. <i>Parasitology Research</i> , <b>2013</b> , 112, 2373-80  | 2.4 | 15  |
| 31 | DNA-barcoding contradicts morphology in quill mite species <i>Torotroglia merulae</i> and <i>T. rubeculi</i> (Prostigmata: Syringophilidae). <i>Folia Parasitologica</i> , <b>2013</b> , 60, 51-60  | 1.8 | 11  |
| 30 | A new feather mite species of the genus <i>Proctophyllodes</i> Robin, 1877 (Astigmata: Proctophyllodidae) from the Long-tailed Tit <i>Aegithalos caudatus</i> (Passeriformes: Aegithalidae) morphological description with DNA barcode data. <i>Zootaxa</i> , <b>2012</b> , 3253, 54    | 0.5 | 111 |
| 29 | A new quill mite <i>Syringophiloidus pseudonigritae</i> sp. nov. (Prostigmata, Syringophilidae) parasitizing <i>Pseudonigrita arnaudi</i> (Passeriformes, Ploceidae) combined description using morphology and DNA barcode data. <i>Zootaxa</i> , <b>2012</b> , 3532, 64                | 0.5 | 11  |
| 28 | <i>Picobia dziabaszweskii</i> sp. nov. (Acari, Syringophilidae) combined description (morphology with DNA barcode data) of a new quill mite species parasitizing <i>Garrulax formosus</i> (Passeriformes: Leiothrichidae). <i>Zootaxa</i> , <b>2012</b> , 3224, 57                      | 0.5 | 12  |
| 27 | The first report of <i>Knemidocoptes intermedius</i> Fain et Macfarlane, 1967 (Acari: Astigmata) in naturally infected European birds. <i>Parasitology Research</i> , <b>2011</b> , 109, 237-40   | 2.4 | 13  |
| 26 | Systematic revision of the feather mite genus <i>Protolichus</i> Trouessart, 1884 (Astigmata, Pterolichidae). <i>Zootaxa</i> , <b>2010</b> , 2526, 1  | 0.5 | 2   |
| 25 | <i>Anhemialges bakeri</i> sp. nov. (Analgoidea, Analgidae) a new species of feather mite from the Common Chiffchaff <i>Phylloscopus collybita</i> (Passeriformes, Sylviidae) from England. <i>Acta Parasitologica</i> , <b>2010</b> , 55,   | 1.7 | 2   |
| 24 | Molecular evidence for species separation in the water mite <i>Hygrobatas nigromaculatus</i> Lebert, 1879 (Acari, Hydrachnidia): evolutionary consequences of the loss of larval parasitism. <i>Aquatic Sciences</i> , <b>2010</b> , 72, 347-360  | 2.5 | 49  |
| 23 | Molecular phylogeny of acariform mites (Acari, Arachnida): strong conflict between phylogenetic signal and long-branch attraction artifacts. <i>Molecular Phylogenetics and Evolution</i> , <b>2010</b> , 56, 222-41  | 4.1 | 209 |
| 22 | A New Species of the Genus <i>Proctophyllodes</i> (Analgoidea: Proctophyllodidae) from Cetti's Warbler <i>Cettia cetti</i> (Passeriformes: Sylviidae) with DNA Barcode Data. <i>Annales Zoologici</i> , <b>2008</b> , 58, 397-402   | 0.6 | 10  |
| 21 | A new species of the genus <i>Coraciacarus</i> (Gabuciniidae, Pterolichoidea) from the huia <i>Heteralocha acutirostris</i> (Callaeatidae, Passeriformes), an extinct bird species from New Zealand. <i>Journal of Natural History</i> , <b>2008</b> , 42, 2763-2776                    | 0.5 | 5   |
| 20 | Two New Species of Mites (Acari: Myobiidae and Myocoptidae) from Skin and Fur of Small Mammals in Poland. <i>Annales Zoologici</i> , <b>2008</b> , 58, 383-390  | 0.6 | 1   |
| 19 | <i>Xoloptes blaszakisp.</i> nov. (Pterolichoidea: Pterolichidae) A New Species of Feather Mite from <i>Alectoris rufa</i> (Galliformes: Phasianidae) from Europe with DNA Barcode Data. <i>Annales Zoologici</i> , <b>2008</b> , 58, 391-396  | 0.6 | 2   |
| 18 | Systematic revision of the feather mite genera <i>Apexolichus</i> Gaud et Atyeo and <i>Titanolichus</i> Gaud et Atyeo (Astigmata, Pterolichidae), parasites of parrots of the Old World (Psittaciformes, Psittacidae). <i>Acta Parasitologica</i> , <b>2008</b> , 53,                   | 1.7 | 1   |
| 17 | <i>Syringoplutarchusia nordmanni</i> , a new genus and species of the feather mite family Syringobiidae (Acari, Astigmata, Pterolichoidea) from the Black-winged Pratincole <i>Glareola nordmanni</i> (Charadriiformes, Glareolidae). <i>Acta Parasitologica</i> , <b>2007</b> , 52, 62 | 1.7 |     |

|    |  |     |    |
|----|--|-----|----|
| 16 | New feather mite species (Acari, Astigmata) from the Sulphur-crested Cockatoo <i>Cacatua galerita</i> and Yellow-crested Cockatoo <i>C. sulphurea</i> (Psittaciformes, Cacatuidae). <i>Acta Parasitologica</i> , <b>2007</b> , 52, 250   | 1.7 | 2  |
| 15 | Three new feather mite genera of the <i>Protolichus</i> generic group (Astigmata, Pterolichidae) from parrots (Aves, Psittaciformes) of the Old World. <i>Acta Parasitologica</i> , <b>2007</b> , 52, 386  | 1.7 | 3  |
| 14 | A new species of the feather mite genus <i>Titanolichus</i> Gaud & Atyeo, 1996 (Acari: Astigmata: Pterolichidae) from the endangered orange-bellied parrot <i>Neophema chrysogaster</i> (Aves: Psittaciformes) from Australia. <i>Australian Journal of Entomology</i> , <b>2006</b> , 45, 206-214 |     | 5  |
| 13 | Observations on the quill mites (Acari: Syringophilidae) from charadriiform birds. <i>Zootaxa</i> , <b>2006</b> , 1156, 51   | 0.5 | 5  |
| 12 | Six new feather mite species (Acari: Astigmata) from the carolina parakeet <i>Conuropsis carolinensis</i> (Psittaciformes: Psittacidae), an extinct parrot of North America. <i>Journal of Natural History</i> , <b>2005</b> , 39, 2257-2278   | 0.5 | 17 |
| 11 | A New Species of the Feather Mite Genus <i>Pterotrogus</i> Gaud (Analgoidea: Pteronyssidae) from the Ivory-billed Woodpecker <i>Campephilus principalis</i> L. (Aves: Piciformes). <i>Annals of the Entomological Society of America</i> , <b>2005</b> , 98, 13-17                                 | 2   | 4  |
| 10 | New feather mite taxa of the <i>Rhytidelasma</i> Gaud, 1966 generic group (Astigmata: Pterolichidae) from the red-flanked lorikeet <i>Charmosyna placentis placentis</i> (Psittacidae). <i>Systematic Parasitology</i> , <b>2004</b> , 58, 91-104  | 1   | 4  |
| 9  | New feather mites of the family Pterolichidae (Acari: Pterolichoidea) from parrots (Aves: Psittaciformes) in Australia. <i>Australian Journal of Entomology</i> , <b>2003</b> , 42, 185-202  |     | 11 |
| 8  | A revised diagnosis of the feather mite genus <i>Magimelia</i> Gaud, 1961 (Pterolichoidea: Pterolichidae: Magimeliinae) and the description of three new species. <i>Systematic Parasitology</i> , <b>2002</b> , 53, 69-79   | 1   | 3  |
| 7  | Phylogeny of feather mite subfamily Avenzoariinae (Acari: Analgoidea: Avenzoariidae) inferred from combined analyses of molecular and morphological data. <i>Molecular Phylogenetics and Evolution</i> , <b>2001</b> , 20, 124-35  | 4.1 | 27 |
| 6  | <i>Picobia polonica</i> sp. n. (Acari: Prostigmata: Syringophilidae), a new species of quill mite from the domestic hen, <i>Gallus gallus domesticus</i> (Aves: Phasianidae). <i>Folia Parasitologica</i> , <b>2001</b> , 48, 154-8  | 1.8 | 6  |
| 5  | Phylogeny and co-speciation in feather mites of the subfamily Avenzoariinae (Analgoidea: Avenzoariidae) <b>1999</b> , 23, 525-549  |     | 16 |
| 4  | Origin and Evolution of Feather Mites (Astigmata). <i>Experimental and Applied Acarology</i> , <b>1999</b> , 23, 437-454.  | 4.1 | 82 |
| 3  | RAPD technique for taxonomic studies of <i>Pellia epiphylla</i> -complex (Hepaticae, Metzgeriales). <i>Genetica</i> , <b>1998</b> , 104, 179-87  | 1.5 | 12 |
| 2  | Morphological adaptation for precopulatory guarding in astigmatic mites (Acari: Acaridida). <i>International Journal of Acarology</i> , <b>1992</b> , 18, 49-54  | 0.6 | 17 |
| 1  | Species boundaries among extremely diverse and sexually dimorphic <i>Arrenurus</i> water mites (Acariformes: Hydrachnidia: Arrenuridae)  |     | 1  |