

# Juliana Carlota Kramer Soares

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8839395/publications.pdf>

Version: 2024-02-01

15  
papers

308  
citations

932766

10  
h-index

1058022

14  
g-index

15  
all docs

15  
docs citations

15  
times ranked

551  
citing authors

#	ARTICLE	IF	CITATIONS
1	Anterior thalamus deep brain stimulation at high current impairs memory in rats. <i>Experimental Neurology</i> , 2010, 225, 154-162.	2.0	71
2	Role of muscarinic M1 receptors in inhibitory avoidance and contextual fear conditioning. <i>Neurobiology of Learning and Memory</i> , 2006, 86, 188-196.	1.0	40
3	A single bout of resistance exercise improves memory consolidation and increases the expression of synaptic proteins in the hippocampus. <i>Hippocampus</i> , 2016, 26, 1096-1103.	0.9	29
4	Effects of nociceptin/orphanin FQ in the acquisition of contextual and tone fear conditioning in rats.. <i>Behavioral Neuroscience</i> , 2008, 122, 98-106.	0.6	23
5	Hippocampal microRNA-mRNA regulatory network is affected by physical exercise. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2018, 1862, 1711-1720.	1.1	23
6	Effects of Long-Term Ayahuasca Administration on Memory and Anxiety in Rats. <i>PLoS ONE</i> , 2015, 10, e0145840.	1.1	21
7	Network supporting contextual fear learning after dorsal hippocampal damage has increased dependence on retrosplenial cortex. <i>PLoS Computational Biology</i> , 2018, 14, e1006207.	1.5	20
8	Effects of pre- or post-training entorhinal cortex AP5 injection on fear conditioning. <i>Physiology and Behavior</i> , 2005, 86, 508-515.	1.0	19
9	Hippocampal NMDA receptor blockade impairs CREB phosphorylation in amygdala after contextual fear conditioning. <i>Hippocampus</i> , 2013, 23, 545-551.	0.9	14
10	Inactivation of muscarinic receptors impairs place and response learning: Implications for multiple memory systems. <i>Neuropharmacology</i> , 2013, 73, 320-326.	2.0	13
11	Involvement of the prelimbic cortex in contextual fear conditioning with temporal and spatial discontinuity. <i>Neurobiology of Learning and Memory</i> , 2017, 144, 1-10.	1.0	10
12	Functional interaction of ventral hippocampal CA1 region and prelimbic cortex contributes to the encoding of contextual fear association of stimuli separated in time. <i>Neurobiology of Learning and Memory</i> , 2020, 171, 107216.	1.0	10
13	M1 muscarinic receptors are necessary for retrieval of remote context fear memory. <i>Physiology and Behavior</i> , 2017, 169, 202-207.	1.0	9
14	Effects of the M1 muscarinic antagonist dicyclomine on emotional memory retrieval.. <i>Behavioral Neuroscience</i> , 2016, 130, 29-35.	0.6	6
15	Decrease of ARC protein expression in the striatum after tone fear conditioning. <i>Revista Neurociencias</i> , 0, 29, 1-22.	0.0	0