

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/8844673/istvan-nagy-publications-by-citations.pdf>  
**Version:** 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

88 papers	3,779 citations	32 h-index	60 g-index
92 ext. papers	4,132 ext. citations	5.8 avg, IF	4.89 L-index

#	Paper	IF	Citations
88	2010 International consensus algorithm for the diagnosis, therapy and management of hereditary angioedema. <i>Allergy, Asthma and Clinical Immunology</i> , <b>2010</b> , 6, 24	3.2	333
87	Cannabinoid 1 receptors are expressed in nociceptive primary sensory neurons. <i>Neuroscience</i> , <b>2000</b> , 100, 685-8	3.9	253
86	TRPV4: Molecular Conductor of a Diverse Orchestra. <i>Physiological Reviews</i> , <b>2016</b> , 96, 911-73	47.9	206
85	Vanilloid receptor 1 expression in the rat urinary tract. <i>Neuroscience</i> , <b>2002</b> , 109, 787-98	3.9	201
84	The role of the vanilloid (capsaicin) receptor (TRPV1) in physiology and pathology. <i>European Journal of Pharmacology</i> , <b>2004</b> , 500, 351-69	5.3	197
83	Anandamide-evoked activation of vanilloid receptor 1 contributes to the development of bladder hyperreflexia and nociceptive transmission to spinal dorsal horn neurons in cystitis. <i>Journal of Neuroscience</i> , <b>2004</b> , 24, 11253-63	6.6	164
82	Hereditary angioedema: a current state-of-the-art review, VII: Canadian Hungarian 2007 International Consensus Algorithm for the Diagnosis, Therapy, and Management of Hereditary Angioedema. <i>Annals of Allergy, Asthma and Immunology</i> , <b>2008</b> , 100, S30-40	3.2	162
81	Canadian 2003 International Consensus Algorithm For the Diagnosis, Therapy, and Management of Hereditary Angioedema. <i>Journal of Allergy and Clinical Immunology</i> , <b>2004</b> , 114, 629-37	11.5	155
80	Anandamide regulates neuropeptide release from capsaicin-sensitive primary sensory neurons by activating both the cannabinoid 1 receptor and the vanilloid receptor 1 in vitro. <i>European Journal of Neuroscience</i> , <b>2003</b> , 17, 2611-8	3.5	151
79	Capsaicin-sensitive sensory fibers in the islets of Langerhans contribute to defective insulin secretion in Zucker diabetic rat, an animal model for some aspects of human type 2 diabetes. <i>European Journal of Neuroscience</i> , <b>2007</b> , 25, 213-23	3.5	123
78	Similarities and differences between the responses of rat sensory neurons to noxious heat and capsaicin. <i>Journal of Neuroscience</i> , <b>1999</b> , 19, 10647-55	6.6	119
77	Noxious heat activates all capsaicin-sensitive and also a sub-population of capsaicin-insensitive dorsal root ganglion neurons. <i>Neuroscience</i> , <b>1999</b> , 88, 995-7	3.9	118
76	The role of neurokinin and N-methyl-D-aspartate receptors in synaptic transmission from capsaicin-sensitive primary afferents in the rat spinal cord in vitro. <i>Neuroscience</i> , <b>1993</b> , 52, 1029-37	3.9	97
75	Systemic inflammation enhances surgery-induced cognitive dysfunction in mice. <i>Neuroscience Letters</i> , <b>2011</b> , 498, 63-6	3.3	95
74	Lignocaine selectively reduces C fibre-evoked neuronal activity in rat spinal cord in vitro by decreasing N-methyl-D-aspartate and neurokinin receptor-mediated post-synaptic depolarizations; implications for the development of novel centrally acting analgesics. <i>Pain</i> , <b>1996</b> , 64, 59-70	8	87
73	Inflammatory mediators convert anandamide into a potent activator of the vanilloid type 1 transient receptor potential receptor in nociceptive primary sensory neurons. <i>Neuroscience</i> , <b>2005</b> , 136, 539-48	3.9	71
72	Activation of capsaicin-sensitive primary sensory neurones induces anandamide production and release. <i>Journal of Neurochemistry</i> , <b>2003</b> , 84, 585-91	6	68

71	Possible mechanisms of cannabinoid-induced antinociception in the spinal cord. <i>European Journal of Pharmacology</i> , <b>2001</b> , 429, 93-100	5.3	67
70	Transient receptor potential ion channels in primary sensory neurons as targets for novel analgesics. <i>British Journal of Pharmacology</i> , <b>2014</b> , 171, 2508-27	8.6	64
69	Cannabinoid 1 receptors are expressed by nerve growth factor- and glial cell-derived neurotrophic factor-responsive primary sensory neurones. <i>Neuroscience</i> , <b>2002</b> , 110, 747-53	3.9	63
68	Functional transient receptor potential vanilloid 1 is expressed in human urothelial cells. <i>Journal of Urology</i> , <b>2009</b> , 182, 2944-50	2.5	56
67	The putative role of vanilloid receptor-like protein-1 in mediating high threshold noxious heat-sensitivity in rat cultured primary sensory neurons. <i>European Journal of Neuroscience</i> , <b>2002</b> , 16, 1483-9	3.5	53
66	TRPV1 function in health and disease. <i>Current Pharmaceutical Biotechnology</i> , <b>2011</b> , 12, 130-44	2.6	49
65	Localization of the endocannabinoid-degrading enzyme fatty acid amide hydrolase in rat dorsal root ganglion cells and its regulation after peripheral nerve injury. <i>Journal of Neuroscience</i> , <b>2009</b> , 29, 3766-80	6.6	46
64	The distribution of sensory fibers immunoreactive for the TRPV1 (capsaicin) receptor in the human prostate. <i>European Urology</i> , <b>2005</b> , 48, 162-7	10.2	43
63	Cannabinoid 1 receptor activation inhibits transient receptor potential vanilloid type 1 receptor-mediated cationic influx into rat cultured primary sensory neurons. <i>Neuroscience</i> , <b>2009</b> , 162, 1202-11	3.9	41
62	Characterisation of cannabinoid 1 receptor expression in the perikarya, and peripheral and spinal processes of primary sensory neurons. <i>Brain Structure and Function</i> , <b>2013</b> , 218, 733-50	4	40
61	Insulin induces cobalt uptake in a subpopulation of rat cultured primary sensory neurons. <i>European Journal of Neuroscience</i> , <b>2003</b> , 18, 2477-86	3.5	40
60	Cobalt accumulation in neurons expressing ionotropic excitatory amino acid receptors in young rat spinal cord: morphology and distribution. <i>Journal of Comparative Neurology</i> , <b>1994</b> , 344, 321-35	3.4	36
59	Pharmacology of the capsaicin receptor, transient receptor potential vanilloid type-1 ion channel. <i>Progress in Drug Research Fortschritte Der Arzneimittelforschung Progres Des Recherches Pharmaceutiques</i> , <b>2014</b> , 68, 39-76		36
58	NK1 and NK2 receptors contribute to C-fibre evoked slow potentials in the spinal cord. <i>NeuroReport</i> , <b>1994</b> , 5, 2105-8	1.7	35
57	Cobalt uptake enables identification of capsaicin- and bradykinin-sensitive subpopulations of rat dorsal root ganglion cells in vitro. <i>Neuroscience</i> , <b>1993</b> , 56, 241-6	3.9	34
56	Neurochemical characterization of insulin receptor-expressing primary sensory neurons in wild-type and vanilloid type 1 transient receptor potential receptor knockout mice. <i>Journal of Comparative Neurology</i> , <b>2007</b> , 503, 334-47	3.4	32
55	Spinal neurokinin NK1 receptor down-regulation and antinociception: effects of spinal NK1 receptor antisense oligonucleotides and NK1 receptor occupancy. <i>Journal of Neurochemistry</i> , <b>1998</b> , 70, 688-98	6	30
54	Role of transient receptor potential and acid-sensing ion channels in peripheral inflammatory pain. <i>Anesthesiology</i> , <b>2010</b> , 112, 729-41	4.3	29

53	Spatial distribution of pre- and postsynaptic sites of axon terminals in the dorsal horn of the frog spinal cord. <i>Neuroscience</i> , <b>1989</b> , 29, 175-88	3.9	28
52	Peripheral orthopaedic surgery down-regulates hippocampal brain-derived neurotrophic factor and impairs remote memory in mouse. <i>Neuroscience</i> , <b>2011</b> , 190, 194-9	3.9	26
51	Peripheral mechanisms of burn injury-associated pain. <i>European Journal of Pharmacology</i> , <b>2013</b> , 716, 169-78	5.3	22
50	The endogenous cannabinoid anandamide inhibits transient receptor potential vanilloid type 1 receptor-mediated currents in rat cultured primary sensory neurons. <i>Acta Physiologica Hungarica</i> , <b>2010</b> , 97, 149-58		22
49	Extracellular signal-regulated kinases in pain of peripheral origin. <i>European Journal of Pharmacology</i> , <b>2011</b> , 650, 8-17	5.3	16
48	Cystitis is associated with TRPV1b-downregulation in rat dorsal root ganglia. <i>NeuroReport</i> , <b>2008</b> , 19, 1469-72	1.7	16
47	Spatial Distribution of the Cannabinoid Type 1 and Capsaicin Receptors May Contribute to the Complexity of Their Crosstalk. <i>Scientific Reports</i> , <b>2016</b> , 6, 33307	4.9	13
46	Isoflurane causes neocortical but not hippocampal-dependent memory impairment in mice. <i>Acta Anaesthesiologica Scandinavica</i> , <b>2012</b> , 56, 1052-7	1.9	13
45	Anandamide in primary sensory neurons: too much of a good thing?. <i>European Journal of Neuroscience</i> , <b>2014</b> , 39, 409-18	3.5	13
44	Morphological and membrane properties of young rat lumbar and thoracic dorsal root ganglion cells with unmyelinated axons. <i>Brain Research</i> , <b>1993</b> , 609, 193-200	3.7	13
43	Capsaicin-sensitive primary sensory neurons in the mouse express N-Acyl phosphatidylethanolamine phospholipase D. <i>Neuroscience</i> , <b>2009</b> , 161, 572-7	3.9	12
42	Comparison of currents activated by noxious heat in rat and chicken primary sensory neurons. <i>Regulatory Peptides</i> , <b>2000</b> , 96, 3-6		12
41	Prolonged exposure to bradykinin and prostaglandin E2 increases TRPV1 mRNA but does not alter TRPV1 and TRPV1b protein expression in cultured rat primary sensory neurons. <i>Neuroscience Letters</i> , <b>2014</b> , 564, 89-93	3.3	11
40	Effects of cannabinoids on capsaicin receptor activity following exposure of primary sensory neurons to inflammatory mediators. <i>Life Sciences</i> , <b>2010</b> , 87, 162-8	6.8	11
39	Alterations of substance P immunoreactivity in lumbar and thoracic segments of rat spinal cord in ultraviolet irradiation induced hyperalgesia of the hindpaw. <i>Brain Research</i> , <b>1998</b> , 786, 248-51	3.7	10
38	Anandamide produced by Ca(2+)-insensitive enzymes induces excitation in primary sensory neurons. <i>Pflügers Archiv European Journal of Physiology</i> , <b>2014</b> , 466, 1421-35	4.6	9
37	The Insulin Receptor Is Colocalized With the TRPV1 Nociceptive Ion Channel and Neuropeptides in Pancreatic Spinal and Vagal Primary Sensory Neurons. <i>Pancreas</i> , <b>2018</b> , 47, 110-115	2.6	9
36	Phosphorylated Histone 3 at Serine 10 Identifies Activated Spinal Neurons and Contributes to the Development of Tissue Injury-Associated Pain. <i>Scientific Reports</i> , <b>2017</b> , 7, 41221	4.9	8

35	Is there a nociceptive carousel?. <i>Trends in Pharmacological Sciences</i> , <b>1997</b> , 18, 223-224	13.2	7
34	The effects of NK-1 and NK-2 receptor antagonists on the capsaicin evoked synaptic response in the rat spinal cord in vitro. <i>Regulatory Peptides</i> , <b>1993</b> , 46, 413-4		7
33	The NA1.7 blocker protoxin II reduces burn injury-induced spinal nociceptive processing. <i>Journal of Molecular Medicine</i> , <b>2018</b> , 96, 75-84	5.5	7
32	Insulin Confers Differing Effects on Neurite Outgrowth in Separate Populations of Cultured Dorsal Root Ganglion Neurons: The Role of the Insulin Receptor. <i>Frontiers in Neuroscience</i> , <b>2018</b> , 12, 732	5.1	7
31	Peripheral inflammation affects modulation of nociceptive synaptic transmission in the spinal cord induced by N-arachidonoylphosphatidylethanolamine. <i>British Journal of Pharmacology</i> , <b>2018</b> , 175, 2322-2336	8.6	6
30	TRPV1 Antagonists as Novel Anti-Diabetic Agents: Regulation of Oral Glucose Tolerance and Insulin Secretion Through Reduction of Low-Grade Inflammation?. <i>Medical Sciences (Basel, Switzerland)</i> , <b>2019</b> , 7,	3.3	6
29	Severe burn injury induces a characteristic activation of extracellular signal-regulated kinase 1/2 in spinal dorsal horn neurons. <i>European Journal of Pain</i> , <b>2011</b> , 15, 683-90	3.7	6
28	Molecular structure of transient receptor potential vanilloid type 1 ion channel (TRPV1). <i>Current Pharmaceutical Biotechnology</i> , <b>2011</b> , 12, 115-21	2.6	6
27	Molecular Mechanisms of TRPV1-Mediated Pain. <i>NeuroImmune Biology</i> , <b>2009</b> , 8, 75-99		6
26	Inflammation of peripheral tissues and injury to peripheral nerves induce differing effects in the expression of the calcium-sensitive N-arachidonoyl ethanolamine-synthesizing enzyme and related molecules in rat primary sensory neurons. <i>Journal of Comparative Neurology</i> , <b>2017</b> , 525, 1778-1796	3.4	5
25	Microdialysis Workflow for Metabotyping Superficial Pathologies: Application to Burn Injury. <i>Analytical Chemistry</i> , <b>2019</b> , 91, 6541-6548	7.8	5
24	Xenon fails to inhibit capsaicin-evoked CGRP release by nociceptors in culture. <i>Neuroscience Letters</i> , <b>2011</b> , 499, 124-6	3.3	5
23	CB receptor-dependent desensitisation of TRPV1 channels contributes to the analgesic effect of dipyrone in sensitised primary sensory neurons. <i>British Journal of Pharmacology</i> , <b>2020</b> , 177, 4615-4626	8.6	4
22	Cyclin-dependent-like kinase 5 is required for pain signaling in human sensory neurons and mouse models. <i>Science Translational Medicine</i> , <b>2020</b> , 12,	17.5	4
21	Chronic neuropathic pain: Pathomechanism and pharmacology. <i>Drug Development Research</i> , <b>2001</b> , 54, 159-166	5.1	4
20	Imidazoline ligand BU224 reverses cognitive deficits, reduces microgliosis and enhances synaptic connectivity in a mouse model of Alzheimer's disease. <i>British Journal of Pharmacology</i> , <b>2021</b> , 178, 654-671	8.6	4
19	The insulin receptor is differentially expressed in somatic and visceral primary sensory neurons. <i>Cell and Tissue Research</i> , <b>2018</b> , 374, 243-249	4.2	3
18	Immunohistochemical localization of neurokinin-1 receptor in the lumbar spinal cord of young rats: morphology and distribution. <i>Somatosensory &amp; Motor Research</i> , <b>1999</b> , 16, 361-8	1.2	3

17	Possible branching of myelinated primary afferent fibres in the dorsal root of the rat. <i>Brain Research</i> , <b>1995</b> , 703, 223-6	3.7	3
16	Combination of cobalt labelling with immunocytochemical reactions for electron microscopic investigations on frog spinal cord. <i>Microscopy Research and Technique</i> , <b>1994</b> , 28, 60-6	2.8	3
15	Fatty acid amide hydrolase inhibition normalises bladder function and reduces pain through normalising the anandamide/palmitoylethanolamine ratio in the inflamed bladder of rats. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , <b>2020</b> , 393, 263-272	3.4	3
14	Xenon reduces activation of transient receptor potential vanilloid type 1 (TRPV1) in rat dorsal root ganglion cells and in human TRPV1-expressing HEK293 cells. <i>Life Sciences</i> , <b>2011</b> , 88, 141-9	6.8	2
13	Taking the sting out of pain. <i>British Journal of Pharmacology</i> , <b>2007</b> , 151, 721-2	8.6	2
12	Leptin and fractalkine: novel subcutaneous cytokines in burn injury. <i>DMM Disease Models and Mechanisms</i> , <b>2020</b> , 13,	4.1	2
11	TRPV1 feed-forward sensitisation depends on COX2 upregulation in primary sensory neurons. <i>Scientific Reports</i> , <b>2021</b> , 11, 3514	4.9	2
10	Histone post-translational modifications as potential therapeutic targets for pain management. <i>Trends in Pharmacological Sciences</i> , <b>2021</b> , 42, 897-911	13.2	2
9	Functional Molecular Biology of the TRPV1 Ion Channel <b>2008</b> , 101-130		2
8	Sensitization of the transient receptor potential vanilloid type 1 ion channel by isoflurane or sevoflurane does not result in extracellular signal-regulated kinase 1/2 activation in rat spinal dorsal horn neurons. <i>Neuroscience</i> , <b>2010</b> , 166, 633-8	3.9	1
7	An Historical Introduction to the Endocannabinoid and Endovanilloid Systems <b>2008</b> , 3-13		1
6	Mechanisms underlying joint pain. <i>Drug Discovery Today Disease Mechanisms</i> , <b>2006</b> , 3, 357-363		1
5	Lactate dehydrogenase activity staining demonstrates time-dependent immune cell infiltration in human ex-vivo burn-injured skin. <i>Scientific Reports</i> , <b>2021</b> , 11, 21249	4.9	1
4	Spinal Excitatory Dynorphinergic Interneurons Contribute to Burn Injury-Induced Nociception Mediated by Phosphorylated Histone 3 at Serine 10 in Rodents. <i>International Journal of Molecular Sciences</i> , <b>2021</b> , 22,	6.3	1
3	Rapid genotyping of genetically modified laboratory animals from whole blood samples without DNA preparation. <i>Acta Biologica Hungarica</i> , <b>2013</b> , 64, 262-5		
2	Hyperexcitability in the Spinal Dorsal Horn: Cooperation of Neuropeptides and Excitatory Amino Acids <b>1994</b> , 379-399		
1	Rare co-occurrence of multiple sclerosis and Wilson's disease - case report.. <i>BMC Neurology</i> , <b>2022</b> , 22, 178	3.1	