

# Minori Koga

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8845199/publications.pdf>

Version: 2024-02-01

15  
papers

112  
citations

1478505

6  
h-index

1474206

9  
g-index

15  
all docs

15  
docs citations

15  
times ranked

172  
citing authors

#	ARTICLE	IF	CITATIONS
1	Risk and resilience in trajectories of post-traumatic stress symptoms among first responders after the 2011 Great East Japan Earthquake: 7-year prospective cohort study. <i>British Journal of Psychiatry</i> , 2022, 221, 668-675.	2.8	4
2	Neural extracellular vesicle-derived miR-17 in blood as a potential biomarker of subthreshold depression. <i>Human Cell</i> , 2021, 34, 1087-1092.	2.7	9
3	Validity and reliability of the Secondary Traumatic Stress Scale—Japanese version. <i>Neuropsychopharmacology Reports</i> , 2021, 41, 476-484.	2.3	3
4	Plasma fatty acid-binding protein 7 concentration correlates with depression/anxiety, cognition, and positive symptom in patients with schizophrenia. <i>Journal of Psychiatric Research</i> , 2021, 144, 304-311.	3.1	5
5	Psychological and physical condition of Japan maritime self-defense force personnel who performed disaster-relief missions after the 2011 great east Japan earthquake. <i>Journal of Psychiatric Research</i> , 2020, 130, 104-111.	3.1	2
6	Impact of a Rice-Centered Diet on the Quality of Sleep in Association with Reduced Oxidative Stress: A Randomized, Open, Parallel-Group Clinical Trial. <i>Nutrients</i> , 2020, 12, 2926.	4.1	2
7	Assessment of Factors Associated With Long-term Posttraumatic Stress Symptoms Among 56 388 First Responders After the 2011 Great East Japan Earthquake. <i>JAMA Network Open</i> , 2020, 3, e2018339.	5.9	14
8	Changes in subunit composition of NMDA receptors in animal models of schizophrenia by repeated administration of methamphetamine. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020, 103, 109984.	4.8	12
9	Investigation of the impact of preconditioning with lipopolysaccharide on inflammation-induced gene expression in the brain and depression-like behavior in male mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020, 103, 109978.	4.8	5
10	Effect of interaction between a specific subtype of child abuse and the FKBP5 rs1360780 SNP on DNA methylation among patients with bipolar disorder. <i>Journal of Affective Disorders</i> , 2020, 272, 417-422.	4.1	14
11	Relationship between the subtypes of child abuse and affective temperaments: Comparison of depression and bipolar disorder patients and healthy controls using the reclassified Child Abuse and Trauma Scale. <i>Journal of Affective Disorders</i> , 2019, 257, 396-403.	4.1	4
12	The effects of eicosapentaenoic acid dietary supplementation on behavioral parameters and expression of hippocampal brain-derived neurotrophic factor in an animal model of post-traumatic stress disorder. <i>European Journal of Pharmacology</i> , 2019, 865, 172751.	3.5	4
13	The reduced bactericidal activity of neutrophils as an incisive indicator of water-immersion restraint stress and impaired exercise performance in mice. <i>Scientific Reports</i> , 2019, 9, 4562.	3.3	8
14	Differential effects of voluntary wheel running and toy rotation on the mRNA expression of neurotrophic factors and FKBP5 in a post-traumatic stress disorder rat model with the shuttle-box task. <i>Biochemical and Biophysical Research Communications</i> , 2018, 501, 307-312.	2.1	14
15	Mediators of the effects of rice intake on health in individuals consuming a traditional Japanese diet centered on rice. <i>PLoS ONE</i> , 2017, 12, e0185816.	2.5	12