

Weijuan Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8853675/publications.pdf>

Version: 2024-02-01

55
papers

2,136
citations

257450

24
h-index

254184

43
g-index

56
all docs

56
docs citations

56
times ranked

3353
citing authors

#	ARTICLE	IF	CITATIONS
1	Tumour-associated macrophages-derived CXCL8 determines immune evasion through autonomous PD-L1 expression in gastric cancer. <i>Gut</i> , 2019, 68, 1764-1773.	12.1	219
2	Blockade of Notch1 Signaling Alleviates Murine Lupus via Blunting Macrophage Activation and M2b Polarization. <i>Journal of Immunology</i> , 2010, 184, 6465-6478.	0.8	157
3	Macrophage Differentiation and Polarization via Phosphatidylinositol 3-Kinase/Akt-ERK Signaling Pathway Conferred by Serum Amyloid P Component. <i>Journal of Immunology</i> , 2011, 187, 1764-1777.	0.8	134
4	AIM2 Facilitates the Apoptotic DNA-induced Systemic Lupus Erythematosus via Arbitrating Macrophage Functional Maturation. <i>Journal of Clinical Immunology</i> , 2013, 33, 925-937.	3.8	123
5	Tumor-associated Macrophage-derived Interleukin-23 Interlinks Kidney Cancer Glutamine Addiction with Immune Evasion. <i>European Urology</i> , 2019, 75, 752-763.	1.9	123
6	Prognostic Value of Diametrically Polarized Tumor-Associated Macrophages in Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2014, 21, 3142-3150.	1.5	98
7	EZH2-mediated loss of miR-622 determines CXCR4 activation in hepatocellular carcinoma. <i>Nature Communications</i> , 2015, 6, 8494.	12.8	95
8	Poor Clinical Outcomes and Immuno-evasive Contexture in Intratumoral IL-10-Producing Macrophages Enriched Gastric Cancer Patients. <i>Annals of Surgery</i> , 2022, 275, e626-e635.	4.2	95
9	Intratumoral TIGIT ⁺ CD8 ⁺ T-cell infiltration determines poor prognosis and immune evasion in patients with muscle-invasive bladder cancer. , 2020, 8, e000978.		81
10	Tumor-infiltrating CD39 ⁺ CD8 ⁺ T cells determine poor prognosis and immune evasion in clear cell renal cell carcinoma patients. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1565-1576.	4.2	72
11	Blockade of DC-SIGN ⁺ Tumor-Associated Macrophages Reactivates Antitumor Immunity and Improves Immunotherapy in Muscle-Invasive Bladder Cancer. <i>Cancer Research</i> , 2020, 80, 1707-1719.	0.9	61
12	DNA-dependent Activator of Interferon-regulatory Factors (DAI) Promotes Lupus Nephritis by Activating the Calcium Pathway. <i>Journal of Biological Chemistry</i> , 2013, 288, 13534-13550.	3.4	51
13	High expression of Solute Carrier Family 1, member 5 (SLC1A5) is associated with poor prognosis in clear-cell renal cell carcinoma. <i>Scientific Reports</i> , 2015, 5, 16954.	3.3	43
14	Tumor infiltrating mast cells determine oncogenic HIF-2 α -conferred immune evasion in clear cell renal cell carcinoma. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 731-741.	4.2	39
15	CD19 ⁺ tumor-infiltrating B-cells prime CD4 ⁺ T-cell immunity and predict platinum-based chemotherapy efficacy in muscle-invasive bladder cancer. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 45-56.	4.2	39
16	Tumor-infiltrating $\gamma\delta$ T cells predict prognosis and adjuvant chemotherapeutic benefit in patients with gastric cancer. <i>Oncotimmunology</i> , 2017, 6, e1353858.	4.6	38
17	Tumor-associated macrophages expressing galectin-9 identify immunoevasive subtype muscle-invasive bladder cancer with poor prognosis but favorable adjuvant chemotherapeutic response. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 2067-2080.	4.2	34
18	Identification and validation of an immunogenic subtype of gastric cancer with abundant intratumoural CD103 ⁺ CD8 ⁺ T cells conferring favourable prognosis. <i>British Journal of Cancer</i> , 2020, 122, 1525-1534.	6.4	34

#	ARTICLE	IF	CITATIONS
19	Galectin-9 predicts postoperative recurrence and survival of patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2015, 36, 5791-5799.	1.8	33
20	CD47 expression in gastric cancer clinical correlates and association with macrophage infiltration. <i>Cancer Immunology, Immunotherapy</i> , 2021, 70, 1831-1840.	4.2	32
21	Amelioration of Lupus Nephritis by Serum Amyloid P Component Gene Therapy with Distinct Mechanisms Varied from Different Stage of the Disease. <i>PLoS ONE</i> , 2011, 6, e22659.	2.5	31
22	P2X7 receptor predicts postoperative cancer-specific survival of patients with clear-cell renal cell carcinoma. <i>Cancer Science</i> , 2015, 106, 1224-1231.	3.9	30
23	Prognostic value of interleukin-6 and interleukin-6 receptor in organ-confined clear-cell renal cell carcinoma: a 5-year conditional cancer-specific survival analysis. <i>British Journal of Cancer</i> , 2015, 113, 1581-1589.	6.4	28
24	Poor clinical outcomes of intratumoral dendritic cell-specific intercellular adhesion molecule 3-grabbing non-integrin-positive macrophages associated with immune evasion in gastric cancer. <i>European Journal of Cancer</i> , 2020, 128, 27-37.	2.8	28
25	Increased expression of colony stimulating factor-1 is a predictor of poor prognosis in patients with clear-cell renal cell carcinoma. <i>BMC Cancer</i> , 2015, 15, 67.	2.6	27
26	Loss of N-Acetylgalactosaminyltransferase-4 Orchestrates Oncogenic MicroRNA-9 in Hepatocellular Carcinoma. <i>Journal of Biological Chemistry</i> , 2017, 292, 3186-3200.	3.4	27
27	High expression of interleukin-11 is an independent indicator of poor prognosis in clear-cell renal cell carcinoma. <i>Cancer Science</i> , 2015, 106, 592-597.	3.9	23
28	Elevated expression of IFN-inducible CXCR3 ligands predicts poor prognosis in patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , 2016, 7, 13976-13983.	1.8	23
29	Association of O ⁶ -Methylguanine-DNA Methyltransferase Protein Expression With Postoperative Prognosis and Adjuvant Chemotherapeutic Benefits Among Patients With Stage II or III Gastric Cancer. <i>JAMA Surgery</i> , 2017, 152, e173120.	4.3	22
30	Intratumoral IL22-producing cells define immunoevasive subtype muscle-invasive bladder cancer with poor prognosis and superior nivolumab responses. <i>International Journal of Cancer</i> , 2020, 146, 542-552.	5.1	22
31	Tumor Suppressive Function of p21-activated Kinase 6 in Hepatocellular Carcinoma. <i>Journal of Biological Chemistry</i> , 2015, 290, 28489-28501.	3.4	20
32	Latency-associated Peptide Identifies Immunoevasive Subtype Gastric Cancer With Poor Prognosis and Inferior Chemotherapeutic Responsiveness. <i>Annals of Surgery</i> , 2022, 275, e163-e173.	4.2	17
33	Interleukin-11 receptor predicts post-operative clinical outcome in patients with early-stage clear-cell renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 202-209.	1.3	16
34	Lauren classification identifies distinct prognostic value and functional status of intratumoral CD8+ T cells in gastric cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1327-1336.	4.2	16
35	CCR5 blockade inflames antitumor immunity in BAP1-mutant clear cell renal cell carcinoma. , 2020, 8, e000228.		15
36	Intratumoral CD103 ⁺ CD4 ⁺ T cell infiltration defines immunoevasive contexture and poor clinical outcomes in gastric cancer patients. <i>Oncolmmunology</i> , 2020, 9, 1844402.	4.6	14

#	ARTICLE	IF	CITATIONS
37	Immune inactivation by APOBEC3B enrichment predicts response to chemotherapy and survival in gastric cancer. <i>Oncolmmunology</i> , 2021, 10, 1975386.	4.6	14
38	Snail predicts recurrence and survival of patients with localized clear cell renal cell carcinoma after surgical resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 69.e1-69.e10.	1.6	13
39	IL-33 is associated with unfavorable postoperative survival of patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2016, 37, 11127-11134.	1.8	13
40	GALNT4 Predicts Clinical Outcome in Patients with Clear Cell Renal Cell Carcinoma. <i>Journal of Urology</i> , 2014, 192, 1534-1541.	0.4	12
41	Prognostic significance of β 1,6-N-acetylglucosaminyltransferase V expression in patients with hepatocellular carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 844-853.	1.3	12
42	High Expression of Colony-Stimulating Factor 1 Receptor Associates with Unfavorable Cancer-Specific Survival of Patients with Clear Cell Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2016, 23, 1044-1052.	1.5	11
43	PAK1 expression determines poor prognosis and immune evasion in metastatic renal cell carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020, 38, 293-304.	1.6	10
44	CXC chemokine receptor 1 predicts postoperative prognosis and chemotherapeutic benefits for TNM II and III resectable gastric cancer patients. <i>Oncotarget</i> , 2017, 8, 20328-20339.	1.8	10
45	β 1,6-N-acetylglucosaminyltransferase V predicts recurrence and survival of patients with clear-cell renal cell carcinoma after surgical resection. <i>World Journal of Urology</i> , 2015, 33, 1791-1799.	2.2	9
46	Decreased expression of Siglec-8 associates with poor prognosis in patients with gastric cancer after surgical resection. <i>Tumor Biology</i> , 2016, 37, 10883-10891.	1.8	9
47	Intratumoral Foxp3 ⁺ ROR γ ⁺ T cell infiltration determines poor prognosis and immunoevasive contexture in gastric cancer patients. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 1-11.	4.2	9
48	High expression of C-C chemokine receptor 2 associates with poor overall survival in gastric cancer patients after surgical resection. <i>Oncotarget</i> , 2016, 7, 23909-23918.	1.8	9
49	p21-Activated kinase 4 predicts early recurrence and poor survival in patients with nonmetastatic clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 205.e13-205.e21.	1.6	8
50	Intratumoral interleukin-9 delineates a distinct immunogenic class of gastric cancer patients with better prognosis and adjuvant chemotherapeutic response. <i>Oncolmmunology</i> , 2020, 9, 1856468.	4.6	8
51	Positive intratumoral chemokine (C-C motif) receptor 8 expression predicts high recurrence risk of post-operation clear-cell renal cell carcinoma patients. <i>Oncotarget</i> , 2016, 7, 8413-8421.	1.8	8
52	Identification and validation of an excellent prognosis subtype of muscle-invasive bladder cancer patients with intratumoral CXCR5 ⁺ CD8 ⁺ T cell abundance. <i>Oncolmmunology</i> , 2020, 9, 1810489.	4.6	7
53	Galectin-8 predicts postoperative recurrence of patients with localized T1 clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 112.e1-112.e8.	1.6	5
54	Beta-1,4-galactosyltransferase II predicts poor prognosis of patients with non-metastatic clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2017, 39, 101042831769141.	1.8	5

#	ARTICLE	IF	CITATIONS
55	C-reactive protein functions as a negative regulator of macrophage activation induced by apoptotic DNA. <i>Protein and Cell</i> , 2011, 2, 672-679.	11.0	2