

Weijuan Zhang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8853675/publications.pdf>

Version: 2024-02-01

55
papers

2,136
citations

257450

24
h-index

254184

43
g-index

56
all docs

56
docs citations

56
times ranked

3353
citing authors

#	ARTICLE	IF	CITATIONS
1	Intratumoral Foxp3+ROR γ t+ T cell infiltration determines poor prognosis and immunoevasive contexture in gastric cancer patients. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 1-11.	4.2	9
2	Latency-associated Peptide Identifies Immunoevasive Subtype Gastric Cancer With Poor Prognosis and Inferior Chemotherapeutic Responsiveness. <i>Annals of Surgery</i> , 2022, 275, e163-e173.	4.2	17
3	Poor Clinical Outcomes and Immunoevasive Contexture in Intratumoral IL-10-Producing Macrophages Enriched Gastric Cancer Patients. <i>Annals of Surgery</i> , 2022, 275, e626-e635.	4.2	95
4	CD47 expression in gastric cancer clinical correlates and association with macrophage infiltration. <i>Cancer Immunology, Immunotherapy</i> , 2021, 70, 1831-1840.	4.2	32
5	Immune inactivation by APOBEC3B enrichment predicts response to chemotherapy and survival in gastric cancer. <i>Oncolmmunology</i> , 2021, 10, 1975386.	4.6	14
6	Intratumoral IL22 α -producing cells define immunoevasive subtype muscle α -invasive bladder cancer with poor prognosis and superior nivolumab responses. <i>International Journal of Cancer</i> , 2020, 146, 542-552.	5.1	22
7	PAK1 expression determines poor prognosis and immune evasion in metastatic renal cell carcinoma patients. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2020, 38, 293-304.	1.6	10
8	Identification and validation of an excellent prognosis subtype of muscle-invasive bladder cancer patients with intratumoral CXCR5 ⁺ CD8 ⁺ T cell abundance. <i>Oncolmmunology</i> , 2020, 9, 1810489.	4.6	7
9	Intratumoral TIGIT ⁺ CD8 ⁺ T-cell infiltration determines poor prognosis and immune evasion in patients with muscle-invasive bladder cancer. , 2020, 8, e000978.		81
10	Intratumoral interleukin-9 delineates a distinct immunogenic class of gastric cancer patients with better prognosis and adjuvant chemotherapeutic response. <i>Oncolmmunology</i> , 2020, 9, 1856468.	4.6	8
11	Intratumoral CD103 ⁺ CD4 ⁺ T cell infiltration defines immunoevasive contexture and poor clinical outcomes in gastric cancer patients. <i>Oncolmmunology</i> , 2020, 9, 1844402.	4.6	14
12	CCR5 blockade inflames antitumor immunity in BAP1-mutant clear cell renal cell carcinoma. , 2020, 8, e000228.		15
13	Lauren classification identifies distinct prognostic value and functional status of intratumoral CD8+ T cells in gastric cancer. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1327-1336.	4.2	16
14	Identification and validation of an immunogenic subtype of gastric cancer with abundant intratumoural CD103+CD8+ T cells conferring favourable prognosis. <i>British Journal of Cancer</i> , 2020, 122, 1525-1534.	6.4	34
15	Poor clinical outcomes of intratumoral dendritic cell α -specific intercellular adhesion molecule 3 α - α grabbing non-integrin α -positive macrophages associated with immune evasion in gastric cancer. <i>European Journal of Cancer</i> , 2020, 128, 27-37.	2.8	28
16	Blockade of DC-SIGN+ Tumor-Associated Macrophages Reactivates Antitumor Immunity and Improves Immunotherapy in Muscle-Invasive Bladder Cancer. <i>Cancer Research</i> , 2020, 80, 1707-1719.	0.9	61
17	Tumor-infiltrating CD39+CD8+ T cells determine poor prognosis and immune evasion in clear cell renal cell carcinoma patients. <i>Cancer Immunology, Immunotherapy</i> , 2020, 69, 1565-1576.	4.2	72
18	Tumor-associated macrophages expressing galectin-9 identify immunoevasive subtype muscle-invasive bladder cancer with poor prognosis but favorable adjuvant chemotherapeutic response. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 2067-2080.	4.2	34

#	ARTICLE	IF	CITATIONS
19	Tumour-associated macrophages-derived CXCL8 determines immune evasion through autonomous PD-L1 expression in gastric cancer. <i>Gut</i> , 2019, 68, 1764-1773.	12.1	219
20	Tumor infiltrating mast cells determine oncogenic HIF-2 α -conferred immune evasion in clear cell renal cell carcinoma. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 731-741.	4.2	39
21	CD19+ tumor-infiltrating B-cells prime CD4+ T-cell immunity and predict platinum-based chemotherapy efficacy in muscle-invasive bladder cancer. <i>Cancer Immunology, Immunotherapy</i> , 2019, 68, 45-56.	4.2	39
22	Tumor-associated Macrophage-derived Interleukin-23 Interlinks Kidney Cancer Glutamine Addiction with Immune Evasion. <i>European Urology</i> , 2019, 75, 752-763.	1.9	123
23	Loss of N-Acetylgalactosaminyltransferase-4 Orchestrates Oncogenic MicroRNA-9 in Hepatocellular Carcinoma. <i>Journal of Biological Chemistry</i> , 2017, 292, 3186-3200.	3.4	27
24	Beta-1,4-galactosyltransferase II predicts poor prognosis of patients with non-metastatic clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2017, 39, 101042831769141.	1.8	5
25	Association of O ⁶ -Methylguanine-DNA Methyltransferase Protein Expression With Postoperative Prognosis and Adjuvant Chemotherapeutic Benefits Among Patients With Stage II or III Gastric Cancer. <i>JAMA Surgery</i> , 2017, 152, e173120.	4.3	22
26	Tumor-infiltrating β 1T cells predict prognosis and adjuvant chemotherapeutic benefit in patients with gastric cancer. <i>Oncotarget</i> , 2017, 6, e1353858.	4.6	38
27	CXC chemokine receptor 1 predicts postoperative prognosis and chemotherapeutic benefits for TNM II and III resectable gastric cancer patients. <i>Oncotarget</i> , 2017, 8, 20328-20339.	1.8	10
28	Decreased expression of Siglec-8 associates with poor prognosis in patients with gastric cancer after surgical resection. <i>Tumor Biology</i> , 2016, 37, 10883-10891.	1.8	9
29	IL-33 is associated with unfavorable postoperative survival of patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2016, 37, 11127-11134.	1.8	13
30	High Expression of Colony-Stimulating Factor 1 Receptor Associates with Unfavorable Cancer-Specific Survival of Patients with Clear Cell Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2016, 23, 1044-1052.	1.5	11
31	Positive intratumoral chemokine (C-C motif) receptor 8 expression predicts high recurrence risk of post-operation clear-cell renal cell carcinoma patients. <i>Oncotarget</i> , 2016, 7, 8413-8421.	1.8	8
32	Elevated expression of IFN-inducible CXCR3 ligands predicts poor prognosis in patients with non-metastatic clear-cell renal cell carcinoma. <i>Oncotarget</i> , 2016, 7, 13976-13983.	1.8	23
33	High expression of C-C chemokine receptor 2 associates with poor overall survival in gastric cancer patients after surgical resection. <i>Oncotarget</i> , 2016, 7, 23909-23918.	1.8	9
34	High expression of interleukin-11 is an independent indicator of poor prognosis in clear-cell renal cell carcinoma. <i>Cancer Science</i> , 2015, 106, 592-597.	3.9	23
35	P2X7 receptor predicts postoperative cancer-specific survival of patients with clear-cell renal cell carcinoma. <i>Cancer Science</i> , 2015, 106, 1224-1231.	3.9	30
36	High expression of Solute Carrier Family 1, member 5 (SLC1A5) is associated with poor prognosis in clear-cell renal cell carcinoma. <i>Scientific Reports</i> , 2015, 5, 16954.	3.3	43

#	ARTICLE	IF	CITATIONS
37	Prognostic significance of β 1,6-N-acetylglucosaminyltransferase V expression in patients with hepatocellular carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 844-853.	1.3	12
38	Prognostic value of interleukin-6 and interleukin-6 receptor in organ-confined clear-cell renal cell carcinoma: a 5-year conditional cancer-specific survival analysis. <i>British Journal of Cancer</i> , 2015, 113, 1581-1589.	6.4	28
39	β 1,6-N-acetylglucosaminyltransferase V predicts recurrence and survival of patients with clear-cell renal cell carcinoma after surgical resection. <i>World Journal of Urology</i> , 2015, 33, 1791-1799.	2.2	9
40	p21-Activated kinase 4 predicts early recurrence and poor survival in patients with nonmetastatic clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 205.e13-205.e21.	1.6	8
41	Interleukin-11 receptor predicts post-operative clinical outcome in patients with early-stage clear-cell renal cell carcinoma. <i>Japanese Journal of Clinical Oncology</i> , 2015, 45, 202-209.	1.3	16
42	Galectin-8 predicts postoperative recurrence of patients with localized T1 clear cell renal cell carcinoma. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 112.e1-112.e8.	1.6	5
43	Galectin-9 predicts postoperative recurrence and survival of patients with clear-cell renal cell carcinoma. <i>Tumor Biology</i> , 2015, 36, 5791-5799.	1.8	33
44	Increased expression of colony stimulating factor-1 is a predictor of poor prognosis in patients with clear-cell renal cell carcinoma. <i>BMC Cancer</i> , 2015, 15, 67.	2.6	27
45	Snail predicts recurrence and survival of patients with localized clear cell renal cell carcinoma after surgical resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2015, 33, 69.e1-69.e10.	1.6	13
46	EZH2-mediated loss of miR-622 determines CXCR4 activation in hepatocellular carcinoma. <i>Nature Communications</i> , 2015, 6, 8494.	12.8	95
47	Tumor Suppressive Function of p21-activated Kinase 6 in Hepatocellular Carcinoma. <i>Journal of Biological Chemistry</i> , 2015, 290, 28489-28501.	3.4	20
48	GALNT4 Predicts Clinical Outcome in Patients with Clear Cell Renal Cell Carcinoma. <i>Journal of Urology</i> , 2014, 192, 1534-1541.	0.4	12
49	Prognostic Value of Diametrically Polarized Tumor-Associated Macrophages in Renal Cell Carcinoma. <i>Annals of Surgical Oncology</i> , 2014, 21, 3142-3150.	1.5	98
50	AIM2 Facilitates the Apoptotic DNA-induced Systemic Lupus Erythematosus via Arbitrating Macrophage Functional Maturation. <i>Journal of Clinical Immunology</i> , 2013, 33, 925-937.	3.8	123
51	DNA-dependent Activator of Interferon-regulatory Factors (DAI) Promotes Lupus Nephritis by Activating the Calcium Pathway. <i>Journal of Biological Chemistry</i> , 2013, 288, 13534-13550.	3.4	51
52	Macrophage Differentiation and Polarization via Phosphatidylinositol 3-Kinase/Akt \rightarrow ERK Signaling Pathway Conferred by Serum Amyloid P Component. <i>Journal of Immunology</i> , 2011, 187, 1764-1777.	0.8	134
53	Amelioration of Lupus Nephritis by Serum Amyloid P Component Gene Therapy with Distinct Mechanisms Varied from Different Stage of the Disease. <i>PLoS ONE</i> , 2011, 6, e22659.	2.5	31
54	C-reactive protein functions as a negative regulator of macrophage activation induced by apoptotic DNA. <i>Protein and Cell</i> , 2011, 2, 672-679.	11.0	2

#	ARTICLE	IF	CITATIONS
55	Blockade of Notch1 Signaling Alleviates Murine Lupus via Blunting Macrophage Activation and M2b Polarization. <i>Journal of Immunology</i> , 2010, 184, 6465-6478.	0.8	157