Alessandro Giuffrida

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8855048/publications.pdf

Version: 2024-02-01

40 papers

1,000 citations

18 h-index 433756 31 g-index

40 all docs

40 docs citations

times ranked

40

1103 citing authors

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Application of charged single isomer derivatives of cyclodextrins in capillary electrophoresis for chiral analysis. Journal of Chromatography A, 2010, 1217, 953-967. | 1.8 | 88 |
| 2 | Determination of formal redox potentials in aqueous solution of copper(II) complexes with ligands having nitrogen and oxygen donor atoms and comparison with their EPR and UV–Vis spectral features. Journal of Inorganic Biochemistry, 2013, 128, 137-145. | 1.5 | 75 |
| 3 | Modified cyclodextrins for fast and sensitive chiralâ€capillary electrophoresisâ€mass spectrometry. Electrophoresis, 2009, 30, 1734-1742. | 1.3 | 69 |
| 4 | Recent advances in chiral separation of amino acids using capillary electromigration techniques. Journal of Chromatography A, 2014, 1363, 41-50. | 1.8 | 61 |
| 5 | A re-investigation of copper coordination in the octa-repeats region of the prion protein. Dalton Transactions, 2005, , 150-158. | 1.6 | 55 |
| 6 | 3-Amino derivative of \hat{l}^2 -cyclodextrin: thermodynamics of copper(II) complexes and exploitation of its enantioselectivity in the separation of amino acid racemates by ligand exchange capillary electrophoresis. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2004, 800, 127-133. | 1.2 | 43 |
| 7 | Ligand exchange chiral separations by cyclodextrin derivatives in capillary electrophoresis. Analyst, The, 2003, 128, 134-136. | 1.7 | 37 |
| 8 | The 6-derivative of \hat{l}^2 -cyclodextrin with succinic acid: a new chiral selector for CD-EKC. Journal of Pharmaceutical and Biomedical Analysis, 2005, 37, 1009-1014. | 1.4 | 32 |
| 9 | Chiral analysis of amino acids from conventional and transgenic yeastsa [~] †. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2008, 875, 243-247. | 1.2 | 32 |
| 10 | High selectivity in new chiral separations of dansyl amino acids by cyclodextrin derivatives in electrokinetic chromatography. Journal of Chromatography A, 2007, 1155, 172-179. | 1.8 | 30 |
| 11 | Chiral capillary zone electrophoresis in enantioseparation and analysis of cinacalcet impurities: Use of Quality by Design principles in method development. Journal of Chromatography A, 2018, 1568, 205-213. | 1.8 | 30 |
| 12 | Coordination properties of 3-functionalized \hat{l}^2 -cyclodextrins. Thermodynamic stereoselectivity of copper(ii) complexes of the A,B-diamino derivative and its exploitation in LECE. Dalton Transactions, 2005, , 2731. | 1.6 | 28 |
| 13 | Ligand exchange capillary electrophoresis by cyclodextrin derivatives, a powerful tool for enantiomeric separations. Electrophoresis, 2006, 27, 1471-1480. | 1.3 | 28 |
| 14 | Ac‣PFFDâ€Th: A Trehaloseâ€Conjugated Peptidomimetic as a Strong Suppressor of Amyloidâ€Î² Oligomer Formation and Cytotoxicity. ChemBioChem, 2016, 17, 1541-1549. | 1.3 | 28 |
| 15 | Simple and mixed complexes of copper(II) with 8-hydroxyquinoline derivatives and amino acids: Characterization in solution and potential biological implications. Journal of Inorganic Biochemistry, 2018, 180, 89-100. | 1.5 | 26 |
| 16 | Chiral separation of amino acids derivatised with fluorescein isothiocyanate by single isomer derivatives 3-monodeoxy-3-monoamino-l̂²- and l̂³-cyclodextrins: the effect of the cavity size. Journal of Chromatography A, 2012, 1269, 360-365. | 1.8 | 23 |
| 17 | Analytical quality by design in the development of a cyclodextrinâ€modified capillary electrophoresis method for the assay of metformin and its related substances. Electrophoresis, 2014, 35, 2538-2545. | 1.3 | 22 |
| 18 | Hemispherodextrins, a new class of cyclodextrin derivatives, in capillary electrophoresis. Journal of Chromatography A, 2001, 916, 61-64. | 1.8 | 18 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Mass spectrometry detection as an innovative and advantageous tool in ligand exchange capillary electrophoresis. Electrophoresis, 2011, 32, 1176-1181. | 1.3 | 18 |
| 20 | Anion Encapsulation Drives the Formation of Dimeric Gd ^{III} [15-metallacrown-5] ³⁺ Complexes in Aqueous Solution. Inorganic Chemistry, 2017, 56, 4771-4774. | 1.9 | 18 |
| 21 | Synthesis and characterisation of the 3-amino-derivative of \hat{I}^3 -cyclodextrin, showing receptor ability and metal ion coordination properties. Tetrahedron Letters, 2008, 49, 4765-4767. | 0.7 | 17 |
| 22 | The 3-amino-derivative of \hat{I}^3 -cyclodextrin as chiral selector of Dns-amino acids in electrokinetic chromatography. Journal of Chromatography A, 2009, 1216, 3678-3686. | 1.8 | 17 |
| 23 | Could 2,6-bis((E)-2-(furan-2-yl)vinyl)-1-methylpyridinium iodide and analog compounds intercalate DNA? A first principle prediction based on structural and electronic properties. Computational and Theoretical Chemistry, 2012, 985, 8-13. | 1.1 | 17 |
| 24 | Simultaneous separation of different enantiomeric pairs in capillary electrophoresis by mixing different hemispherodextrins, a very versatile class of receptors. Journal of Chromatography A, 2002, 979, 137-145. | 1.8 | 16 |
| 25 | The Contribution of Electrospray Mass Spectrometry to the Study of Metal Complexes: The Case of Copper(II)–Dipeptide Systems. European Journal of Inorganic Chemistry, 2009, 2009, 2612-2620. | 1.0 | 16 |
| 26 | New cyclodextrin derivatives as chiral selectors in capillary electrophoresis. Fresenius' Journal of Analytical Chemistry, 2001, 370, 363-366. | 1.5 | 15 |
| 27 | Diaminotrehalose $\hat{\epsilon}$ capped $\hat{l}^2\hat{a}$ cyclodextrin, a new member of hemispherodextrins: Synthesis, thermodynamic and spectroscopic characterization and its exploitation in chiral electrokinetic chromatography. Journal of Separation Science, 2011, 34, 70-76. | 1.3 | 15 |
| 28 | Neuroprotective effects of the monoamine oxidase inhibitor tranylcypromine and its amide derivatives against $A\hat{l}^2(1\hat{a}\in 42)$ -induced toxicity. European Journal of Pharmacology, 2015, 764, 256-263. | 1.7 | 14 |
| 29 | Chiral separation of terbutaline and non-steroidal anti-inflammatory drugs by using a new lysineâ€ ⁴ bridged hemispherodextrin in capillary electrophoresis. Journal of Pharmaceutical and Biomedical Analysis, 2017, 145, 734-741. | 1.4 | 14 |
| 30 | Synthesis and NMR characterization of βâ€alanineâ€bridged hemispherodextrin, a very efficient chiral selector in EKC. Electrophoresis, 2007, 28, 2580-2588. | 1.3 | 13 |
| 31 | The pivotal role of copper(II) in the enantiorecognition of tryptophan and histidine by gold nanoparticles. Analytical and Bioanalytical Chemistry, 2014, 406, 481-491. | 1.9 | 12 |
| 32 | Semax, an ACTH4-10 peptide analog with high affinity for copper(II) ion and protective ability against metal induced cell toxicity. Journal of Inorganic Biochemistry, 2015, 142, 39-46. | 1.5 | 12 |
| 33 | Coordination Properties of 3-Functionalised \hat{l}^2 -Cyclodextrins: Thermodynamic Stereoselectivity of Copper(II) Complexes of the 3-Histamine Derivative and Its Exploitation in Ligand-Exchange Capillary Electrophoresis. European Journal of Inorganic Chemistry, 2014, 2014, 377-383. | 1.0 | 11 |
| 34 | Separation and quantitation of metal ions by 4-(2-pyridylazo)resorcinol complexation in capillary electrophoresis–electrospray ionisation mass spectrometry. Journal of Chromatography A, 2008, 1179, 17-23. | 1.8 | 10 |
| 35 | Synthesis and characterization of new tyrosine capped anisotropic silver nanoparticles and their exploitation for the selective determination of iodide ions. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 2017, 529, 128-136. | 2.3 | 9 |
| 36 | Optimisation methodology in the chiral and achiral separation in electrokinetic chromatography in the case of a multicomponent sample of dansyl amino acids. Journal of Pharmaceutical and Biomedical Analysis, 2013, 85, 55-60. | 1.4 | 8 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | Exploring the intermolecular interactions acting in solvent-modified MEKC by Molecular Dynamics and NMR: The effect of n-butanol on the separation of diclofenac and its impurities. Journal of Pharmaceutical and Biomedical Analysis, 2018, 149, 249-257. | 1.4 | 8 |
| 38 | MV-Flow and LumiFlow: new Doppler tools for evaluating the microvasculature of the fetal head. Radiologia Brasileira, 2021, 54, 348-349. | 0.3 | 7 |
| 39 | Influence of the N-terminus acetylation of Semax, a synthetic analog of ACTH(4-10), on copper(II) and zinc(II) coordination and biological properties. Journal of Inorganic Biochemistry, 2016, 164, 59-69. | 1.5 | 5 |
| 40 | Preclinical evidence of enhanced analgesic activity of duloxetine complexed with succinyl- \hat{l}^2 -cyclodextrin: A comparative study with cyclodextrin complexes. International Journal of Pharmaceutics, 2019, 566, 391-399. | 2.6 | 3 |