

Alessandro Giuffrida

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8855048/publications.pdf>

Version: 2024-02-01

40
papers

1,000
citations

430442

18
h-index

433756

31
g-index

40
all docs

40
docs citations

40
times ranked

1103
citing authors

#	ARTICLE	IF	CITATIONS
1	MV-Flow and LumiFlow: new Doppler tools for evaluating the microvasculature of the fetal head. <i>Radiologia Brasileira</i> , 2021, 54, 348-349.	0.3	7
2	Preclinical evidence of enhanced analgesic activity of duloxetine complexed with succinyl- β -cyclodextrin: A comparative study with cyclodextrin complexes. <i>International Journal of Pharmaceutics</i> , 2019, 566, 391-399.	2.6	3
3	Exploring the intermolecular interactions acting in solvent-modified MEKC by Molecular Dynamics and NMR: The effect of n-butanol on the separation of diclofenac and its impurities. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018, 149, 249-257.	1.4	8
4	Simple and mixed complexes of copper(II) with 8-hydroxyquinoline derivatives and amino acids: Characterization in solution and potential biological implications. <i>Journal of Inorganic Biochemistry</i> , 2018, 180, 89-100.	1.5	26
5	Chiral capillary zone electrophoresis in enantioseparation and analysis of cinacalcet impurities: Use of Quality by Design principles in method development. <i>Journal of Chromatography A</i> , 2018, 1568, 205-213.	1.8	30
6	Anion Encapsulation Drives the Formation of Dimeric Gd ^{III} [15-metallacrown-5] ³⁺ Complexes in Aqueous Solution. <i>Inorganic Chemistry</i> , 2017, 56, 4771-4774.	1.9	18
7	Synthesis and characterization of new tyrosine capped anisotropic silver nanoparticles and their exploitation for the selective determination of iodide ions. <i>Colloids and Surfaces A: Physicochemical and Engineering Aspects</i> , 2017, 529, 128-136.	2.3	9
8	Chiral separation of terbutaline and non-steroidal anti-inflammatory drugs by using a new lysine-bridged hemispherodextrin in capillary electrophoresis. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2017, 145, 734-741.	1.4	14
9	Acetyl-PFFD β Th: A Trehalose-Conjugated Peptidomimetic as a Strong Suppressor of Amyloid β Oligomer Formation and Cytotoxicity. <i>ChemBioChem</i> , 2016, 17, 1541-1549.	1.3	28
10	Influence of the N-terminus acetylation of Semax, a synthetic analog of ACTH(4-10), on copper(II) and zinc(II) coordination and biological properties. <i>Journal of Inorganic Biochemistry</i> , 2016, 164, 59-69.	1.5	5
11	Neuroprotective effects of the monoamine oxidase inhibitor tranylcypromine and its amide derivatives against A β (1-42)-induced toxicity. <i>European Journal of Pharmacology</i> , 2015, 764, 256-263.	1.7	14
12	Semax, an ACTH4-10 peptide analog with high affinity for copper(II) ion and protective ability against metal induced cell toxicity. <i>Journal of Inorganic Biochemistry</i> , 2015, 142, 39-46.	1.5	12
13	Coordination Properties of 3-Functionalised β -Cyclodextrins: Thermodynamic Stereoselectivity of Copper(II) Complexes of the 3-Histamine Derivative and Its Exploitation in Ligand-Exchange Capillary Electrophoresis. <i>European Journal of Inorganic Chemistry</i> , 2014, 2014, 377-383.	1.0	11
14	The pivotal role of copper(II) in the enantio-recognition of tryptophan and histidine by gold nanoparticles. <i>Analytical and Bioanalytical Chemistry</i> , 2014, 406, 481-491.	1.9	12
15	Recent advances in chiral separation of amino acids using capillary electromigration techniques. <i>Journal of Chromatography A</i> , 2014, 1363, 41-50.	1.8	61
16	Analytical quality by design in the development of a cyclodextrin-modified capillary electrophoresis method for the assay of metformin and its related substances. <i>Electrophoresis</i> , 2014, 35, 2538-2545.	1.3	22
17	Optimisation methodology in the chiral and achiral separation in electrokinetic chromatography in the case of a multicomponent sample of dansyl amino acids. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2013, 85, 55-60.	1.4	8
18	Determination of formal redox potentials in aqueous solution of copper(II) complexes with ligands having nitrogen and oxygen donor atoms and comparison with their EPR and UV-Vis spectral features. <i>Journal of Inorganic Biochemistry</i> , 2013, 128, 137-145.	1.5	75

#	ARTICLE	IF	CITATIONS
19	Chiral separation of amino acids derivatised with fluorescein isothiocyanate by single isomer derivatives 3-monodeoxy-3-monoamino- β - and γ -cyclodextrins: the effect of the cavity size. <i>Journal of Chromatography A</i> , 2012, 1269, 360-365.	1.8	23
20	Could 2,6-bis((E)-2-(furan-2-yl)vinyl)-1-methylpyridinium iodide and analog compounds intercalate DNA? A first principle prediction based on structural and electronic properties. <i>Computational and Theoretical Chemistry</i> , 2012, 985, 8-13.	1.1	17
21	Diaminotrehalose- α -capped β -cyclodextrin, a new member of hemispherodextrins: Synthesis, thermodynamic and spectroscopic characterization and its exploitation in chiral electrokinetic chromatography. <i>Journal of Separation Science</i> , 2011, 34, 70-76.	1.3	15
22	Mass spectrometry detection as an innovative and advantageous tool in ligand exchange capillary electrophoresis. <i>Electrophoresis</i> , 2011, 32, 1176-1181.	1.3	18
23	Application of charged single isomer derivatives of cyclodextrins in capillary electrophoresis for chiral analysis. <i>Journal of Chromatography A</i> , 2010, 1217, 953-967.	1.8	88
24	The Contribution of Electrospray Mass Spectrometry to the Study of Metal Complexes: The Case of Copper(II)-Dipeptide Systems. <i>European Journal of Inorganic Chemistry</i> , 2009, 2009, 2612-2620.	1.0	16
25	Modified cyclodextrins for fast and sensitive chiral capillary electrophoresis-mass spectrometry. <i>Electrophoresis</i> , 2009, 30, 1734-1742.	1.3	69
26	The 3-amino-derivative of γ -cyclodextrin as chiral selector of Dns-amino acids in electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2009, 1216, 3678-3686.	1.8	17
27	Synthesis and characterisation of the 3-amino-derivative of β -cyclodextrin, showing receptor ability and metal ion coordination properties. <i>Tetrahedron Letters</i> , 2008, 49, 4765-4767.	0.7	17
28	Chiral analysis of amino acids from conventional and transgenic yeasts. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008, 875, 243-247.	1.2	32
29	Separation and quantitation of metal ions by 4-(2-pyridylazo)resorcinol complexation in capillary electrophoresis-electrospray ionisation mass spectrometry. <i>Journal of Chromatography A</i> , 2008, 1179, 17-23.	1.8	10
30	Synthesis and NMR characterization of β -alanine-bridged hemispherodextrin, a very efficient chiral selector in EKC. <i>Electrophoresis</i> , 2007, 28, 2580-2588.	1.3	13
31	High selectivity in new chiral separations of dansyl amino acids by cyclodextrin derivatives in electrokinetic chromatography. <i>Journal of Chromatography A</i> , 2007, 1155, 172-179.	1.8	30
32	Ligand exchange capillary electrophoresis by cyclodextrin derivatives, a powerful tool for enantiomeric separations. <i>Electrophoresis</i> , 2006, 27, 1471-1480.	1.3	28
33	The 6-derivative of β -cyclodextrin with succinic acid: a new chiral selector for CD-EKC. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2005, 37, 1009-1014.	1.4	32
34	Coordination properties of 3-functionalized β -cyclodextrins. Thermodynamic stereoselectivity of copper(II) complexes of the A,B-diamino derivative and its exploitation in LECE. <i>Dalton Transactions</i> , 2005, , 2731.	1.6	28
35	A re-investigation of copper coordination in the octa-repeats region of the prion protein. <i>Dalton Transactions</i> , 2005, , 150-158.	1.6	55
36	3-Amino derivative of β -cyclodextrin: thermodynamics of copper(II) complexes and exploitation of its enantioselectivity in the separation of amino acid racemates by ligand exchange capillary electrophoresis. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2004, 800, 127-133.	1.2	43

#	ARTICLE	IF	CITATIONS
37	Ligand exchange chiral separations by cyclodextrin derivatives in capillary electrophoresis. <i>Analyst</i> , 2003, 128, 134-136.	1.7	37
38	Simultaneous separation of different enantiomeric pairs in capillary electrophoresis by mixing different hemispherodextrins, a very versatile class of receptors. <i>Journal of Chromatography A</i> , 2002, 979, 137-145.	1.8	16
39	New cyclodextrin derivatives as chiral selectors in capillary electrophoresis. <i>Fresenius' Journal of Analytical Chemistry</i> , 2001, 370, 363-366.	1.5	15
40	Hemispherodextrins, a new class of cyclodextrin derivatives, in capillary electrophoresis. <i>Journal of Chromatography A</i> , 2001, 916, 61-64.	1.8	18