

Jeffrey R Schelling

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8857875/jeffrey-r-schelling-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

75
papers

4,259
citations

32
h-index

65
g-index

77
ext. papers

4,877
ext. citations

7.4
avg, IF

4.74
L-index

#	Paper	IF	Citations
75	Ventricular arrhythmias in mouse models of diabetic kidney disease. <i>Scientific Reports</i> , 2021 , 11, 20570	4.9	0
74	Social Support in Older Adults With CKD: A Report From the CRIC (Chronic Renal Insufficiency Cohort) Study. <i>Kidney Medicine</i> , 2021 , 3, 776-784.e1	2.8	
73	Association of Multiple Plasma Biomarker Concentrations with Progression of Prevalent Diabetic Kidney Disease: Findings from the Chronic Renal Insufficiency Cohort (CRIC) Study. <i>Journal of the American Society of Nephrology: JASN</i> , 2021 , 32, 115-126	12.7	26
72	Urine Biomarkers of Kidney Tubule Health, Injury, and Inflammation are Associated with Progression of CKD in Children. <i>Journal of the American Society of Nephrology: JASN</i> , 2021 , 32, 2664-2677	12.7	3
71	Plasma Biomarkers of Tubular Injury and Inflammation Are Associated with CKD Progression in Children. <i>Journal of the American Society of Nephrology: JASN</i> , 2020 , 31, 1067-1077	12.7	22
70	Fatty acid transport protein-2 regulates glycemic control and diabetic kidney disease progression. <i>JCI Insight</i> , 2020 , 5,	9.9	7
69	Longitudinal Changes in Health-Related Quality of Life in Primary Glomerular Disease: Results From the CureGN Study. <i>Kidney International Reports</i> , 2020 , 5, 1679-1689	4.1	4
68	Health-related quality of life in glomerular disease. <i>Kidney International</i> , 2019 , 95, 1209-1224	9.9	20
67	Erythrocytosis is associated with intradialytic hypotension: a case series. <i>BMC Nephrology</i> , 2019 , 20, 235	2.7	0
66	Incident Type 2 Diabetes Among Individuals With CKD: Findings From the Chronic Renal Insufficiency Cohort (CRIC) Study. <i>American Journal of Kidney Diseases</i> , 2019 , 73, 72-81	7.4	14
65	CureGN Study Rationale, Design, and Methods: Establishing a Large Prospective Observational Study of Glomerular Disease. <i>American Journal of Kidney Diseases</i> , 2019 , 73, 218-229	7.4	39
64	Kidney Proximal Tubule Lipoapoptosis Is Regulated by Fatty Acid Transporter-2 (FATP2). <i>Journal of the American Society of Nephrology: JASN</i> , 2018 , 29, 81-91	12.7	29
63	Clinical Characteristics and Treatment Patterns of Children and Adults With IgA Nephropathy or IgA Vasculitis: Findings From the CureGN Study. <i>Kidney International Reports</i> , 2018 , 3, 1373-1384	4.1	23
62	Urine Kidney Injury Biomarkers and Risks of Cardiovascular Disease Events and All-Cause Death: The CRIC Study. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2017 , 12, 761-771	6.9	34
61	Urine biomarkers of tubular injury do not improve on the clinical model predicting chronic kidney disease progression. <i>Kidney International</i> , 2017 , 91, 196-203	9.9	53
60	GFR Estimation Using β -Trace Protein and β -Microglobulin in CKD. <i>American Journal of Kidney Diseases</i> , 2016 , 67, 40-8	7.4	94
59	Tubular atrophy in the pathogenesis of chronic kidney disease progression. <i>Pediatric Nephrology</i> , 2016 , 31, 693-706	3.2	78

58	Variants in genes belonging to the fibroblast growth factor family are associated with lower extremity amputation in non-Hispanic whites: Findings from the chronic renal insufficiency cohort study. <i>Wound Repair and Regeneration</i> , 2016 , 24, 705-11	3.6	1
57	Serum β Trace Protein and β -Microglobulin as Predictors of ESRD, Mortality, and Cardiovascular Disease in Adults With CKD in the Chronic Renal Insufficiency Cohort (CRIC) Study. <i>American Journal of Kidney Diseases</i> , 2016 , 68, 68-76	7.4	50
56	Lipidomic Signature of Progression of Chronic Kidney Disease in the Chronic Renal Insufficiency Cohort. <i>Kidney International Reports</i> , 2016 , 1, 256-268	4.1	50
55	Genome-Wide Association and Trans-ethnic Meta-Analysis for Advanced Diabetic Kidney Disease: Family Investigation of Nephropathy and Diabetes (FIND). <i>PLoS Genetics</i> , 2015 , 11, e1005352	6	84
54	Na ⁺ /H ⁺ exchanger-1 (NHE1) regulation in kidney proximal tubule. <i>Cellular and Molecular Life Sciences</i> , 2015 , 72, 2061-74	10.3	18
53	Relation of serum lipids and lipoproteins with progression of CKD: The CRIC study. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014 , 9, 1190-8	6.9	90
52	Mesangial cell α 8-integrin regulates glomerular capillary integrity and repair. <i>American Journal of Physiology - Renal Physiology</i> , 2014 , 306, F1400-9	4.3	10
51	Treatment of albuminuria due to diabetic nephropathy: recent trial results. <i>Clinical Investigation</i> , 2014 , 4, 327-341		1
50	Lipotoxic disruption of NHE1 interaction with PI(4,5)P2 expedites proximal tubule apoptosis. <i>Journal of Clinical Investigation</i> , 2014 , 124, 1057-68	15.9	34
49	Association of N-terminal pro-B-type natriuretic peptide with left ventricular structure and function in chronic kidney disease (from the Chronic Renal Insufficiency Cohort [CRIC]). <i>American Journal of Cardiology</i> , 2013 , 111, 432-8	3	42
48	Diabetes, lower extremity amputation, loss of protective sensation, and neuronal nitric oxide synthase associated protein in the chronic renal insufficiency cohort study. <i>Wound Repair and Regeneration</i> , 2013 , 21, 17-24	3.6	10
47	Renal function and proteinuria after successful immunosuppressive therapies in patients with FSGS. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2013 , 8, 211-8	6.9	16
46	A genome-wide search for linkage of estimated glomerular filtration rate (eGFR) in the Family Investigation of Nephropathy and Diabetes (FIND). <i>PLoS ONE</i> , 2013 , 8, e81888	3.7	23
45	Mesangial cell integrin α 8 provides glomerular endothelial cell cytoprotection by sequestering TGF- β and regulating PECAM-1. <i>American Journal of Pathology</i> , 2011 , 178, 609-20	5.8	37
44	APOL1 genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy. <i>Journal of the American Society of Nephrology: JASN</i> , 2011 , 22, 2129-37	12.7	540
43	Genomewide linkage scan for diabetic renal failure and albuminuria: the FIND study. <i>American Journal of Nephrology</i> , 2011 , 33, 381-9	4.6	48
42	Use of diffusion tensor MRI to identify early changes in diabetic nephropathy. <i>American Journal of Nephrology</i> , 2011 , 34, 476-82	4.6	82
41	Phosphoinositide binding differentially regulates NHE1 Na ⁺ /H ⁺ exchanger-dependent proximal tubule cell survival. <i>Journal of Biological Chemistry</i> , 2011 , 286, 42435-42445	5.4	24

40	Podocyte injury induces nuclear translocation of WTIP via microtubule-dependent transport. <i>Journal of Biological Chemistry</i> , 2010 , 285, 9995-10004	5.4	21
39	Ouabain stimulates Na-K-ATPase through a sodium/hydrogen exchanger-1 (NHE-1)-dependent mechanism in human kidney proximal tubule cells. <i>American Journal of Physiology - Renal Physiology</i> , 2010 , 299, F77-90	4.3	49
38	Tissue transglutaminase inhibition as treatment for diabetic glomerular scarring: it's good to be glueless. <i>Kidney International</i> , 2009 , 76, 363-5	9.9	9
37	Genome-wide linkage scans for type 2 diabetes mellitus in four ethnically diverse populations-significant evidence for linkage on chromosome 4q in African Americans: the Family Investigation of Nephropathy and Diabetes Research Group. <i>Diabetes/Metabolism Research and Reviews</i> , 2009 , 25, 740-7	7.5	9
36	Transient Elevations of Microalbuminuria May Represent a Marker of Systemic Vascular Injury in Sickle Cell Disease Patients.. <i>Blood</i> , 2009 , 114, 4611-4611	2.2	
35	MYH9 is a major-effect risk gene for focal segmental glomerulosclerosis. <i>Nature Genetics</i> , 2008 , 40, 1175-84	36.3	559
34	MYH9 is associated with nondiabetic end-stage renal disease in African Americans. <i>Nature Genetics</i> , 2008 , 40, 1185-92	36.3	518
33	Risk factors for development and progression of diabetic kidney disease and treatment patterns among diabetic siblings of patients with diabetic kidney disease. <i>American Journal of Kidney Diseases</i> , 2008 , 51, 29-37	7.4	21
32	Genome-wide scan for estimated glomerular filtration rate in multi-ethnic diabetic populations: the Family Investigation of Nephropathy and Diabetes (FIND). <i>Diabetes</i> , 2008 , 57, 235-43	0.9	75
31	Regulation of cell survival by Na ⁺ /H ⁺ exchanger-1. <i>American Journal of Physiology - Renal Physiology</i> , 2008 , 295, F625-32	4.3	71
30	The WT1 Interacting Protein: A molecular messenger between the slit diaphragm and podocyte nucleus. <i>FASEB Journal</i> , 2008 , 22, 1218.9	0.9	
29	NPHS2 variation in sporadic focal segmental glomerulosclerosis. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 2987-95	12.7	49
28	Genome-wide scans for diabetic nephropathy and albuminuria in multiethnic populations: the family investigation of nephropathy and diabetes (FIND). <i>Diabetes</i> , 2007 , 56, 1577-85	0.9	125
27	Beta8 integrin binds Rho GDP dissociation inhibitor-1 and activates Rac1 to inhibit mesangial cell myofibroblast differentiation. <i>Journal of Biological Chemistry</i> , 2006 , 281, 19688-99	5.4	32
26	A dipstick protein and specific gravity algorithm accurately predicts pathological proteinuria. <i>American Journal of Kidney Diseases</i> , 2005 , 45, 833-41	7.4	32
25	Diabetes and the kidney. <i>American Journal of Kidney Diseases</i> , 2005 , 46, 766-73	7.4	30
24	Variants in the WilmsTumor gene are associated with focal segmental glomerulosclerosis in the African American population. <i>Physiological Genomics</i> , 2005 , 21, 212-21	3.6	47
23	The metabolic syndrome as a risk factor for chronic kidney disease: more than a fat chance?. <i>Journal of the American Society of Nephrology: JASN</i> , 2004 , 15, 2773-4	12.7	30

22	The NHE1 Na ⁺ /H ⁺ exchanger recruits ezrin/radixin/moesin proteins to regulate Akt-dependent cell survival. <i>Journal of Biological Chemistry</i> , 2004 , 279, 26280-6	5.4	128
21	A WT1 co-regulator controls podocyte phenotype by shuttling between adhesion structures and nucleus. <i>Journal of Biological Chemistry</i> , 2004 , 279, 14398-408	5.4	67
20	Renal phenotype is exacerbated in Os and lpr double mutant mice. <i>Kidney International</i> , 2004 , 66, 1029-35	3.9	3
19	Kidney disease, genotype and the pathogenesis of vasculopathy. <i>Current Opinion in Nephrology and Hypertension</i> , 2003 , 12, 71-8	3.5	10
18	DNA expression analysis: serial analysis of gene expression, microarrays and kidney disease. <i>Current Opinion in Nephrology and Hypertension</i> , 2003 , 12, 407-14	3.5	12
17	Nucleic acid-based techniques for post-transcriptional regulation of molecular targets. <i>Current Opinion in Nephrology and Hypertension</i> , 2003 , 12, 415-21	3.5	6
16	Renal tubular epithelial cell apoptosis is associated with caspase cleavage of the NHE1 Na ⁺ /H ⁺ exchanger. <i>American Journal of Physiology - Renal Physiology</i> , 2003 , 284, F829-39	4.3	57
15	Generation of kidney transcriptomes using serial analysis of gene expression. <i>Nephron Experimental Nephrology</i> , 2002 , 10, 82-92		10
14	Approaches to understanding susceptibility to nephropathy: from genetics to genomics. <i>Kidney International</i> , 2002 , 61, S61-7	9.9	13
13	Fas activation induces renal tubular epithelial cell beta 8 integrin expression and function in the absence of apoptosis. <i>Journal of Biological Chemistry</i> , 2002 , 277, 47826-33	5.4	31
12	Myofibroblast differentiation: plasma membrane microdomains and cell phenotype. <i>Nephron Experimental Nephrology</i> , 2002 , 10, 313-9		7
11	Apoptosis and JNK activation are differentially regulated by Fas expression level in renal tubular epithelial cells. <i>Kidney International</i> , 2001 , 60, 65-76	9.9	22
10	Activation of EphA receptor tyrosine kinase inhibits the Ras/MAPK pathway. <i>Nature Cell Biology</i> , 2001 , 3, 527-30	23.4	273
9	A family-based strategy to identify genes for diabetic nephropathy. <i>American Journal of Kidney Diseases</i> , 2001 , 37, 638-47	7.4	11
8	Serum C-peptide concentrations poorly phenotype type 2 diabetic end-stage renal disease patients. <i>Kidney International</i> , 2000 , 58, 1742-50	9.9	40
7	Inaccuracy of clinical phenotyping parameters for hypertensive nephrosclerosis. <i>Nephrology Dialysis Transplantation</i> , 2000 , 15, 1801-7	4.3	53
6	Interleukin-1 stimulates Jun N-terminal/stress-activated protein kinase by an arachidonate-dependent mechanism in mesangial cells. <i>Kidney International</i> , 1999 , 55, 1740-9	9.9	20
5	Involvement of Fas-dependent apoptosis in renal tubular epithelial cell deletion in chronic renal failure. <i>Kidney International</i> , 1999 , 56, 1313-6	9.9	47

4	Genetic susceptibility to end-stage renal disease. <i>Current Opinion in Nephrology and Hypertension</i> , 1999 , 8, 465-72	3.5	42
3	Management of tumor lysis syndrome with standard continuous arteriovenous hemodialysis: case report and a review of the literature. <i>Renal Failure</i> , 1998 , 20, 635-44	2.9	13
2	Glucocorticoid uncoupling of angiotensin II-dependent phospholipase C activation in rat vascular smooth muscle cells. <i>Kidney International</i> , 1994 , 46, 675-82	9.9	14
1	Increased osmolal gap in alcoholic ketoacidosis and lactic acidosis. <i>Annals of Internal Medicine</i> , 1990 , 113, 580-2	8	88