

# Faisal A Almobarak

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8858387/publications.pdf>

Version: 2024-02-01

29  
papers

540  
citations

1039880

9  
h-index

642610

23  
g-index

31  
all docs

31  
docs citations

31  
times ranked

592  
citing authors

#	ARTICLE	IF	CITATIONS
1	Enhanced Detection of Open-angle Glaucoma with an Anatomically Accurate Optical Coherence Tomographyâ€‘Derived Neuroretinal Rim Parameter. <i>Ophthalmology</i> , 2013, 120, 535-543.	2.5	323
2	Association of increased levels of plasma tumor necrosis factor alpha with primary open-angle glaucoma. <i>Clinical Ophthalmology</i> , 2018, Volume 12, 701-706.	0.9	34
3	Elevated levels of plasma tumor necrosis factor alpha in patients with pseudoexfoliation glaucoma. <i>Clinical Ophthalmology</i> , 2018, Volume 12, 153-159.	0.9	26
4	Intermediate and Long-term Outcomes of Mitomycin Câ€‘enhanced Trabeculectomy as a First Glaucoma Procedure in Uveitic Glaucoma. <i>Journal of Glaucoma</i> , 2017, 26, 478-485.	0.8	19
5	Polymorphism rs10483727 in the SIX1/SIX6 Gene Locus Is a Risk Factor for Primary Open Angle Glaucoma in a Saudi Cohort. <i>Genetic Testing and Molecular Biomarkers</i> , 2018, 22, 74-78.	0.3	14
6	Polymorphism rs11656696 in <i>GAS7</i> Is Not Associated with Primary Open Angle Glaucoma in a Saudi Cohort. <i>Genetic Testing and Molecular Biomarkers</i> , 2017, 21, 754-758.	0.3	11
7	Association of endothelial nitric oxide synthase (NOS3) gene polymorphisms with primary open-angle glaucoma in a Saudi cohort. <i>PLoS ONE</i> , 2020, 15, e0227417.	1.1	11
8	Analysis of <i>Cyclin-Dependent Kinase Inhibitor-2B</i> rs1063192 Polymorphism in Saudi Patients with Primary Open-Angle Glaucoma. <i>Genetic Testing and Molecular Biomarkers</i> , 2016, 20, 637-641.	0.3	10
9	Elevated Plasma Level of 8-Hydroxy-2â€‘deoxyguanosine Is Associated with Primary Open-Angle Glaucoma. <i>Journal of Ophthalmology</i> , 2020, 2020, 1-8.	0.6	10
10	The influence of mitomycin C concentration on the outcome of trabeculectomy in uveitic glaucoma. <i>International Ophthalmology</i> , 2018, 38, 2371-2379.	0.6	9
11	Incidence and management of haemorrhagic Descemet membrane detachment in canaloplasty and phacocanaloplasty. <i>Acta Ophthalmologica</i> , 2016, 94, e298-304.	0.6	8
12	Association analysis of polymorphisms rs12997 in ACVR1 and rs1043784 in BMP6 genes involved in bone morphogenic protein signaling pathway in primary angle-closure and pseudoexfoliation glaucoma patients of Saudi origin. <i>BMC Medical Genetics</i> , 2020, 21, 145.	2.1	8
13	Outcomes of Trabeculectomy With Mitomycin-C in Uveitis Associated With Vogt-Koyanagi-Harada Disease. <i>Journal of Glaucoma</i> , 2016, 25, 528-532.	0.8	7
14	The Influence of Phacoemulsification on Intraocular Pressure Control and Trabeculectomy Survival in Uveitic Glaucoma. <i>Journal of Glaucoma</i> , 2017, 26, 444-449.	0.8	7
15	Plexin domain containing 2 (PLXDC2) gene polymorphism rs7081455 may not influence POAG risk in a Saudi cohort. <i>BMC Research Notes</i> , 2018, 11, 733.	0.6	7
16	Analysis of Polymorphism rs1900004 in Atonal bHLH Transcription Factor 7 in Saudi Patients with Primary Open Angle Glaucoma. <i>Genetic Testing and Molecular Biomarkers</i> , 2016, 20, 715-718.	0.3	6
17	Polymorphism rs7961953 in TMTC2 gene is not associated with primary open-angle glaucoma in a Saudi cohort. <i>Ophthalmic Genetics</i> , 2019, 40, 74-76.	0.5	6
18	Ultrasound Cyclo Plasty After Failed Glaucoma Surgery: Outcomes and Complications. <i>Ophthalmology and Therapy</i> , 2022, 11, 1601-1610.	1.0	5

#	ARTICLE	IF	CITATIONS
19	Lack of Association between Variant rs7916697 in ATOH7 and Primary Open Angle Glaucoma in a Saudi Cohort. <i>Genetics Research International</i> , 2018, 2018, 1-6.	2.0	4
20	Lack of Association Between Polymorphisms in TXNRD2 and LMX1B and Primary Open-Angle Glaucoma in a Saudi Cohort. <i>Frontiers in Genetics</i> , 2021, 12, 690780.	1.1	3
21	Polymorphism rs3742330 in microRNA Biogenesis Gene DICER1 Is Associated with Pseudoexfoliation Glaucoma in Saudi Cohort. <i>Genes</i> , 2022, 13, 489.	1.0	3
22	Association of Rs12997 Variant in the <i>Acvr1</i> Gene: A Member of Bone Morphogenic Protein Signaling Pathway with Primary Open-Angle Glaucoma in a Saudi Cohort. <i>Journal of Investigative Medicine</i> , 2021, 69, 402-407.	0.7	2
23	Evaluation of ABCA1 and FNDC3B Gene Polymorphisms Associated With Pseudoexfoliation Glaucoma and Primary Angle-Closure Glaucoma in a Saudi Cohort. <i>Frontiers in Genetics</i> , 0, 13, .	1.1	2
24	Aqueous misdirection after Nd:YAG goniotomy in deep sclerectomy treated with Nd:YAG irido-zonulo-hyaloidotomy. <i>European Journal of Ophthalmology</i> , 2021, , 112067212110195.	0.7	1
25	Long-term outcomes of initial trabeculectomy in glaucoma associated with granulomatous and non-granulomatous uveitis. <i>International Ophthalmology</i> , 2021, 41, 3459-3470.	0.6	1
26	Association analysis of variants rs35934224 in TXNRD2 and rs6478746 in LMX1B in primary angle-closure and pseudoexfoliation glaucoma. <i>European Journal of Ophthalmology</i> , 2021, , 112067212110425.	0.7	1
27	Successful Pars Plana Vitrectomy with Zonulo-hyaloidectomy Performed 4 Years after the Onset of Chronic Low-Grade Aqueous Misdirection. <i>Middle East African Journal of Ophthalmology</i> , 2021, 28, 137-139.	0.5	1
28	Phacoemulsification after trabeculectomy in uveitis associated with Vogt-Koyanagi-Harada disease: intermediate-term visual outcome, IOP control and trabeculectomy survival. <i>BMC Ophthalmology</i> , 2022, 22, 210.	0.6	1
29	Genetic variant rs613872 in transcription factor 4 (TCF4) is not associated with primary open-angle glaucoma. <i>Eye</i> , 2021, 35, 696-697.	1.1	0