

# Rodrigue Lescouezec

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8861317/publications.pdf>

Version: 2024-02-01

85  
papers

4,515  
citations

116194

36  
h-index

116156

66  
g-index

93  
all docs

93  
docs citations

93  
times ranked

2673  
citing authors

#	ARTICLE	IF	CITATIONS
1	Molecular Magnetic Materials Based on $\{Co^{III}(Tp^*)(CN)_3\}^{\sim}$ Cyanidometallate: Combined Magnetic, Structural and $^{59}Co$ NMR Study. Chemistry - A European Journal, 2022, 28, .	1.7	2
2	Enlightening the Alkali Ion Role in the Photomagnetic Effect of FeCo Prussian Blue Analogues. Journal of the American Chemical Society, 2022, 144, 10888-10901.	6.6	16
3	The Atypical Hysteresis of $[Fe(C_6F_5Tp)_2]$ : Overlay of Spin Crossovers and Symmetry-Breaking Phase Transition. Angewandte Chemie - International Edition, 2021, 60, 8803-8807.	7.2	7
4	The Atypical Hysteresis of $[Fe(C_6F_5Tp)_2]$ : Overlay of Spin Crossovers and Symmetry-Breaking Phase Transition. Angewandte Chemie, 2021, 133, 8885-8889.	1.6	1
5	Field-induced single ion magnet behaviour of discrete and one-dimensional complexes containing $[bis(1\text{-methylimidazol-2-yl})\text{ketone}]cobalt$ building units. Dalton Transactions, 2021, 50, 16353-16363.	1.6	6
6	Observation of Protonation-Induced Spin State Switching in a Cyanido-Bridged $\{Fe_2Co_2\}$ Molecular Square. Inorganic Chemistry, 2021, 60, 17705-17714.	1.9	7
7	Thermo- and electro-switchable $Cs\{[Fe_4]^{Fe_4}\}$ cubic cage: spin-transition and electrochromism. Chemical Communications, 2020, 56, 10950-10953.	2.2	20
8	Two-Step Thermoinduced Metal-to-Metal Electron Transfer and ON/OFF Photoswitching in a Molecular $[Fe_2Co_2]$ Square Complex. Inorganic Chemistry, 2020, 59, 11879-11888.	1.9	36
9	Electrospray ionization: an efficient approach to deposit polymetallic molecular switches onto gold surfaces. Chemical Communications, 2020, 56, 6587-6589.	2.2	3
10	Pressure-Induced Conversion of a Paramagnetic FeCo Complex into a Molecular Magnetic Switch with Tuneable Hysteresis. Angewandte Chemie, 2020, 132, 17425-17429.	1.6	10
11	Pressure-Induced Conversion of a Paramagnetic FeCo Complex into a Molecular Magnetic Switch with Tuneable Hysteresis. Angewandte Chemie - International Edition, 2020, 59, 17272-17276.	7.2	29
12	Electron Transfer in the $Cs\{[Mn_4Fe_4]\}$ Cubic Switch: A Soluble Molecular Model of the MnFe Prussian Blue Analogues. Angewandte Chemie, 2020, 132, 8166-8170.	1.6	9
13	Electron Transfer in the $Cs\{[Mn_4Fe_4]\}$ Cubic Switch: A Soluble Molecular Model of the MnFe Prussian Blue Analogues. Angewandte Chemie - International Edition, 2020, 59, 8089-8093.	7.2	32
14	Probing the Local Magnetic Structure of the $[Fe^{III}(Tp)(CN)_3]^{\sim}$ Building Block Via Solid-State NMR Spectroscopy, Polarized Neutron Diffraction, and First-Principle Calculations. Chemistry - A European Journal, 2019, 25, 12120-12136.	1.7	9
15	Tetranuclear $[FeII_2FeIII_2]^{2+}$ molecular switches: $[FeII(bik)_2(N^{\ominus})_2]$ spin-crossover complexes containing $[FeIII(Tp)(CN)_3]^{\ominus}$ metalloligands as N-donor. Comptes Rendus Chimie, 2019, 22, 516-524.	0.2	12
16	Solid-state electrochemistry of metal cyanides. Comptes Rendus Chimie, 2019, 22, 483-489.	0.2	5
17	Direct Observation of Charge Transfer and Magnetism in $Fe_4Co_4$ Cyanide-Bridged Molecular Cubes. Journal of Physical Chemistry Letters, 2019, 10, 1799-1804.	2.1	15
18	A soluble cyanide-bridged $\{Fe_4Ni_4\}$ box encapsulating a $Cs^+$ ion: synthesis, structure and electronic properties. Journal of Coordination Chemistry, 2018, 71, 601-614.	0.8	10

#	ARTICLE	IF	CITATIONS
19	A [Fe <sup>III</sup> (Tp)(CN) <sub>3</sub> ] <sup>+</sup> scorpionate-based complex as a building block for designing ion storage hosts (Tp: hydrotrispyrazolylborate). <i>Chemical Communications</i> , 2018, 54, 5189-5192.	2.2	14
20	Solution and Solid-State Study of the Spin-Crossover [Fe <sup>II</sup> (R <sup>bik</sup> ) <sub>3</sub> ](BF <sub>4</sub> ) <sub>2</sub> Complexes (R = Me, Et, Vinyl). <i>European Journal of Inorganic Chemistry</i> , 2018, 2018, 414-428.	1.0	28
21	Thermally-Induced Spin Crossover and LIESST Effect in the Neutral [Fe <sup>II</sup> (Mebik) <sub>2</sub> (NCX) <sub>2</sub> ] Complexes: Variable-Temperature Structural, Magnetic, and Optical Studies (X = S, Se; Mebik =) <i>Tj ETQq1 1 0.784314 rgBT /Overlock 10 1650 657</i>	1.6	9
22	Synthesis, characterization and magnetism of homoleptic bis(5-aryl-2-iminopyrrolyl) complexes of iron(II) and cobalt(II). <i>Polyhedron</i> , 2018, 152, 179-187.	1.0	18
23	Probing the local structure of Prussian blue electrodes by <sup>113</sup> Cd NMR spectroscopy. <i>Dalton Transactions</i> , 2017, 46, 6159-6162.	1.6	9
24	A new {Fe <sub>4</sub> Co <sub>4</sub> } soluble switchable nanomagnet encapsulating Cs <sup>+</sup> : enhancing the stability and redox flexibility and tuning the photomagnetic effect.. <i>Dalton Transactions</i> , 2017, 46, 15549-15557.	1.6	32
25	K <sup>+</sup> , { [Fe <sup>II</sup> (Tp)(CN) <sub>3</sub> ] <sub>4</sub> [Co <sup>III</sup> ( <sup>pz</sup> Tp)] <sub>3</sub> [Co <sup>II</sup> ( <sup>ks</sup> )] <sub>3</sub> } a neutral soluble model complex of photomagnetic Prussian blue analogues. <i>Chemical Science</i> , 2016, 7, 4825-4831.	3.7	69
26	An {Fe <sub>6</sub> } tetrahedral cage: building nanoscopic molecular assemblies through cyanometallate and alkoxo linkers. <i>Dalton Transactions</i> , 2016, 45, 17610-17615.	1.6	15
27	Polarized Neutron Diffraction to Probe Local Magnetic Anisotropy of a Low-Spin Fe(III) Complex. <i>Angewandte Chemie</i> , 2016, 128, 4031-4035.	1.6	5
28	Polarized Neutron Diffraction to Probe Local Magnetic Anisotropy of a Low-Spin Fe(III) Complex. <i>Angewandte Chemie - International Edition</i> , 2016, 55, 3963-3967.	7.2	31
29	Innenr <sup>Ä</sup> 4cktitelbild: Polarized Neutron Diffraction to Probe Local Magnetic Anisotropy of a Low-Spin Fe(III) Complex ( <i>Angew. Chem.</i> 12/2016). <i>Angewandte Chemie</i> , 2016, 128, 4171-4171.	1.6	0
30	One synthesis: two redox states. Temperature-oriented crystallization of a charge transfer {Fe <sub>2</sub> Co <sub>2</sub> } square complex in a {Fe <sup>II</sup> LSCo <sup>III</sup> L} <sub>2</sub> diamagnetic or {Fe <sup>III</sup> LSCo <sup>II</sup> HS} <sub>2</sub> paramagnetic state. <i>RSC Advances</i> , 2016, 6, 17456-17459.	1.7	39
31	Versatile nano-platforms for hybrid systems: expressing spin-transition behavior on nanoparticles. <i>Journal of Materials Chemistry C</i> , 2015, 3, 3350-3355.	2.7	8
32	Photo- and thermo-induced spin crossover in a cyanide-bridged {Mo <sup>V</sup> <sub>2</sub> Fe <sup>II</sup> <sub>2</sub> } rhombus molecule. <i>Chemical Communications</i> , 2014, 50, 2893-2895.	2.2	59
33	Paramagnetic Hexacyanometalates. The Diversity of Spin Distribution Studied by <sup>13</sup> C and <sup>15</sup> N MAS NMR Spectroscopy. <i>Inorganic Chemistry</i> , 2013, 52, 12634-12644.	1.9	8
34	Combining Cyanometalates and Coordination Clusters: An Alternative Synthetic Route toward Original Molecular Materials. <i>Crystal Growth and Design</i> , 2013, 13, 4190-4194.	1.4	6
35	A cyanide and hydroxo-bridged nanocage: a new generation of coordination clusters. <i>Chemical Communications</i> , 2013, 49, 1181.	2.2	27
36	On/Off Photoswitching in a Cyanide-Bridged {Fe <sub>2</sub> Co <sub>2</sub> } Magnetic Molecular Square. <i>Journal of the American Chemical Society</i> , 2013, 135, 1653-1656.	6.6	143

#	ARTICLE	IF	CITATIONS
37	W $\pi$ Co Discrete Complex Exhibiting Photo- and Thermo-Induced Magnetisation. <i>Chemistry - A European Journal</i> , 2013, 19, 7682-7685.	1.7	44
38	Self-assembly, metal binding ability, and magnetic properties of dinickel(ii) and dicobalt(ii) triple mesocates. <i>CrystEngComm</i> , 2012, 14, 5639.	1.3	14
39	Photomagnetic effect in a cyanide-bridged mixed-valence {FeII2FeIII2} molecular square. <i>Chemical Communications</i> , 2012, 48, 5653.	2.2	84
40	Synthesis, Crystal Structures, and Magnetic Properties of a New Family of Heterometallic Cyanide-Bridged Fe <sup>III</sup> <sub>2</sub> M <sup>II</sup> <sub>2</sub> (M = Mn, Ni, and Co) Square Complexes. <i>Inorganic Chemistry</i> , 2011, 50, 6250-6262.	1.9	67
41	Probing Spin Density and Local Structure in the Prussian Blue Analogues CsCd[Fe/Co(CN) <sub>6</sub> ] $\cdot$ 0.5H <sub>2</sub> O and Cd <sub>3</sub> [Fe/Co(CN) <sub>6</sub> ] <sub>2</sub> $\cdot$ 15H <sub>2</sub> O with Solid-State MAS NMR Spectroscopy. <i>Chemistry - A European Journal</i> , 2011, 17, 11567-11575.	1.7	24
42	Rational Enantioselective Design of Chiral Heterobimetallic Single-Chain Magnets: Synthesis, Crystal Structures and Magnetic Properties of Oxamato-Bridged M <sup>II</sup> Cu <sup>II</sup> Chains (M=Mn, Co). <i>Chemistry - A European Journal</i> , 2011, 17, 12482-12494.	1.7	78
43	Single chain magnet behaviour in an enantiopure chiral cobalt(ii)-copper(ii) one-dimensional compound. <i>Chemical Communications</i> , 2010, 46, 2322.	2.2	100
44	Supramolecular coordination chemistry of aromatic polyoxalamide ligands: A metallosupramolecular approach toward functional magnetic materials. <i>Coordination Chemistry Reviews</i> , 2010, 254, 2281-2296.	9.5	178
45	Oligo-phenyleneoxalamide Copper(II) Mesocates as Electro-Switchable Ferromagnetic Metal-Organic Wires. <i>Chemistry - A European Journal</i> , 2010, 16, 12838-12851.	1.7	30
46	Synthesis, crystal structure and magnetic properties of two oxalato-bridged dimetallic trinuclear complexes combined with a polar cation. <i>Dalton Transactions</i> , 2010, 39, 4951.	1.6	35
47	Dimensionality Switching Through a Thermally Induced Reversible Single-Crystal-to-Single-Crystal Phase Transition in a Cyanide Complex. <i>Inorganic Chemistry</i> , 2010, 49, 11045-11056.	1.9	38
48	Tuning the Spin Ground State in Heterononanuclear Nickel(II)-Copper(II) Cylinders with a Triangular Metallacyclophane Core. <i>Inorganic Chemistry</i> , 2010, 49, 11264-11266.	1.9	5
49	Variation of the ground spin state in homo- and hetero-octanuclear copper(ii) and nickel(ii) double-star complexes with a meso-helicate-type metallacryptand core. <i>Dalton Transactions</i> , 2010, 39, 4786.	1.6	11
50	One dimensional assembly of Mn <sub>6</sub> single molecule magnets linked by oligothiophene bridges. <i>Dalton Transactions</i> , 2010, 39, 4751.	1.6	29
51	[FeII <sub>2</sub> CoII <sub>2</sub> LS] $^{2+}$ [FeII <sub>2</sub> LSCoII <sub>2</sub> HS] <sub>2</sub> photoinduced conversion in a cyanide-bridged heterobimetallic molecular square. <i>Chemical Communications</i> , 2010, 46, 8995.	2.2	113
52	Revisiting Prussian Blue Analogues with Solid-State MAS NMR Spectroscopy: Spin Density and Local Structure in [Cd <sub>3</sub> {Fe(CN) <sub>6</sub> ] <sub>2</sub> $\cdot$ 15H <sub>2</sub> O. <i>Angewandte Chemie - International Edition</i> , 2009, 48, 1673-1676.	7.2	35
53	Ferromagnetic Coupling by Spin Polarization in a Trinuclear Copper(II) Metallacyclophane with a Triangular Cage-Like Structure. <i>Inorganic Chemistry</i> , 2009, 48, 5244-5249.	1.9	47
54	Redox Switch-Off of the Ferromagnetic Coupling in a Mixed-Spin Tricobalt(II) Triple Mesocate. <i>Journal of the American Chemical Society</i> , 2009, 131, 14614-14615.	6.6	39

#	ARTICLE	IF	CITATIONS
55	Molecular-Programmed Self-Assembly of Homo- and Heterometallic Tetranuclear Coordination Compounds: Synthesis, Crystal Structures, and Magnetic Properties of Rack-Type $\text{Cu}_2\text{M}_2$ Complexes (M = Cu and Ni) with Tetranucleating Phenylenedioxamate Bridging Ligands. <i>Inorganic Chemistry</i> , 2009, 48, 4661-4673.	1.9	22
56	A Metallacryptand-Based Manganese(II)-Cobalt(II) Ferrimagnet with a Three-Dimensional Honeycomb Open-Framework Architecture. <i>Angewandte Chemie - International Edition</i> , 2008, 47, 4211-4216.	7.2	41
57	Rational design of a new class of heterobimetallic molecule-based magnets: Synthesis, crystal structures, and magnetic properties of oxamate-bridged ( $\text{M}^2=\text{LiI}$ and $\text{MnII}$ ; $\text{M}=\text{NiII}$ and $\text{CoII}$ ) open-frameworks with a three-dimensional honeycomb architecture. <i>Inorganica Chimica Acta</i> , 2008, 361, 3394-3402.	1.2	49
58	Synthesis, crystal structure and magnetic properties of a new cyanide-bridged iron(III)-nickel(II) ferromagnetic chain. <i>Inorganica Chimica Acta</i> , 2008, 361, 3912-3918.	1.2	25
59	Ligand design for multidimensional magnetic materials: a metallosupramolecular perspective. <i>Dalton Transactions</i> , 2008, , 2780.	1.6	244
60	4,2-Ribbon like ferromagnetic cyano-bridged $\text{FeIII}_2\text{NiII}$ chains: a magneto-structural study. <i>Dalton Transactions</i> , 2007, , 3690.	1.6	43
61	Magnetic Face-to-Face Interaction and Electrocommunication in Chromium Sandwich Compounds. <i>Chemistry - A European Journal</i> , 2007, 13, 1191-1200.	1.7	10
62	Solid-State Anion-Guest Encapsulation by Metallosupramolecular Capsules Made from Two Tetranuclear Copper(II) Complexes. <i>European Journal of Inorganic Chemistry</i> , 2007, 2007, 4569-4573.	1.0	9
63	$[\text{Fe}(\text{bpym})(\text{CN})_4]$ : A New Building Block for Designing Single-Chain Magnets. <i>Journal of the American Chemical Society</i> , 2006, 128, 4842-4853.	6.6	248
64	Design of single cyanide-bridged tetranuclear bimetallic rectangles exhibiting ferromagnetic coupling. <i>Inorganic Chemistry Communication</i> , 2005, 8, 382-385.	1.8	29
65	Design of single chain magnets through cyanide-bearing six-coordinate complexes. <i>Coordination Chemistry Reviews</i> , 2005, 249, 2691-2729.	9.5	417
66	Charge transfer salts containing a paramagnetic cyano-complex and iodine substituted organic donor involving $\text{I}^{\text{I}}(\text{donor})\cdots\text{N}^{\text{II}}(\text{anion})$ -interactions. <i>Comptes Rendus Chimie</i> , 2005, 8, 1286-1297.	0.2	27
67	$[\text{Cr}^{\text{III}}(\text{L})(\text{CN})_4]^{2-}$ : a new building block in designing cyanide-bridged 4,2-ribbon-like chains $\{[\text{Cr}^{\text{III}}(\text{L})(\text{CN})_4]_2\text{Mn}(\text{H}_2\text{O})_2\}_n \cdot n\text{H}_2\text{O}$ [L = 2-aminomethylpyridine (n = 6) and 1,10-phenanthroline (n = 4)]. <i>New Journal of Chemistry</i> , 2005, 29, 210-219.	1.4	41
68	Synthesis, crystal structures and magnetic properties of cyanide- and phenolate-bridged $[\text{M}^{\text{III}}\text{Ni}^{\text{II}}]_2$ tetranuclear complexes (M = Fe and Cr). <i>Dalton Transactions</i> , 2005, , 1357-1364.	1.6	65
69	Highly Resolved Spin-Density Distribution in the Prussian-Blue Precursors $\text{Cs}_2\text{K}[\text{Fe}(\text{CN})_6]$ and $\text{Cs}_2\text{K}[\text{Mn}(\text{CN})_6]$ . <i>Angewandte Chemie - International Edition</i> , 2004, 43, 2571-2573.	7.2	34
70	Highly Resolved Spin-Density Distribution in the Prussian-Blue Precursors $\text{Cs}_2\text{K}[\text{Fe}(\text{CN})_6]$ and $\text{Cs}_2\text{K}[\text{Mn}(\text{CN})_6]$ . <i>ChemInform</i> , 2004, 35, no.	0.1	0
71	Nuclearity Controlled Cyanide-Bridged Bimetallic $\text{Cr}^{\text{III}}\text{-Mn}^{\text{II}}$ Compounds: Synthesis, Crystal Structures, Magnetic Properties and Theoretical Calculations. <i>Chemistry - A European Journal</i> , 2004, 10, 6130-6145.	1.7	94
72	mer- $[\text{Fe}^{\text{III}}(\text{bpca})(\text{CN})_3]$ : A New Low-Spin Iron(III) Complex to Build Heterometallic Ladder-like Chains. <i>Inorganic Chemistry</i> , 2004, 43, 2234-2236.	1.9	101

#	ARTICLE	IF	CITATIONS
73	Title is missing!. <i>Angewandte Chemie</i> , 2003, 115, 1521-1524.	1.6	79
74	Cyanide-Bridged Iron(III)–Cobalt(II) Double Zigzag Ferromagnetic Chains: Two New Molecular Magnetic Nanowires. <i>Angewandte Chemie - International Edition</i> , 2003, 42, 1483-1486.	7.2	353
75	[Cr(AA)(C <sub>2</sub> O <sub>4</sub> ) <sub>2</sub> ] <sup>n-</sup> and [Cu(bpca)] <sup>+</sup> as building blocks in designing new oxalato-bridged Cr(III)–Cu(I) compounds [AA=2,2'-bipyridine and 1,10-phenanthroline; bpca=bis(2-pyridylcarbonyl)amide anion]. <i>Inorganica Chimica Acta</i> , 2003, 350, 131-142.	1.2	53
76	Cyanide-bridged Fe(III)–Co(II) bis double zigzag chains with a slow relaxation of the magnetisation. <i>Chemical Communications</i> , 2003, , 1850-1851.	2.2	154
77	Ferromagnetic Coupling between Low- and High-Spin Iron(III) Ions in the Tetranuclear Complex fac-[Fe(III)(HB(pz) <sub>3</sub> )(CN) <sub>2</sub> ] <sub>3</sub> [Fe(III)(H <sub>2</sub> O) <sub>3</sub> ] <sub>3</sub> ·6H <sub>2</sub> O ([HB(pz) <sub>3</sub> ]= Hydrotris(1-pyrazolyl)borate). <i>Inorganic Chemistry</i> , 2002, 41, 5943-5945.	1.9	124
78	[Fe(bipy)(CN) <sub>4</sub> ]- as a Versatile Building Block for the Design of Heterometallic Systems: Synthesis, Crystal Structure, and Magnetic Properties of PPh <sub>4</sub> [Fe(III)(bipy)(CN) <sub>4</sub> ] <sup>-</sup> ·H <sub>2</sub> O, [Fe(III)(bipy)(CN) <sub>4</sub> ] <sup>-</sup> ·2Mn(II)(H <sub>2</sub> O) <sub>4</sub> ·4H <sub>2</sub> O, and [Fe(III)(bipy)(CN) <sub>4</sub> ] <sup>-</sup> ·2Zn(II)·2H <sub>2</sub> O [bipy = 2,2'-Bipyridine; M = Mn and Zn]. <i>Inorganic Chemistry</i> , 2002, 41, 818-826.	1.9	95
79	[Mn <sub>2</sub> (bipym)(H <sub>2</sub> O) <sub>8</sub> ] <sup>4+</sup> and [Fe(bipy)(CN) <sub>4</sub> ] <sup>-</sup> as building blocks in designing novel bipym- and cyanide-bridged heterobimetallic complexes (bipym = 2,2'-bipyrimidine and bipy = 2,2'-bipyridine). <i>Dalton Transactions RSC</i> , 2002, , 3171-3176.	2.3	46
80	[Cr(bpym)(C <sub>2</sub> O <sub>4</sub> ) <sub>2</sub> ] <sup>n-</sup> in designing heterometallic complexes. Crystal structures and magnetic properties of PPh <sub>4</sub> [Cr(III)(C <sub>2</sub> O <sub>4</sub> ) <sub>2</sub> ] <sup>n-</sup> ·H <sub>2</sub> O and [Ag(bpym)][Cr(C <sub>2</sub> O <sub>4</sub> ) <sub>2</sub> ](H <sub>2</sub> O) <sub>2</sub> ·2H <sub>2</sub> O (bpym=2,2'-bipyrimidine). <i>Inorganica Chimica Acta</i> , 2002, 336, 46-54.	1.2	44
81	[Fe(Phen)(CN) <sub>4</sub> ] <sup>-</sup> : A Versatile Building Block for the Design of Heterometallic Systems. Crystal Structures and Magnetic Properties of PPh <sub>4</sub> [Fe(III)(Phen)(CN) <sub>4</sub> ] <sup>-</sup> ·2H <sub>2</sub> O and [Fe(III)(Phen)(CN) <sub>4</sub> ] <sup>-</sup> ·2Mn(II)(H <sub>2</sub> O) <sub>2</sub> ·4H <sub>2</sub> O [Phen = 1,10-Phenanthroline; M = Mn(II) and Zn(II)]. <i>Inorganic Chemistry</i> , 2001, 40, 2065-2072.	1.9	107
82	[Cr(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> : a new bis-oxalato building block for the design of heteropolymetallic systems. Crystal structures and magnetic properties of PPh <sub>4</sub> [Cr(III)(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> , AsPh <sub>4</sub> [Cr(III)(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> , Hdpa[Cr(III)(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> ·4H <sub>2</sub> O, Rad[Cr(III)(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> ·H <sub>2</sub> O and Sr[Cr(III)(dpa)(ox) <sub>2</sub> ] <sup>n-</sup> ·8H <sub>2</sub> O (dpa = 2,2'-dipyridylamine). <i>New Journal of Chemistry</i> , 2001, 25, 1224-1235.	1.4	42
83	[Cr(phen)(ox) <sub>2</sub> ] <sup>n-</sup> : a versatile bis-oxalato building block for the design of heteropolymetallic systems. Crystal structures and magnetic properties of AsPh <sub>4</sub> [Cr(III)(phen)(ox) <sub>2</sub> ] <sup>n-</sup> ·H <sub>2</sub> O, [NaCr(III)(phen)(ox) <sub>2</sub> ] <sup>n-</sup> ·2H <sub>2</sub> O and {[Cr(III)(phen)(ox) <sub>2</sub> ] <sup>n-</sup> ·2[Mn <sub>2</sub> (bpy) <sub>2</sub> (H <sub>2</sub> O) <sub>2</sub> (ox)] <sup>n-</sup> ·6H <sub>2</sub> O. <i>New Journal of Chemistry</i> , 2000, 24, 527-536.	1.4	90
84	Two-Dimensional Assembling of (2,2'-Bipyrimidine)bis(oxalato)chromate(III) Units through Alkaline Cations. <i>Inorganic Chemistry</i> , 1999, 38, 2234-2237.	1.9	53
85	Building responsive materials by assembling {Fe <sub>4</sub> Co <sub>4</sub> } switchable molecular cubes. <i>Journal of Materials Chemistry C</i> , 0, , .	2.7	8