

Grethe Skretting

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8868738/publications.pdf>

Version: 2024-02-01

24
papers

302
citations

840776

11
h-index

888059

17
g-index

24
all docs

24
docs citations

24
times ranked

427
citing authors

#	ARTICLE	IF	CITATIONS
1	Downregulation of TFPI in breast cancer cells induces tyrosine phosphorylation signaling and increases metastatic growth by stimulating cell motility. <i>BMC Cancer</i> , 2011, 11, 357.	2.6	40
2	TFPI ¹ and TFPI ² are expressed at the surface of breast cancer cells and inhibit TF-FVIIa activity. <i>Journal of Hematology and Oncology</i> , 2013, 6, 5.	17.0	27
3	EPAS1/HIF-2 alpha-mediated downregulation of tissue factor pathway inhibitor leads to a pro-thrombotic potential in endothelial cells. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2016, 1862, 670-678.	3.8	27
4	Overexpression of both TFPI ¹ and TFPI ² induces apoptosis and expression of genes involved in the death receptor pathway in breast cancer cells. <i>Molecular Carcinogenesis</i> , 2010, 49, 951-963.	2.7	25
5	A novel hypoxia response element regulates oxygen-related repression of tissue factor pathway inhibitor in the breast cancer cell line MCF-7. <i>Thrombosis Research</i> , 2017, 157, 111-116.	1.7	21
6	Syndecan-3 and TFPI Colocalize on the Surface of Endothelial-, Smooth Muscle-, and Cancer Cells. <i>PLoS ONE</i> , 2015, 10, e0117404.	2.5	21
7	Functional characterization of polymorphisms in the human TFPI gene. <i>Biochemical and Biophysical Research Communications</i> , 2010, 397, 106-111.	2.1	19
8	Tissue factor pathway inhibitor attenuates ER stress-induced inflammation in human M2-polarized macrophages. <i>Biochemical and Biophysical Research Communications</i> , 2017, 491, 442-448.	2.1	19
9	Oestrogen induced downregulation of TFPI expression is mediated by ER ¹ . <i>Thrombosis Research</i> , 2014, 134, 138-143.	1.7	14
10	Tissue factor pathway inhibitor polymorphisms in women with and without a history of venous thrombosis and the effects of postmenopausal hormone therapy. <i>Blood Coagulation and Fibrinolysis</i> , 2010, 21, 516-521.	1.0	12
11	Factor VII deficiency: Unveiling the cellular and molecular mechanisms underlying three model alterations of the enzyme catalytic domain. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018, 1864, 660-667.	3.8	11
12	The chemical chaperone sodium 4-phenylbutyrate improves the secretion of the protein CA267T mutant in CHO-K1 cells through the GRASP55 pathway. <i>Cell and Bioscience</i> , 2015, 5, 57.	4.8	9
13	Normalization of disrupted clock gene expression in males with tetraplegia: a crossover randomized placebo-controlled trial of melatonin supplementation. <i>Spinal Cord</i> , 2018, 56, 1076-1083.	1.9	9
14	Determinants of acquired activated protein C resistance and D-dimer in breast cancer. <i>Thrombosis Research</i> , 2016, 145, 78-83.	1.7	8
15	The effect of the chemical chaperone 4-phenylbutyrate on secretion and activity of the p.Q160R missense variant of coagulation factor FVII. <i>Cell and Bioscience</i> , 2019, 9, 69.	4.8	8
16	Indirect regulation of TFPI-2 expression by miR-494 in breast cancer cells. <i>Scientific Reports</i> , 2020, 10, 4036.	3.3	7
17	Estrogen induced expression of tissue factor pathway inhibitor-2 in MCF7 cells involves lysine-specific demethylase 1. <i>Molecular and Cellular Endocrinology</i> , 2017, 443, 80-88.	3.2	6
18	Oestrogens Downregulate Tissue Factor Pathway Inhibitor through Oestrogen Response Elements in the 5' Flanking Region. <i>PLoS ONE</i> , 2016, 11, e0152114.	2.5	6

#	ARTICLE	IF	CITATIONS
19	Increased expression of TFPI in human carotid stenosis. <i>Thrombosis Research</i> , 2017, 155, 31-37.	1.7	4
20	Activation of Endoplasmic Reticulum Stress and Unfolded Protein Response in Congenital Factor VII Deficiency. <i>Thrombosis and Haemostasis</i> , 2018, 47, 664-675.	3.4	4
21	Transcription factor FOXP3: A repressor of the <i>TFPI</i> gene?. <i>Journal of Cellular Biochemistry</i> , 2019, 120, 12924-12936.	2.6	3
22	Overexpression of tissue factor pathway inhibitor in CHO-K1 cells results in increased activation of NF- κ B and apoptosis mediated by a caspase-3 independent pathway. <i>Molecular Biology Reports</i> , 2012, 39, 10089-10096.	2.3	2
23	Candidate Gene Polymorphisms and the Risk for Pregnancy Related Venous Thrombosis. <i>Blood</i> , 2010, 116, 4203-4203.	1.4	0
24	The Chemical Chaperone 4-Phenylbutyrate Increases Secretion and Activity of Missense and Elongated Factor VII Mutants. <i>Blood</i> , 2018, 132, 3777-3777.	1.4	0