

# Serhat ÆolakoÄlu

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8869130/publications.pdf>

Version: 2024-02-01

18  
papers

83  
citations

1683354

5  
h-index

1473754

9  
g-index

18  
all docs

18  
docs citations

18  
times ranked

131  
citing authors

#	ARTICLE	IF	CITATIONS
1	Proximate and Elemental Composition of <i>Chamelea gallina</i> from the Southern Coast of the Marmara Sea (Turkey). <i>Biological Trace Element Research</i> , 2011, 143, 983-991.	1.9	16
2	Some population parameters of <i>Ruditapes philippinarum</i> (Bivalvia, Veneridae) on the southern coast of the Marmara Sea, Turkey. <i>Helgoland Marine Research</i> , 2014, 68, 539-548.	1.3	16
3	Gross pathology, blood chemistry, lipid and peroxide contents in rainbow trout ( <i>Oncorhynchus</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock temperature. <i>Acta Veterinaria Hungarica</i> , 2009, 57, 305-317.	0.2	9
4	Metal levels in economically important bivalve species from Turkey. <i>Food Additives and Contaminants: Part B Surveillance</i> , 2012, 5, 272-278.	1.3	7
5	Title is missing!. <i>Turkish Journal of Fisheries and Aquatic Sciences</i> , 2014, 14, .	0.4	6
6	Bycatch and discards from two types of bivalve dredges targeting <i>Donax trunculus</i> and <i>Chamelea gallina</i> used in the southern coast of the Marmara Sea, Turkey. <i>Fisheries Science</i> , 2020, 86, 995-1004.	0.7	6
7	Depuration Times of <i>Donax trunculus</i> and <i>Tapes decussatus</i> . <i>Brazilian Journal of Microbiology</i> , 2014, 45, 1017-1022.	0.8	5
8	Population Structure and Dynamics of Warty Venus, <i>Venus verrucosa</i> (Bivalvia, Veneridae), in North Aegean Sea, Turkey. <i>Journal of Shellfish Research</i> , 2015, 34, 347-354.	0.3	4
9	Bioaccumulation monitoring of chemical contaminants in mussels <i>Mytilus galloprovincialis</i> from the southern coast of the Marmara Sea, Turkey. <i>Turkish Journal of Veterinary and Animal Sciences</i> , 2020, 44, 235-243.	0.2	4
10	Levels of PAHs, PCBs, and toxic metals in <i>Ruditapes philippinarum</i> and <i>Donax trunculus</i> in Marmara Sea, Turkey. <i>Journal of the Science of Food and Agriculture</i> , 2021, 101, 1167-1173.	1.7	3
11	Catch composition of set net (fixed stake trap) fisheries in the coastal waters of Saros Bay, North Aegean Sea. <i>Su ĀcerĀ¼nleri Dergisi</i> , 2015, 32, 53-58.	0.1	3
12	Distribution of Macrozoobenthic Invertebrates in the South Marmara Sea. <i>Journal of Animal and Veterinary Advances</i> , 2010, 9, 471-473.	0.1	2
13	Antioxidant Activity of <i>Mytilus galloprovincialis</i> and <i>Ruditapes philippinarum</i> . <i>Turkish Journal of Agriculture: Food Science and Technology</i> , 0, 7, 150-153.	0.1	2
14	BatĀ± Marmaraâ€™da Kum Midyesi ( <i>Chamelea gallina</i> L., 1758) ve Kum ĀžĀ±rlanĀ± ( <i>Donax trunculus</i> L., 1758) PopulasyonlarĀ±nĀ±n BĀ¼yĀ¼me Ā–zelliklerinin Ā°ncelenmesi. <i>Journal of Fisheriesciencescom</i> , 2014, , .	0.2	0
15	Length-weight relationships of fishes caught by stationary uncovered pound nets in the coastal waters of Saros Bay, North Aegean Sea (Turkey). <i>Su ĀcerĀ¼nleri Dergisi</i> , 2021, 38, 95-99.	0.1	0
16	TĀ¼rkiyeâ€™de ticari Āšift kabuklu yumuĀ¼akĀša avcĀ±lĀ±Ā¼Ā±nda kullanĀ±lan av araĀšlarĀ±. <i>Su ĀcerĀ¼nleri Dergisi</i> , 2017, 34, 415-421.	0.1	0
17	Seasonal variations and human health risk assessment of trace elements in the bivalve ecosystem in the Sea of Marmara. <i>Food Additives and Contaminants: Part B Surveillance</i> , 0, , 1-14.	1.3	0
18	Stuffed mussels in Turkish culinary culture: Ottoman Empire period. <i>Su ĀcerĀ¼nleri Dergisi</i> , 2022, 39, 170-173.	0.1	0