Tieba Millogo

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8879623/publications.pdf

Version: 2024-02-01

840585 839398 36 435 11 18 citations h-index g-index papers 38 38 38 561 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Evaluation of the quality of maternal death review cycles in Burkina Faso. International Journal of Gynecology and Obstetrics, 2022, , .	1.0	2
2	Time to Long-Acting Reversible Contraceptive Uptake Over Twelve Months Postpartum: Findings of the Yam Daabo Cluster Randomized-Controlled Trial in Burkina Faso and the Democratic Republic of the Congo. Open Access Journal of Contraception, 2021, Volume 12, 73-82.	0.6	3
3	Yam Daabo interventions' effects on postpartum family planning use in Burkina Faso at 24 months after childbirth. BMC Public Health, 2021, 21, 946.	1.2	2
4	Effectiveness of the World Health Organization Safe Childbirth Checklist (WHO-SCC) in preventing poor childbirth outcomes: a study protocol for a matched-pair cluster randomized control trial. BMC Public Health, 2021, 21, 1652.	1.2	1
5	Determinants of childbirth care quality along the care continuum in limited resource settings: A structural equation modeling analysis of cross-sectional data from Burkina Faso and CÃte d'lvoire. BMC Pregnancy and Childbirth, 2021, 21, 848.	0.9	0
6	Quality of maternal and newborn care in limited-resource settings: a facility-based cross-sectional study in Burkina Faso and Cà te d'Ivoire. BMJ Open, 2020, 10, e036121.	0.8	7
7	Discontinuation and switching of postpartum contraceptive methods over twelve months in Burkina Faso and the Democratic Republic of the Congo: a secondary analysis of the Yam Daabo trial. Contraception and Reproductive Medicine, 2020, 5, 35.	0.7	4
8	Feasibility study on the adoption of the WHO safe childbirth checklist by front-line healthcare providers and managers in Burkina Faso and Cà te d'lvoire. Pilot and Feasibility Studies, 2020, 6, 150.	0.5	6
9	Prevalence and risk factors for overweight and obesity: a cross-sectional countrywide study in Burkina Faso. BMJ Open, 2020, 10, e032953.	0.8	11
10	Post-partum family planning in Burkina Faso (Yam Daabo): a two group, multi-intervention, single-blinded, cluster-randomised controlled trial. The Lancet Global Health, 2019, 7, e1109-e1117.	2.9	21
11	Tobacco use and associated risk factors in Burkina Faso: results from a population-based cross-sectional survey. BMC Public Health, 2019, 19, 1466.	1.2	11
12	Distribution of ABO and RHD blood group antigens in blood donors in Burkina Faso. International Journal of Immunogenetics, 2019, 46, 1-6.	0.8	11
13	Task sharing for family planning services, Burkina Faso. Bulletin of the World Health Organization, 2019, 97, 783-788.	1.5	10
14	Diabetes and abnormal glucose regulation in the adult population of Burkina Faso: prevalence and predictors. BMC Public Health, 2018, 18, 350.	1.2	24
15	Birth spacing and informed family planning choices after childbirth in Burkina Faso and the Democratic Republic of Congo: Participatory action research to design and evaluate a decision-making tool for providers and their clients. Patient Education and Counseling, 2018, 101, 1871-1875.	1.0	7
16	Traceability of Blood Transfusions and Reporting of Adverse Reactions in Developing Countries: A Six-Year Postpilot Phase Experience in Burkina Faso. Advances in Hematology, 2018, 2018, 1-9.	0.6	6
17	Impact Evaluation of Seasonal Malaria Chemoprevention under Routine Program Implementation: A Quasi-Experimental Study in Burkina Faso. American Journal of Tropical Medicine and Hygiene, 2018, 98, 524-533.	0.6	46
18	Continuation of subcutaneous or intramuscular injectable contraception when administered by facility-based and community health workers: findings from a prospective cohort study in Burkina Faso and Uganda. Contraception, 2018, 98, 423-429.	0.8	20

#	Article	IF	CITATIONS
19	Participatory action research to identify a package of interventions to promote postpartum family planning in Burkina Faso and the Democratic Republic of Congo. BMC Women's Health, 2018, 18, 122.	0.8	9
20	Prevalence and factors associated with hypertension in Burkina Faso: a countrywide cross-sectional study. BMC Public Health, 2017, 17, 64.	1.2	49
21	Clinical outreach refresher trainings in crisis settings (S-CORT): <i>clinical management of sexual violence survivors</i> and <i>manual vacuum aspiration</i> in Burkina Faso, Nepal, and South Sudan. Reproductive Health Matters, 2017, 25, 103-113.	1.3	7
22	Inadequate programming, insufficient communication and non-compliance with the basic principles of maternal death audits in health districts in Burkina Faso: a qualitative study. Reproductive Health, 2017, 14, 121.	1.2	10
23	Evaluation of the implementation fidelity of the seasonal malaria chemoprevention intervention in Kaya health district, Burkina Faso. PLoS ONE, 2017, 12, e0187460.	1.1	17
24	Factors associated with very early neonatal mortality in Burkina Faso: A matched case–control study. International Journal of Gynecology and Obstetrics, 2016, 135, S93-S97.	1.0	6
25	Rate of and factors associated with indications for cesarean deliveries: Results of a national review in Burkina Faso. International Journal of Gynecology and Obstetrics, 2016, 135, S51-S57.	1.0	8
26	Trends and changes in home deliveries in Kassena–Nankana districts in northern Ghana: Results from repeated crossâ€sectional surveys. International Journal of Gynecology and Obstetrics, 2016, 135, S33-S38.	1.0	1
27	Factors associated with fresh stillbirths: A hospitalâ€based, matched, case–control study in Burkina Faso. International Journal of Gynecology and Obstetrics, 2016, 135, S98-S102.	1.0	8
28	Availability and utilization of obstetric and newborn care in Guinea: A national needs assessment. International Journal of Gynecology and Obstetrics, 2016, 135, S2-S6.	1.0	10
29	Predictors of mortality of low birth weight newborns during the neonatal period: A cohort study in two health districts of Burkina Faso. International Journal of Gynecology and Obstetrics, 2016, 135, S89-S92.	1.0	13
30	A case–control study of risk factors for maternal mortality in Burkina Faso in 2014. International Journal of Gynecology and Obstetrics, 2016, 135, S79-S83.	1.0	3
31	A case-control study of risk factors for surgical site infection after cesarean delivery in eastern Burkina Faso. International Journal of Gynecology and Obstetrics, 2016, 135, S107-S110.	1.0	8
32	Abolishing Fees at Health Centers in the Context of Community Case Management of Malaria: What Effects on Treatment-Seeking Practices for Febrile Children in Rural Burkina Faso?. PLoS ONE, 2015, 10, e0141306.	1.1	23
33	Domestic Larval Control Practices and Malaria Prevalence among Under-Five Children in Burkina Faso. PLoS ONE, 2015, 10, e0141784.	1.1	4
34	Audit of cesarean delivery in Burkina Faso. International Journal of Gynecology and Obstetrics, 2014, 125, 214-218.	1.0	19
35	The Kaya HDSS, Burkina Faso: a platform for epidemiological studies and health programme evaluation. International Journal of Epidemiology, 2013, 42, 741-749.	0.9	36
36	Assessment of the maternal death surveillance and response implementation process in Burkina Faso. International Journal of Gynecology and Obstetrics, 0, , .	1.0	4