

Karl-Norbert Klotz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8881982/publications.pdf>

Version: 2024-02-01

145
papers

5,715
citations

66343
42
h-index

91884
69
g-index

154
all docs

154
docs citations

154
times ranked

5059
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | 8-Cyclopentyl-1,3-dipropylxanthine (DPCPX) ? a selective high affinity antagonist radioligand for A1 adenosine receptors. Naunyn-Schmiedeberg's Archives of Pharmacology, 1987, 336, 204-210. | 3.0 | 367 |
| 2 | Adenosine receptors and their ligands. Naunyn-Schmiedeberg's Archives of Pharmacology, 2000, 362, 382-391. | 3.0 | 284 |
| 3 | Comparative pharmacology of human α_2 -adrenergic receptor subtypes?characterization of stably transfected receptors in CHO cells. Naunyn-Schmiedeberg's Archives of Pharmacology, 2004, 369, 151-159. | 3.0 | 279 |
| 4 | Analysis of receptor oligomerization by FRAP microscopy. Nature Methods, 2009, 6, 225-230. | 19.0 | 187 |
| 5 | Ectonucleotidases CD39 and CD73 on OvCA cells are potent adenosine-generating enzymes responsible for adenosine receptor 2A-dependent suppression of T cell function and NK cell cytotoxicity. Cancer Immunology, Immunotherapy, 2011, 60, 1405-1418. | 4.2 | 163 |
| 6 | The Small Antitumoral Immune Response Modifier Imiquimod Interacts with Adenosine Receptor Signaling in a TLR7- and TLR8-Independent Fashion. Journal of Investigative Dermatology, 2006, 126, 1338-1347. | 0.7 | 138 |
| 7 | Adenosine Receptors as Mediators of Both Cell Proliferation and Cell Death of Cultured Human Melanoma Cells. Journal of Investigative Dermatology, 2002, 119, 923-933. | 0.7 | 134 |
| 8 | Water-Soluble Phosphate Prodrugs of 1-Propargyl-8-styrylxanthine Derivatives, A2A-Selective Adenosine Receptor Antagonists. Journal of Medicinal Chemistry, 2000, 43, 440-448. | 6.4 | 129 |
| 9 | 2-Chloro-N6-[3H]cyclopentyladenosine ([3HCCPA) ?a high affinity agonist radioligand for A1 adenosine receptors. Naunyn-Schmiedeberg's Archives of Pharmacology, 1989, 340, 679-683. | 3.0 | 124 |
| 10 | 2-Chloro-N6-cyclopentyladenosine: a highly selective agonist at A1 adenosine receptors. Naunyn-Schmiedeberg's Archives of Pharmacology, 1988, 337, 687-9. | 3.0 | 120 |
| 11 | Adenosine A ₁ Receptor-Mediated Activation of Phospholipase C in Cultured Astrocytes Depends on the Level of Receptor Expression. Journal of Neuroscience, 1997, 17, 4956-4964. | 3.6 | 114 |
| 12 | N6-Alkyl-2-alkynyl Derivatives of Adenosine as Potent and Selective Agonists at the Human Adenosine A3 Receptor and a Starting Point for Searching A2B Ligands. Journal of Medicinal Chemistry, 2002, 45, 3271-3279. | 6.4 | 104 |
| 13 | Synthesis, Biological Activity, and Molecular Modeling Investigation of New Pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine Derivatives as Human A3 Adenosine Receptor Antagonists. Journal of Medicinal Chemistry, 2002, 45, 770-780. | 6.4 | 99 |
| 14 | 2-Alkynyl derivatives of adenosine and adenosine-5'-N-ethyluronamide as selective agonists at A2 adenosine receptors. Journal of Medicinal Chemistry, 1992, 35, 2363-2368. | 6.4 | 98 |
| 15 | Pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine Derivatives as Highly Potent and Selective Human A3 Adenosine Receptor Antagonists: A Influence of the Chain at the N8 Pyrazole Nitrogen. Journal of Medicinal Chemistry, 2000, 43, 4768-4780. | 6.4 | 89 |
| 16 | New substituted 9-alkylpurines as adenosine receptor ligands. Bioorganic and Medicinal Chemistry, 1998, 6, 523-533. | 3.0 | 82 |
| 17 | synthesis of new α_2 -isoxazoline derivatives and their pharmacological characterization as α_2 -adrenergic receptor antagonists. Bioorganic and Medicinal Chemistry, 1998, 6, 401-408. | 3.0 | 81 |
| 18 | Medicinal Chemistry and Pharmacology of A2B Adenosine Receptors. Current Topics in Medicinal Chemistry, 2003, 3, 427-443. | 2.1 | 81 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Pyrazolo[4,3-e]-1,2,4-triazolo[1,5-c]pyrimidine Derivatives as Highly Potent and Selective Human A3Adenosine Receptor Antagonists. <i>Journal of Medicinal Chemistry</i> , 1999, 42, 4473-4478. | 6.4 | 80 |
| 20 | Human breast cancer cell line MDA-MB-231 expresses endogenous A2B adenosine receptors mediating a Ca ²⁺ signal. <i>British Journal of Pharmacology</i> , 2005, 145, 211-218. | 5.4 | 65 |
| 21 | Role of the β 1-Adrenergic Pathway in Anesthetic and Ischemic Preconditioning against Myocardial Infarction in the Rabbit Heart In Vivo. <i>Anesthesiology</i> , 2006, 105, 503-510. | 2.5 | 64 |
| 22 | [3H]HEMADO a novel tritiated agonist selective for the human adenosine A3 receptor. <i>European Journal of Pharmacology</i> , 2007, 556, 14-18. | 3.5 | 60 |
| 23 | Neutrophil chemoattractant receptors and the membrane skeleton. <i>BioEssays</i> , 1994, 16, 193-198. | 2.5 | 59 |
| 24 | Expression, Pharmacological Profile, and Functional Coupling of A2B Receptors in a Recombinant System and in Peripheral Blood Cells Using a Novel Selective Antagonist Radioligand, [3H]MRE 2029-F20. <i>Molecular Pharmacology</i> , 2005, 67, 2137-2147. | 2.3 | 58 |
| 25 | Separation of solubilized A2 adenosine receptors of human platelets from non-receptor [3H]NECA binding sites by gel filtration. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1988, 337, 64-8. | 3.0 | 55 |
| 26 | Improving Potency, Selectivity, and Water Solubility of Adenosine A1 Receptor Antagonists: Xanthines Modified at Position 3 and Related Pyrimido[1,2,3-cd]purinediones. <i>ChemMedChem</i> , 2006, 1, 891-902. | 3.2 | 54 |
| 27 | Pharmacological characterization of novel adenosine ligands in recombinant and native human A2B receptors. <i>Biochemical Pharmacology</i> , 2005, 70, 1601-1612. | 4.4 | 53 |
| 28 | Cytoskeletal regulation of chemotactic receptors: Molecular complexation of N-formyl peptide receptors with G proteins and actin. <i>European Journal of Haematology</i> , 1993, 51, 288-293. | 2.2 | 53 |
| 29 | 8-Bromo-9-alkyl adenine derivatives as tools for developing new adenosine A2A and A2B receptors ligands. <i>Bioorganic and Medicinal Chemistry</i> , 2009, 17, 2812-2822. | 3.0 | 53 |
| 30 | Carbamazepine distinguishes between adenosine receptors that mediate different second messenger responses. <i>European Journal of Pharmacology</i> , 1991, 206, 285-290. | 2.6 | 52 |
| 31 | Comparison of A1 adenosine receptors in brain from different species by radioligand binding and photoaffinity labelling. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1991, 343, 196-201. | 3.0 | 52 |
| 32 | Agonist-Induced Internalization and Recycling of the Human A3 Adenosine Receptors. <i>Journal of Neurochemistry</i> , 2002, 75, 1493-1501. | 3.9 | 52 |
| 33 | Characterization of the Solubilized A1Adenosine Receptor from Rat Brain Membranes. <i>Journal of Neurochemistry</i> , 1986, 46, 1528-1534. | 3.9 | 51 |
| 34 | A New Synthesis of Sulfonamides by Aminolysis of p-Nitrophenylsulfonates Yielding Potent and Selective Adenosine A2BReceptor Antagonists. <i>Journal of Medicinal Chemistry</i> , 2006, 49, 4384-4391. | 6.4 | 50 |
| 35 | Adenosine receptor agonists: synthesis and biological evaluation of 1-deaza analogs of adenosine derivatives. <i>Journal of Medicinal Chemistry</i> , 1988, 31, 1179-1183. | 6.4 | 49 |
| 36 | 7-Deazaadenines Bearing Polar Substituents: Structure-Activity Relationships of New A1 and A3 Adenosine Receptor Antagonists. <i>Journal of Medicinal Chemistry</i> , 2000, 43, 4636-4646. | 6.4 | 49 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | 2-(Benzimidazol-2-yl)quinoxalines: A Novel Class of Selective Antagonists at Human A1 and A3 Adenosine Receptors Designed by 3D Database Searching. <i>Journal of Medicinal Chemistry</i> , 2005, 48, 8253-8260. | 6.4 | 49 |
| 38 | 2-Substituted N-ethylcarboxamidoadenosine derivatives as high-affinity agonists at human A3 adenosine receptors. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1999, 360, 103-108. | 3.0 | 48 |
| 39 | 8-(Sulfostyryl)xanthines: water-soluble A2A-selective adenosine receptor antagonists Preliminary results were presented at the International Symposium "Purines" in Milan, Italy; abstract published in <i>Drug Dev. Res.</i> 1996, 37, 112.. <i>Bioorganic and Medicinal Chemistry</i> , 1998, 6, 707-719. | 3.0 | 46 |
| 40 | Discovery of novel A3 adenosine receptor ligands based on chromone scaffold. <i>Biochemical Pharmacology</i> , 2012, 84, 21-29. | 4.4 | 46 |
| 41 | N6-Cycloalkyl- and N6-Bicycloalkyl-C5'-(C2')-modified Adenosine Derivatives as High-Affinity and Selective Agonists at the Human A1 Adenosine Receptor with Antinociceptive Effects in Mice. <i>Journal of Medicinal Chemistry</i> , 2009, 52, 2393-2406. | 6.4 | 44 |
| 42 | Chalcone-based derivatives as new scaffolds for A3 adenosine receptor antagonists. <i>Journal of Pharmacy and Pharmacology</i> , 2013, 65, 697-703. | 2.4 | 44 |
| 43 | Reduction of postischemic leukocyte-endothelium interaction by adenosine via A2 receptor. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1992, 346, 234-237. | 3.0 | 43 |
| 44 | 2',3'-Dideoxy-N6-cyclohexyladenosine: an adenosine derivative with antagonist properties at adenosine receptors. <i>European Journal of Pharmacology</i> , 1988, 156, 157-160. | 3.5 | 40 |
| 45 | Effector coupling of stably transfected human A3 adenosine receptors in CHO cells. <i>Biochemical Pharmacology</i> , 2002, 64, 61-65. | 4.4 | 40 |
| 46 | The Significance of 2-Furyl Ring Substitution with a 2-(<i>para</i> -substituted) Aryl Group in a New Series of Pyrazolo-triazolo-pyrimidines as Potent and Highly Selective hA ₃ Adenosine Receptors Antagonists: New Insights into Structure-Affinity Relationship and Receptor-Antagonist Recognition. <i>Journal of Medicinal Chemistry</i> , 2010, 53, 3361-3375. | 6.4 | 40 |
| 47 | Medicinal Chemistry and Therapeutic Potential of Agonists, Antagonists and Allosteric Modulators of A1 Adenosine Receptor: Current Status and Perspectives. <i>Current Pharmaceutical Design</i> , 2019, 25, 2697-2715. | 1.9 | 39 |
| 48 | Involvement of mitogen protein kinase cascade in agonist-mediated human A3 adenosine receptor regulation. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2002, 1591, 55-62. | 4.1 | 38 |
| 49 | N6-Methoxy-2-alkynyladenosine Derivatives as Highly Potent and Selective Ligands at the Human A3 Adenosine Receptor. <i>Journal of Medicinal Chemistry</i> , 2007, 50, 1222-1230. | 6.4 | 38 |
| 50 | Characterization of potent ligands at human recombinant adenosine receptors. <i>Drug Development Research</i> , 1998, 45, 176-181. | 2.9 | 34 |
| 51 | Synthesis, Biological Evaluation, and Molecular Modeling of Ribose-Modified Adenosine Analogues as Adenosine Receptor Agonists. <i>Journal of Medicinal Chemistry</i> , 2005, 48, 1550-1562. | 6.4 | 34 |
| 52 | Synthesis and Biological Evaluation of 2-Alkynyl-N6-methyl-5'-N-methylcarboxamidoadenosine Derivatives as Potent and Highly Selective Agonists for the Human Adenosine A3 Receptor. <i>Journal of Medicinal Chemistry</i> , 2009, 52, 7897-7900. | 6.4 | 34 |
| 53 | 9-Ethyladenine derivatives as adenosine receptor antagonists: 2- and 8-substitution results in distinct selectivities. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2003, 367, 629-634. | 3.0 | 33 |
| 54 | Synthesis, adenosine receptor binding and molecular modelling studies of novel thieno[2,3-d]pyrimidine derivatives. <i>Chemical Biology and Drug Design</i> , 2018, 91, 962-969. | 3.2 | 33 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Adenosine A _{2A} Receptor Antagonists: New 8-Substituted 9-Ethyladenines as Tools for inâ€¦vivo Rat Models of Parkinson's Disease. ChemMedChem, 2009, 4, 1010-1019. | 3.2 | 32 |
| 56 | Guanine Nucleotide Effects on 8-Cyclopentyl-1,3-[3H]Dipropylxanthine Binding to Membrane-Bound and Solubilized A1 Adenosine Receptors of Rat Brain. Journal of Neurochemistry, 1990, 54, 1988-1994. | 3.9 | 30 |
| 57 | Synthesis and Biological Evaluation of a New Series of 1,2,4-Triazolo[1,5- <i>a</i>]-1,3,5-triazines as Human A _{2A} Adenosine Receptor Antagonists with Improved Water Solubility. Journal of Medicinal Chemistry, 2011, 54, 877-889. | 6.4 | 30 |
| 58 | New insight into adenosine receptors selectivity derived from a novel series of [5-substituted-4-phenyl-1,3-thiazol-2-yl] benzamides and furamides. European Journal of Medicinal Chemistry, 2013, 63, 924-934. | 5.5 | 30 |
| 59 | The glycoprotein nature of A1 adenosine receptors. Biochemical and Biophysical Research Communications, 1986, 140, 406-413. | 2.1 | 29 |
| 60 | The application of a 3D-QSAR (autoMEP/PLS) approach as an efficient pharmacodynamic-driven filtering method for small-sized virtual library: Application to a lead optimization of a human A3 adenosine receptor antagonist. Bioorganic and Medicinal Chemistry, 2006, 14, 4923-4932. | 3.0 | 29 |
| 61 | Synthesis and pharmacological characterization of a new series of 5,7-disubstituted-[1,2,4]triazolo[1,5- <i>a</i>][1,3,5]triazine derivatives as adenosine receptor antagonists: A preliminary inspection of ligandâ€“receptor recognition process. Bioorganic and Medicinal Chemistry, 2010, 18, 2524-2536. | 3.0 | 29 |
| 62 | Affinities of barbiturates for the GABA-receptor complex and A1 adenosine receptors: a possible explanation of their excitatory effects. Naunyn-Schmiedeberg's Archives of Pharmacology, 1987, 336, 211-217. | 3.0 | 27 |
| 63 | A carboxyl-terminal tail peptide of neutrophil chemotactic receptor disrupts its physical complex with G protein. Journal of Leukocyte Biology, 1993, 54, 572-577. | 3.3 | 26 |
| 64 | Combining selectivity and affinity predictions using an integrated Support Vector Machine (SVM) approach: An alternative tool to discriminate between the human adenosine A2A and A3 receptor pyrazolo-triazolo-pyrimidine antagonists binding sites. Bioorganic and Medicinal Chemistry, 2009, 17, 5259-5274. | 3.0 | 26 |
| 65 | 7-Amino-2-aryl/hetero-aryl-5,8-dihydro[1,2,4]triazolo[1,5- <i>a</i>]pyridine-6-carbonitriles: Synthesis and adenosine receptor binding studies. Chemical Biology and Drug Design, 2019, 94, 1568-1573. | 3.2 | 26 |
| 66 | Binding thermodynamics at the human A3 adenosine receptor. Biochemical Pharmacology, 2002, 63, 157-161. | 4.4 | 25 |
| 67 | Novel thiazoleâ€“thiophene conjugates as adenosine receptor antagonists: Synthesis, biological evaluation and docking studies. Bioorganic and Medicinal Chemistry Letters, 2015, 25, 1306-1309. | 2.2 | 25 |
| 68 | Regulation of p42/p44 mitogen-activated protein kinase by the human adenosine A3 receptor in transfected CHO cells. European Journal of Pharmacology, 2001, 420, 19-26. | 3.5 | 24 |
| 69 | 5- ² -Carbamoyl derivatives of 2- ² -C-methyl-purine nucleosides as selective A1 adenosine receptor agonists: Affinity, efficacy, and selectivity for A1 receptor from different species. Bioorganic and Medicinal Chemistry, 2008, 16, 336-353. | 3.0 | 24 |
| 70 | Synthesis and adenosine receptors binding studies of new fluorinated analogues of pyrido[2,3- <i>d</i>]pyrimidines and quinazolines. Medicinal Chemistry Research, 2018, 27, 756-767. | 2.4 | 24 |
| 71 | Neuroprotective potential of adenosine A1 receptor partial agonists in experimental models of cerebral ischemia. Journal of Neurochemistry, 2019, 149, 211-230. | 3.9 | 24 |
| 72 | Discovery of 7-(Prolinol-N-yl)-2-phenylamino-thiazolo[5,4- <i>d</i>]pyrimidines as Novel Non-Nucleoside Partial Agonists for the A2A Adenosine Receptor: Prediction from Molecular Modeling. Journal of Medicinal Chemistry, 2016, 59, 5922-5928. | 6.4 | 23 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Discovery of indolylpiperazinympyrimidines with dual-target profiles at adenosine A2A and dopamine D2 receptors for Parkinson's disease treatment. <i>PLoS ONE</i> , 2018, 13, e0188212. | 2.5 | 23 |
| 74 | A2A-selective adenosine receptor antagonists: Development of water-soluble prodrugs and a new tritiated radioligand. <i>Drug Development Research</i> , 1998, 45, 190-197. | 2.9 | 22 |
| 75 | Novel chiral isoxazole derivatives: Synthesis and pharmacological characterization at human β_2 -adrenergic receptor subtypes. <i>Bioorganic and Medicinal Chemistry</i> , 2007, 15, 2533-2543. | 3.0 | 22 |
| 76 | Cytotoxic purine nucleoside analogues bind to A1, A2A, and A3 adenosine receptors. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2012, 385, 519-525. | 3.0 | 22 |
| 77 | 5- ϵ -Ethyl-tetrazolyl-N ⁶ -Substituted Adenosine and 2-Chloro-adenosine Derivatives as Highly Potent Dual Acting A ₁ Adenosine Receptor Agonists and A ₃ Adenosine Receptor Antagonists. <i>Journal of Medicinal Chemistry</i> , 2015, 58, 2560-2566. | 6.4 | 22 |
| 78 | Discovery of Novel Adenosine Receptor Antagonists through a Combined Structure- and Ligand-Based Approach Followed by Molecular Dynamics Investigation of Ligand Binding Mode. <i>Journal of Chemical Information and Modeling</i> , 2018, 58, 794-815. | 5.4 | 22 |
| 79 | Pyrazolo[4,3-e]1,2,4-triazolo[1,5-c]pyrimidine derivatives as adenosine receptor ligands: A starting point for searching A2B adenosine receptor antagonists. <i>Drug Development Research</i> , 2001, 53, 225-235. | 2.9 | 21 |
| 80 | A1 adenosine receptor antagonists, 3-aryl[1,2,4]triazino[4,3-a]benzimidazol-4-(10H)-ones (ATBIs) and N-alkyl and N-acyl-(7-substituted-2-phenylimidazo[1,2-a][1,3,5]triazin-4-yl)amines (ITAs): Different recognition of bovine and human binding sites. <i>Drug Development Research</i> , 2004, 63, 1-7. | 2.9 | 20 |
| 81 | Exploring Potency and Selectivity Receptor Antagonist Profiles Using a Multilabel Classification Approach: The Human Adenosine Receptors as a Key Study. <i>Journal of Chemical Information and Modeling</i> , 2009, 49, 2820-2836. | 5.4 | 19 |
| 82 | Different efficacy of adenosine and NECA derivatives at the human A3 adenosine receptor: Insight into the receptor activation switch. <i>Biochemical Pharmacology</i> , 2014, 87, 321-331. | 4.4 | 19 |
| 83 | Guanosine exerts antiplatelet and antithrombotic properties through an adenosine-related cAMP-PKA signaling. <i>International Journal of Cardiology</i> , 2017, 248, 294-300. | 1.7 | 19 |
| 84 | 7-Nitrobenzofurazan (NBD) derivatives of 5- ϵ -N-ethylcarboxamidoadenosine (NECA) as new fluorescent probes for human A3 adenosine receptors. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2001, 11, 3023-3026. | 2.2 | 18 |
| 85 | 5,7-Disubstituted-[1,2,4]triazolo[1,5-a][1,3,5]triazines as pharmacological tools to explore the antagonist selectivity profiles toward adenosine receptors. <i>European Journal of Medicinal Chemistry</i> , 2016, 108, 529-541. | 5.5 | 18 |
| 86 | Exploring the Directionality of 5-Substitutions in a New Series of 5-Alkylaminopyrazolo[4,3-e]1,2,4-triazolo[1,5-c]pyrimidine as a Strategy To Design Novel Human A ₃ Adenosine Receptor Antagonists.. <i>Journal of Medicinal Chemistry</i> , 2012, 55, 9654-9668. | 6.4 | 17 |
| 87 | Effects of Adenosine on Histamine Release from Human Lung Fragments. <i>International Archives of Allergy and Immunology</i> , 1992, 98, 50-56. | 2.1 | 16 |
| 88 | Synthesis and adenosine receptors binding affinities of a series of 3-arylcoumarins. <i>Journal of Pharmacy and Pharmacology</i> , 2013, 65, 1590-1597. | 2.4 | 16 |
| 89 | Revisiting a Receptor-Based Pharmacophore Hypothesis for Human A _{2A} Adenosine Receptor Antagonists. <i>Journal of Chemical Information and Modeling</i> , 2013, 53, 1620-1637. | 5.4 | 16 |
| 90 | Discovery of simplified N2-substituted pyrazolo[3,4-d]pyrimidine derivatives as novel adenosine receptor antagonists: Efficient synthetic approaches, biological evaluations and molecular docking studies. <i>Bioorganic and Medicinal Chemistry</i> , 2014, 22, 1751-1765. | 3.0 | 16 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Pharmacological characterisation of novel adenosine A ₃ receptor antagonists. Scientific Reports, 2020, 10, 20781. | 3.3 | 16 |
| 92 | Peptide G protein agonists from a phage display library. Biochemical Pharmacology, 2003, 65, 961-967. | 4.4 | 15 |
| 93 | Synthesis of enantiopure 1 ² -isoxazoline derivatives and evaluation of their affinity and efficacy profiles at human 1 ² -adrenergic receptor subtypes. Bioorganic and Medicinal Chemistry, 2006, 14, 4393-4401. | 3.0 | 15 |
| 94 | Organoruthenium Antagonists of Human A ₃ Adenosine Receptors. Chemistry - A European Journal, 2013, 19, 8321-8330. | 3.3 | 15 |
| 95 | Exploring the Role of N ⁶ -Substituents in Potent Dual Acting 5 ² -C-Ethyltetrazolyladenosine Derivatives: Synthesis, Binding, Functional Assays, and Antinociceptive Effects in Mice. Journal of Medicinal Chemistry, 2017, 60, 4327-4341. | 6.4 | 15 |
| 96 | Synthesis and Biological Activity of Trisubstituted Adenines as A _{2A} Adenosine Receptor Antagonists. Nucleosides, Nucleotides and Nucleic Acids, 2007, 26, 1443-1446. | 1.1 | 14 |
| 97 | 8-(2-Furyl)adenine derivatives as A _{2A} adenosine receptor ligands. European Journal of Medicinal Chemistry, 2013, 70, 525-535. | 5.5 | 14 |
| 98 | Structure-Based Optimization of Coumarin hA ₃ Adenosine Receptor Antagonists. Journal of Medicinal Chemistry, 2020, 63, 2577-2587. | 6.4 | 14 |
| 99 | Targeting adenosine receptors with coumarins: synthesis and binding activities of amide and carbamate derivatives. Journal of Pharmacy and Pharmacology, 2012, 65, 30-34. | 2.4 | 13 |
| 100 | Insight into the Interactions between Novel Coumarin Derivatives and Human A ₃ Adenosine Receptors. ChemMedChem, 2014, 9, 2245-2253. | 3.2 | 13 |
| 101 | Scaffold Decoration at Positions 5 and 8 of 1,2,4-Triazolo[1,5-c]Pyrimidines to Explore the Antagonist Profiling on Adenosine Receptors: A Preliminary Structure-Activity Relationship Study. Journal of Medicinal Chemistry, 2014, 57, 6210-6225. | 6.4 | 13 |
| 102 | The A _{2B} adenosine receptor in MDA-MB-231 breast cancer cells diminishes ERK1/2 phosphorylation by activation of MAPK-phosphatase-1. PLoS ONE, 2018, 13, e0202914. | 2.5 | 13 |
| 103 | [1,2,4]Triazolo[1,5-c]pyrimidines as adenosine receptor antagonists: Modifications at the 8 position to reach selectivity towards A ₃ adenosine receptor subtype. European Journal of Medicinal Chemistry, 2018, 157, 837-851. | 5.5 | 13 |
| 104 | New 2,6,9-trisubstituted adenines as adenosine receptor antagonists: a preliminary SAR profile. Purinergic Signalling, 2007, 3, 339-346. | 2.2 | 12 |
| 105 | The Length and Flexibility of the 2-Substituent of 9-Ethyladenine Derivatives Modulate Affinity and Selectivity for the Human A _{2A} Adenosine Receptor. ChemMedChem, 2016, 11, 1829-1839. | 3.2 | 12 |
| 106 | 2- and 8-alkynyl-9-ethyladenines: Synthesis and biological activity at human and rat adenosine receptors. Purinergic Signalling, 2005, 1, 173-181. | 2.2 | 11 |
| 107 | Does the combination of optimal substitutions at the C2-, N5- and N8-positions of the pyrazolo-triazolo-pyrimidine scaffold guarantee selective modulation of the human A ₃ adenosine receptors?. Bioorganic and Medicinal Chemistry, 2011, 19, 6120-6134. | 3.0 | 11 |
| 108 | New potent and selective A ₁ adenosine receptor antagonists as potential tools for the treatment of gastrointestinal diseases. European Journal of Medicinal Chemistry, 2018, 151, 199-213. | 5.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | A novel small molecule A2A adenosine receptor agonist, indirubin-3- β -monoxime, alleviates lipid-induced inflammation and insulin resistance in 3T3-L1 adipocytes. <i>Biochemical Journal</i> , 2019, 476, 2371-2391. | 3.7 | 11 |
| 110 | Adenosine receptors and human melanoma. <i>Drug Development Research</i> , 2003, 58, 377-385. | 2.9 | 10 |
| 111 | Pyrazolo-triazolo-pyrimidines as adenosine receptor antagonists: Effect of the N-5 bond type on the affinity and selectivity at the four adenosine receptor subtypes. <i>Purinergic Signalling</i> , 2008, 4, 39-46. | 2.2 | 10 |
| 112 | Receptor crosstalk: haloperidol treatment enhances A2A adenosine receptor functioning in a transfected cell model. <i>Purinergic Signalling</i> , 2010, 6, 373-381. | 2.2 | 10 |
| 113 | 2-(N-Acyl) and 2-N-acyl-N6-substituted analogues of adenosine and their affinity at the human adenosine receptors. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2004, 14, 1495-1498. | 2.2 | 9 |
| 114 | Novel Mutants of the Human A_{2A} -Adrenergic Receptor Reveal Amino Acids Relevant for Receptor Activation. <i>Journal of Biological Chemistry</i> , 2006, 281, 18120-18125. | 3.4 | 9 |
| 115 | Development of novel adenosine receptor ligands based on the 3-amidocoumarin scaffold. <i>Bioorganic Chemistry</i> , 2015, 61, 1-6. | 4.1 | 9 |
| 116 | Synthesis and Evaluation of a New Series of 8- β -(2-Nitroaryl)Xanthines as Adenosine Receptor Ligands. <i>Drug Development Research</i> , 2016, 77, 241-250. | 2.9 | 9 |
| 117 | Structure-Based Design, Synthesis, and In Vivo Antinociceptive Effects of Selective $\text{A}_{1/2}$ Adenosine Receptor Agonists. <i>Journal of Medicinal Chemistry</i> , 2018, 61, 305-318. | 6.4 | 9 |
| 118 | Pyrazolo[4,3- <i>e</i>][1,2,4]triazolo[1,5- <i>c</i>]pyrimidines to develop functionalized ligands to target adenosine receptors: fluorescent ligands as an example. <i>MedChemComm</i> , 2019, 10, 1094-1108. | 3.4 | 9 |
| 119 | Radiation inactivation analysis of the A_1 adenosine receptor of rat brain Decrease in radiation inactivation size in the presence of guanine nucleotide. <i>FEBS Letters</i> , 1989, 252, 125-128. | 2.8 | 8 |
| 120 | Synthesis of (R)-3,4-dihydro-2H-pyran-2-carboxaldehyde: application to the synthesis of potent adenosine A_{2A} and A_3 receptor agonist. <i>Tetrahedron Letters</i> , 2009, 50, 2693-2696. | 1.4 | 8 |
| 121 | New 9-methyl-8-(4-hydroxyphenyl)adenine derivatives as A_1 adenosine receptor antagonists. <i>Collection of Czechoslovak Chemical Communications</i> , 2011, 76, 1379-1393. | 1.0 | 8 |
| 122 | Synthesis and pharmacological characterization of novel xanthine carboxylate amides as A_{2A} adenosine receptor ligands exhibiting bronchospasmolytic activity. <i>Bioorganic Chemistry</i> , 2016, 65, 26-37. | 4.1 | 8 |
| 123 | Adenosine Receptor Ligands: Coumarin-Chalcone Hybrids as Modulating Agents on the Activity of hARs. <i>Molecules</i> , 2020, 25, 4306. | 3.8 | 8 |
| 124 | Synthesis of 2-Azido-(R)-N ⁶ -p-hydroxyphenylisopropyladenosine (R-AHPIA) as Potential Photoaffinity Probe for $\text{A}_{1/2}$ Adenosine Receptors. <i>Nucleosides & Nucleotides</i> , 1986, 5, 213-222. | 0.5 | 7 |
| 125 | G-protein-coupled A_1 adenosine receptors in coated vesicles of mammalian brain: Characterization by radioligand binding and photoaffinity labelling. <i>Cellular Signalling</i> , 1992, 4, 737-745. | 3.6 | 7 |
| 126 | The interaction of N-formyl peptide chemoattractant receptors with the membrane skeleton is energy-dependent. <i>Cellular Signalling</i> , 1994, 6, 943-947. | 3.6 | 7 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 127 | Physical coupling of N-formyl peptide chemoattractant receptors to G protein is unaffected by desensitization. <i>Biochemical Pharmacology</i> , 1994, 48, 1297-1300. | 4.4 | 7 |
| 128 | Synthesis, biological studies and molecular modeling investigation of 1,3-dimethyl-2,4-dioxo-6-methyl-8-(substituted) 1,2,3,4-tetrahydro [1,2,4]-triazolo [3,4-f]-purines as potential adenosine receptor antagonists. <i>Il Farmaco</i> , 2005, 60, 299-306. | 0.9 | 7 |
| 129 | Synthesis, biological evaluation and molecular modelling studies of 1,3,7,8-tetrasubstituted xanthines as potent and selective A2A AR ligands with in vivo efficacy against animal model of Parkinson's disease. <i>Bioorganic Chemistry</i> , 2019, 87, 601-612. | 4.1 | 7 |
| 130 | Molecular modelling study of 2-phenylethynyladenosine (PEAdo) derivatives as highly selective A3 adenosine receptor ligands. <i>Purinergic Signalling</i> , 2006, 2, 589-594. | 2.2 | 6 |
| 131 | Inhibitors of membranous adenylyl cyclases with affinity for adenosine receptors. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2016, 389, 349-352. | 3.0 | 6 |
| 132 | The Influence of the 1-(3-Trifluoromethyl-Benzyl)-1H-Pyrazole-4-yl Moiety on the Adenosine Receptors Affinity Profile of Pyrazolo[4,3-e][1,2,4]Triazolo[1,5-c]Pyrimidine Derivatives. <i>PLoS ONE</i> , 2015, 10, e0143504. | 2.5 | 6 |
| 133 | Coumarins and adenosine receptors: New perceptions in structure-affinity relationships. <i>Chemical Biology and Drug Design</i> , 2018, 91, 245-256. | 3.2 | 5 |
| 134 | A2A Adenosine Receptor Antagonists: Are Triazotriazine and Purine Scaffolds Interchangeable?. <i>Molecules</i> , 2022, 27, 2386. | 3.8 | 5 |
| 135 | Discovery of 2-aminoimidazole and 2-amino imidazolyl-thiazoles as non-xanthine human adenosine A3receptor antagonists: SAR and molecular modeling studies. <i>MedChemComm</i> , 2018, 9, 676-684. | 3.4 | 4 |
| 136 | New A2A adenosine receptor antagonists: a structure-based upside-down interaction in the receptor cavity. <i>Bioorganic Chemistry</i> , 2019, 92, 103183. | 4.1 | 4 |
| 137 | Targeting G Protein-Coupled Receptors with Magnetic Carbon Nanotubes: The Case of the A3 Adenosine Receptor. <i>ChemMedChem</i> , 2020, 15, 1909-1920. | 3.2 | 4 |
| 138 | Bronchospasmolytic activity and adenosine receptor binding of some newer 1,3-dipropyl-8-phenyl substituted xanthine derivatives. <i>Chemical Biology and Drug Design</i> , 2020, 95, 600-609. | 3.2 | 4 |
| 139 | A3 adenosine receptors: Synthesis and biological evaluation of new potent and selective ligands. , 2005, , . | | 4 |
| 140 | Agonists activate different A2B adenosine receptor signaling pathways in MBA-MD-231 breast cancer cells with distinct potencies. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 2019, 392, 1515-1521. | 3.0 | 3 |
| 141 | Adenosine Receptor Biology in the Central Nervous System. , 2017, , 3-22. | | 2 |
| 142 | Pharmacology and Molecular Biology of A3 Adenosine Receptors. , 2010, , 49-58. | | 1 |
| 143 | Synthesis, Biological Studies and Molecular Modeling Investigation of 1,3-Dimethyl-2,4-dioxo-6-methyl-8-(substituted) 1,2,3,4-Tetrahydro-[1,2,4]-triazolo-[3,4-f]-purines as Potential Adenosine Receptor Antagonists.. <i>ChemInform</i> , 2005, 36, no. | 0.0 | 0 |
| 144 | Design, Synthesis and Evaluation of New Indolylpyrimidylpiperazines for Gastrointestinal Cancer Therapy. <i>Molecules</i> , 2019, 24, 3661. | 3.8 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Potent and selective A ₃ adenosine receptor antagonists bearing aminoesters as heterobifunctional moieties. RSC Medicinal Chemistry, 2021, 12, 254-262. | 3.9 | 0 |