

Hitoshi Ashida

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8886355/hitoshi-ashida-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

175
papers

5,110
citations

37
h-index

64
g-index

178
ext. papers

5,641
ext. citations

3.8
avg, IF

5.62
L-index

#	Paper	IF	Citations
175	Simultaneous determination of all polyphenols in vegetables, fruits, and teas. <i>Journal of Agricultural and Food Chemistry</i> , 2003 , 51, 571-81	5.7	490
174	Multitargeted cancer prevention by quercetin. <i>Cancer Letters</i> , 2008 , 269, 315-25	9.9	489
173	Cyanidin 3-glucoside ameliorates hyperglycemia and insulin sensitivity due to downregulation of retinol binding protein 4 expression in diabetic mice. <i>Biochemical Pharmacology</i> , 2007 , 74, 1619-27	6	225
172	Tea catechin suppresses adipocyte differentiation accompanied by down-regulation of PPARgamma2 and C/EBPalpha in 3T3-L1 cells. <i>Bioscience, Biotechnology and Biochemistry</i> , 2004 , 68, 2353-9	2.1	212
171	Flavones and flavonols at dietary levels inhibit a transformation of aryl hydrocarbon receptor induced by dioxin. <i>FEBS Letters</i> , 2000 , 476, 213-7	3.8	135
170	Rapid preparation of a plasma membrane fraction from adipocytes and muscle cells: application to detection of translocated glucose transporter 4 on the plasma membrane. <i>Bioscience, Biotechnology and Biochemistry</i> , 2007 , 71, 2343-6	2.1	128
169	Black soybean seed coat extract ameliorates hyperglycemia and insulin sensitivity via the activation of AMP-activated protein kinase in diabetic mice. <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 5558-64	5.7	99
168	Epigallocatechin gallate promotes GLUT4 translocation in skeletal muscle. <i>Biochemical and Biophysical Research Communications</i> , 2008 , 377, 286-90	3.4	95
167	Prevention mechanisms of glucose intolerance and obesity by cacao liquor procyanidin extract in high-fat diet-fed C57BL/6 mice. <i>Archives of Biochemistry and Biophysics</i> , 2012 , 527, 95-104	4.1	93
166	A black soybean seed coat extract prevents obesity and glucose intolerance by up-regulating uncoupling proteins and down-regulating inflammatory cytokines in high-fat diet-fed mice. <i>Journal of Agricultural and Food Chemistry</i> , 2011 , 59, 8985-93	5.7	87
165	Anti-obesity actions of green tea: possible involvements in modulation of the glucose uptake system and suppression of the adipogenesis-related transcription factors. <i>BioFactors</i> , 2004 , 22, 135-40	6.1	81
164	D-pinitol and myo-inositol stimulate translocation of glucose transporter 4 in skeletal muscle of C57BL/6 mice. <i>Bioscience, Biotechnology and Biochemistry</i> , 2010 , 74, 1062-7	2.1	77
163	Rutin potentiates insulin receptor kinase to enhance insulin-dependent glucose transporter 4 translocation. <i>Molecular Nutrition and Food Research</i> , 2014 , 58, 1168-76	5.9	69
162	Dietary flavonoids as cancer-preventive and therapeutic biofactors. <i>Frontiers in Bioscience - Scholar</i> , 2011 , 3, 1332-62	2.4	64
161	β-Glucan from <i>Lentinus edodes</i> inhibits nitric oxide and tumor necrosis factor-α production and phosphorylation of mitogen-activated protein kinases in lipopolysaccharide-stimulated murine RAW 264.7 macrophages. <i>Journal of Biological Chemistry</i> , 2012 , 287, 871-8	5.4	58
160	Rat L6 myotubes as an in vitro model system to study GLUT4-dependent glucose uptake stimulated by inositol derivatives. <i>Cytotechnology</i> , 2007 , 55, 103-8	2.2	58
159	Oolong, black and pu-erh tea suppresses adiposity in mice via activation of AMP-activated protein kinase. <i>Food and Function</i> , 2014 , 5, 2420-9	6.1	57

158	Green and black tea suppress hyperglycemia and insulin resistance by retaining the expression of glucose transporter 4 in muscle of high-fat diet-fed C57BL/6J mice. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 12916-23	5.7	55
157	Antimutagenicity of flavones and flavonols to heterocyclic amines by specific and strong inhibition of the cytochrome P450 1A family. <i>Bioscience, Biotechnology and Biochemistry</i> , 1998 , 62, 970-7	2.1	55
156	Target enzymes on hepatic dysfunction caused by dietary products of lipid peroxidation. <i>Archives of Biochemistry and Biophysics</i> , 1991 , 288, 71-8	4.1	53
155	Measurement of Glucose Uptake in Cultured Cells. <i>Current Protocols in Pharmacology</i> , 2015 , 71, 12.14.1-12.14.26	4.1	50
154	4-Hydroxyderricin and xanthoangelol from Ashitaba (<i>Angelica keiskei</i>) suppress differentiation of preadipocytes to adipocytes via AMPK and MAPK pathways. <i>Molecular Nutrition and Food Research</i> , 2013 , 57, 1729-40	5.9	49
153	Immunomodulatory beta-glucan from <i>Lentinus edodes</i> activates mitogen-activated protein kinases and nuclear factor-kappaB in murine RAW 264.7 macrophages. <i>Journal of Biological Chemistry</i> , 2011 , 286, 31194-8	5.4	47
152	Cinnamtannin A2, a tetrameric procyanidin, increases GLP-1 and insulin secretion in mice. <i>Bioscience, Biotechnology and Biochemistry</i> , 2013 , 77, 888-91	2.1	46
151	Activation of the aryl hydrocarbon receptor induces hepatic steatosis via the upregulation of fatty acid transport. <i>Archives of Biochemistry and Biophysics</i> , 2010 , 504, 221-7	4.1	45
150	Curcumin suppresses the transformation of an aryl hydrocarbon receptor through its phosphorylation. <i>Archives of Biochemistry and Biophysics</i> , 2007 , 466, 267-73	4.1	45
149	Interaction between the aryl hydrocarbon receptor and its antagonists, flavonoids. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 359, 822-7	3.4	45
148	Prenylated chalcones 4-hydroxyderricin and xanthoangelol stimulate glucose uptake in skeletal muscle cells by inducing GLUT4 translocation. <i>Molecular Nutrition and Food Research</i> , 2011 , 55, 467-75	5.9	42
147	Procyanidin Promotes Translocation of Glucose Transporter 4 in Muscle of Mice through Activation of Insulin and AMPK Signaling Pathways. <i>PLoS ONE</i> , 2016 , 11, e0161704	3.7	42
146	Suppression mechanisms of flavonoids on aryl hydrocarbon receptor-mediated signal transduction. <i>Archives of Biochemistry and Biophysics</i> , 2010 , 501, 134-41	4.1	41
145	Pigments in green tea leaves (<i>Camellia sinensis</i>) suppress transformation of the aryl hydrocarbon receptor induced by dioxin. <i>Journal of Agricultural and Food Chemistry</i> , 2004 , 52, 2499-506	5.7	41
144	Propolis extract promotes translocation of glucose transporter 4 and glucose uptake through both PI3K- and AMPK-dependent pathways in skeletal muscle. <i>BioFactors</i> , 2013 , 39, 457-66	6.1	40
143	A novel method using 8-hydroperoxy-2'-deoxyguanosine formation for evaluating antioxidative potency. <i>Free Radical Research</i> , 2002 , 36, 307-16	4	39
142	Chalcones suppress fatty acid-induced lipid accumulation through a LKB1/AMPK signaling pathway in HepG2 cells. <i>Food and Function</i> , 2014 , 5, 1134-41	6.1	38
141	Fermented tea improves glucose intolerance in mice by enhancing translocation of glucose transporter 4 in skeletal muscle. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 11366-71	5.7	38

140	2,3,7,8-tetrachlorodibenzo-p-dioxin impairs an insulin signaling pathway through the induction of tumor necrosis factor-alpha in adipocytes. <i>Toxicological Sciences</i> , 2010 , 115, 482-91	4.4	38
139	TCDD-induced CYP1A1 expression, an index of dioxin toxicity, is suppressed by flavonoids permeating the human intestinal Caco-2 cell monolayers. <i>Journal of Agricultural and Food Chemistry</i> , 2006 , 54, 8891-8	5.7	38
138	Luteolin modulates expression of drug-metabolizing enzymes through the AhR and Nrf2 pathways in hepatic cells. <i>Archives of Biochemistry and Biophysics</i> , 2014 , 557, 36-46	4.1	36
137	Tea catechins modulate the glucose transport system in 3T3-L1 adipocytes. <i>Food and Function</i> , 2010 , 1, 167-73	6.1	36
136	An update on the dietary ligands of the AhR. <i>Expert Opinion on Drug Metabolism and Toxicology</i> , 2008 , 4, 1429-47	5.5	36
135	Cacao liquor procyanidin extract improves glucose tolerance by enhancing GLUT4 translocation and glucose uptake in skeletal muscle. <i>Journal of Nutritional Science</i> , 2012 , 1, e2	2.7	35
134	Inhibitory effects of caffeine and its metabolites on intracellular lipid accumulation in murine 3T3-L1 adipocytes. <i>BioFactors</i> , 2008 , 34, 293-302	6.1	35
133	Quercetin and its metabolite isorhamnetin promote glucose uptake through different signalling pathways in myotubes. <i>Scientific Reports</i> , 2019 , 9, 2690	4.9	34
132	Comparative analysis of carbohydrate-binding specificities of two anti-glycogen monoclonal antibodies using ELISA and surface plasmon resonance. <i>Carbohydrate Research</i> , 2012 , 350, 49-54	2.9	33
131	β-Glucan from <i>Saccharomyces cerevisiae</i> reduces lipopolysaccharide-induced inflammatory responses in RAW264.7 macrophages. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2012 , 1820, 1656-63	4.3	31
130	Molokhia (<i>Corchorus olitorius</i> L.) extract suppresses transformation of the aryl hydrocarbon receptor induced by dioxins. <i>Food and Chemical Toxicology</i> , 2006 , 44, 250-60	4.7	31
129	Suppressive effects of flavonoids on dioxin toxicity. <i>BioFactors</i> , 2000 , 12, 201-6	6.1	31
128	Green Tea Ameliorates Hyperglycemia by Promoting the Translocation of Glucose Transporter 4 in the Skeletal Muscle of Diabetic Rodents. <i>International Journal of Molecular Sciences</i> , 2019 , 20,	6.3	30
127	Antagonistic and agonistic effects of indigoids on the transformation of an aryl hydrocarbon receptor. <i>Archives of Biochemistry and Biophysics</i> , 2008 , 470, 187-99	4.1	30
126	Suppressive effect of polysaccharides from the edible and medicinal mushrooms, <i>Lentinus edodes</i> and <i>Agaricus blazei</i> , on the expression of cytochrome P450s in mice. <i>Bioscience, Biotechnology and Biochemistry</i> , 2002 , 66, 1610-4	2.1	30
125	Ameliorative effects of polyunsaturated fatty acids against palmitic acid-induced insulin resistance in L6 skeletal muscle cells. <i>Lipids in Health and Disease</i> , 2012 , 11, 36	4.4	29
124	Ashitaba (<i>Angelica keiskei</i>) extract prevents adiposity in high-fat diet-fed C57BL/6 mice. <i>Food and Function</i> , 2015 , 6, 135-45	6.1	28
123	Inhibitory effects of 4-hydroxyderricin and xanthoangelol on lipopolysaccharide-induced inflammatory responses in RAW264 macrophages. <i>Journal of Agricultural and Food Chemistry</i> , 2014 , 62, 462-7	5.7	26

122	Suppression of lipopolysaccharide and galactosamine-induced hepatic inflammation by red grape pomace. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 9315-20	5.7	26
121	Subcellular localization of flavonol aglycone in hepatocytes visualized by confocal laser scanning fluorescence microscope. <i>Cytotechnology</i> , 2009 , 59, 177-82	2.2	26
120	Licorice flavonoid oil enhances muscle mass in KK-A mice. <i>Life Sciences</i> , 2018 , 205, 91-96	6.8	26
119	Epigallocatechin gallate induces GLUT4 translocation in skeletal muscle through both PI3K- and AMPK-dependent pathways. <i>Food and Function</i> , 2018 , 9, 4223-4233	6.1	25
118	Absorption and metabolism of 4-hydroxyderricin and xanthoangelol after oral administration of <i>Angelica keiskei</i> (Ashitaba) extract in mice. <i>Archives of Biochemistry and Biophysics</i> , 2012 , 521, 71-6	4.1	25
117	Cardamonin stimulates glucose uptake through translocation of glucose transporter-4 in L6 myotubes. <i>Phytotherapy Research</i> , 2011 , 25, 1218-24	6.7	25
116	An enzymatic fluorimetric assay to quantitate 2-deoxyglucose and 2-deoxyglucose-6-phosphate for in vitro and in vivo use. <i>Analytical Biochemistry</i> , 2010 , 404, 238-40	3.1	25
115	2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD)-induced changes in activities of nuclear protein kinases and phosphatases affecting DNA binding activity of c-Myc and AP-1 in the livers of guinea pigs. <i>Biochemical Pharmacology</i> , 2000 , 59, 741-51	6	25
114	Black soybean seed coat polyphenols prevent B(a)P-induced DNA damage through modulating drug-metabolizing enzymes in HepG2 cells and ICR mice. <i>Mutation Research - Genetic Toxicology and Environmental Mutagenesis</i> , 2013 , 752, 34-41	3	24
113	Green tea prevents obesity by increasing expression of insulin-like growth factor binding protein-1 in adipose tissue of high-fat diet-fed mice. <i>Journal of Agricultural and Food Chemistry</i> , 2012 , 60, 8917-23	5.7	24
112	Measurement of glucose uptake in cultured cells. <i>Current Protocols in Pharmacology</i> , 2011 , Chapter 12, Unit 12.14.1-22	4.1	24
111	Effect of in vivo administered 2,3,7,8-tetrachlorodibenzo-p-dioxin on DNA-binding activities of nuclear transcription factors in liver of guinea pigs. <i>Journal of Biochemical and Molecular Toxicology</i> , 1998 , 12, 191-204	3.4	24
110	Enzymatically synthesized glycogen inhibits colitis through decreasing oxidative stress. <i>Free Radical Biology and Medicine</i> , 2017 , 106, 355-367	7.8	22
109	Piperine Promotes Glucose Uptake through ROS-Dependent Activation of the CAMKK/AMPK Signaling Pathway in Skeletal Muscle. <i>Molecular Nutrition and Food Research</i> , 2018 , 62, e1800086	5.9	22
108	Suppressive effects of ethanolic extracts from propolis and its main botanical origin on dioxin toxicity. <i>Journal of Agricultural and Food Chemistry</i> , 2005 , 53, 10306-9	5.7	22
107	A physiological concentration of luteolin induces phase II drug-metabolizing enzymes through the ERK1/2 signaling pathway in HepG2 cells. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 663, 151-159	4.1	21
106	Screening plant derived dietary phenolic compounds for bioactivity related to cardiovascular disease. <i>Phytotherapy Research</i> , 2018 , 32, 22-28	3.2	20
105	Glabridin induces glucose uptake via the AMP-activated protein kinase pathway in muscle cells. <i>Molecular and Cellular Endocrinology</i> , 2014 , 393, 99-108	4.4	20

104	3-O-Acyl-epicatechins Increase Glucose Uptake Activity and GLUT4 Translocation through Activation of PI3K Signaling in Skeletal Muscle Cells. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 16288-99	6.3	20
103	A new southwestern chemistry-based ELISA for detection of aryl hydrocarbon receptor transformation: application to the screening of its receptor agonists and antagonists. <i>Journal of Immunological Methods</i> , 2004 , 287, 187-201	2.5	19
102	Black tea theaflavins suppress dioxin-induced transformation of the aryl hydrocarbon receptor. <i>Bioscience, Biotechnology and Biochemistry</i> , 2005 , 69, 883-90	2.1	18
101	Metabolites of galangin by 2,3,7,8-tetrachlorodibenzo-p-dioxin-inducible cytochrome P450 1A1 in human intestinal epithelial Caco-2 cells and their antagonistic activity toward aryl hydrocarbon receptor. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 8111-8	5.7	17
100	Rapid Preparation of a Plasma Membrane Fraction: Western Blot Detection of Translocated Glucose Transporter 4 from Plasma Membrane of Muscle and Adipose Cells and Tissues. <i>Current Protocols in Protein Science</i> , 2016 , 85, 29.18.1-29.18.12	3.1	17
99	The chalcones cardamonin and flavokawain B inhibit the differentiation of preadipocytes to adipocytes by activating ERK. <i>Archives of Biochemistry and Biophysics</i> , 2014 , 554, 44-54	4.1	16
98	High-throughput evaluation of aryl hydrocarbon receptor-binding sites selected via chromatin immunoprecipitation-based screening in Hepa-1c1c7 cells stimulated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Genes and Genetic Systems</i> , 2008 , 83, 455-68	1.4	16
97	Theobromine suppresses adipogenesis through enhancement of CCAAT-enhancer-binding protein degradation by adenosine receptor A1. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017 , 1864, 2438-2448	4.9	15
96	Immunomodulatory activity of enzymatically synthesized glycogen and its digested metabolite in a co-culture system consisting of differentiated Caco-2 cells and RAW264.7 macrophages. <i>Food and Function</i> , 2013 , 4, 1387-93	6.1	15
95	Detection of orally administered inositol stereoisomers in mouse blood plasma and their effects on translocation of glucose transporter 4 in skeletal muscle cells. <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 4850-4	5.7	15
94	Absorption, metabolism, distribution and faecal excretion of B-type procyanidin oligomers in mice after a single oral administration of black soybean seed coat extract. <i>Food and Function</i> , 2018 , 9, 5362-5370	6.1	15
93	Glabridin inhibits dexamethasone-induced muscle atrophy. <i>Archives of Biochemistry and Biophysics</i> , 2019 , 664, 157-166	4.1	14
92	The interaction of auraptene and other oxyprenylated phenylpropanoids with glucose transporter type 4. <i>Phytomedicine</i> , 2017 , 32, 74-79	6.5	14
91	Metabolic fate of orally administered enzymatically synthesized glycogen in rats. <i>Food and Function</i> , 2011 , 2, 183-9	6.1	14
90	Involvement of SREBPs in 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced disruption of lipid metabolism in male guinea pig. <i>Toxicology and Applied Pharmacology</i> , 2008 , 229, 281-9	4.6	14
89	Tryptophan pyrolysis products, Trp-P-1 and Trp-P-2 induce apoptosis in primary cultured rat hepatocytes. <i>Bioscience, Biotechnology and Biochemistry</i> , 1998 , 62, 2283-7	2.1	14
88	Effects of Microbial Metabolites of (-)-Epigallocatechin Gallate on Glucose Uptake in L6 Skeletal Muscle Cell and Glucose Tolerance in ICR Mice. <i>Biological and Pharmaceutical Bulletin</i> , 2019 , 42, 212-221	2.3	13
87	Black soybean seed coat polyphenols prevent AAPH-induced oxidative DNA-damage in HepG2 cells. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2017 , 60, 108-114	3.1	13

86	Enzymatically synthesized glycogen reduces lipid accumulation in diet-induced obese rats. <i>Nutrition Research</i> , 2013 , 33, 743-52	4	13
85	Oral toxicological studies of black soybean (<i>Glycine max</i>) hull extract: acute studies in rats and mice, and chronic studies in mice. <i>Food and Chemical Toxicology</i> , 2011 , 49, 3272-8	4.7	13
84	Aryl hydrocarbon receptor-mediated induction of the cytosolic phospholipase A(2)alpha gene by 2,3,7,8-tetrachlorodibenzo-p-dioxin in mouse hepatoma Hepa-1c1c7 cells. <i>Journal of Bioscience and Bioengineering</i> , 2009 , 108, 277-81	3.3	13
83	Inhibition of P-glycoprotein enhances the suppressive effect of kaempferol on transformation of the aryl hydrocarbon receptor. <i>Bioscience, Biotechnology and Biochemistry</i> , 2009 , 73, 1635-9	2.1	13
82	Suppression of cytochrome P450 1A1 expression induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin in mouse hepatoma hepa-1c1c7 cells treated with serum of (-)-epigallocatechin-3-gallate- and green tea extract-administered rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 2009 , 73, 1206-8	2.1	13
81	Cacao polyphenol extract suppresses transformation of an aryl hydrocarbon receptor in C57BL/6 mice. <i>Journal of Agricultural and Food Chemistry</i> , 2008 , 56, 10399-405	5.7	13
80	Enzymatically modified isoquercitrin promotes energy metabolism through activating AMPK in male C57BL/6 mice. <i>Food and Function</i> , 2019 , 10, 5188-5202	6.1	12
79	Structure-activity relationships of anthraquinones on the suppression of DNA-binding activity of the aryl hydrocarbon receptor induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Journal of Bioscience and Bioengineering</i> , 2009 , 107, 296-300	3.3	12
78	Suppression of dioxin mediated aryl hydrocarbon receptor transformation by ethanolic extracts of propolis. <i>Bioscience, Biotechnology and Biochemistry</i> , 2004 , 68, 935-8	2.1	12
77	Cacao liquor procyanidins prevent postprandial hyperglycaemia by increasing glucagon-like peptide-1 activity and AMP-activated protein kinase in mice. <i>Journal of Nutritional Science</i> , 2019 , 8, e2	2.7	11
76	Anthocyanins fail to suppress transformation of aryl hydrocarbon receptor induced by dioxin. <i>Bioscience, Biotechnology and Biochemistry</i> , 2005 , 69, 896-903	2.1	11
75	Substitution at the C-3 Position of Catechins Has an Influence on the Binding Affinities against Serum Albumin. <i>Molecules</i> , 2017 , 22,	4.8	10
74	Black Tea Polyphenols Promotes GLUT4 Translocation through Both PI3K-and AMPK-dependent Pathways in Skeletal Muscle Cells. <i>Food Science and Technology Research</i> , 2015 , 21, 489-494	0.8	10
73	Catechins in tea suppress the activity of cytochrome P450 1A1 through the aryl hydrocarbon receptor activation pathway in rat livers. <i>International Journal of Food Sciences and Nutrition</i> , 2015 , 66, 300-7	3.7	10
72	3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1) induces apoptosis in rat splenocytes and thymocytes by different mechanisms. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2000 , 457, 57-67	3.3	10
71	Low dose of luteolin activates Nrf2 in the liver of mice at start of the active phase but not that of the inactive phase. <i>PLoS ONE</i> , 2020 , 15, e0231403	3.7	10
70	Curcumin and its derivatives inhibit 2,3,7,8,-tetrachloro-dibenzo-p-dioxin-induced expression of drug metabolizing enzymes through aryl hydrocarbon receptor-mediated pathway. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 616-628	2.1	10
69	Black soybean improves the vascular function through an increase in nitric oxide and a decrease in oxidative stress in healthy women. <i>Archives of Biochemistry and Biophysics</i> , 2020 , 688, 108408	4.1	9

68	Protective action of dehydroascorbic acid on the Ah receptor-dependent and receptor-independent induction of lipid peroxidation in adipose tissue of male guinea pig caused by TCDD administration. <i>Journal of Biochemical Toxicology</i> , 1996 , 11, 269-78		9
67	Prevention effect of quercetin and its glycosides on obesity and hyperglycemia through activating AMPK in high-fat diet-fed ICR mice. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 74-83	3.1	9
66	An analysis method for flavan-3-ols using high performance liquid chromatography coupled with a fluorescence detector. <i>Journal of Food and Drug Analysis</i> , 2017 , 25, 478-487	7	8
65	6-(Methylsulfinyl)hexyl isothiocyanate protects acetaldehyde-caused cytotoxicity through the induction of aldehyde dehydrogenase in hepatocytes. <i>Archives of Biochemistry and Biophysics</i> , 2020 , 686, 108329	4.1	8
64	Evaluation of Intestinal Dioxin Permeability Using Human Caco-2 Cell Monolayers. <i>Food Science and Technology Research</i> , 2003 , 9, 364-366	0.8	8
63	Black soybean seed coat polyphenols promote nitric oxide production in the aorta through glucagon-like peptide-1 secretion from the intestinal cells. <i>Food and Function</i> , 2019 , 10, 7875-7882	6.1	8
62	Liquorice flavonoid oil suppresses hyperglycaemia accompanied by skeletal muscle myocellular GLUT4 recruitment to the plasma membrane in KK-A mice. <i>International Journal of Food Sciences and Nutrition</i> , 2019 , 70, 294-302	3.7	8
61	Lycii fructus extract ameliorates hydrogen peroxide-induced cytotoxicity through indirect antioxidant action. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 1812-1820	2.1	8
60	Methylxanthine Derivative-Rich Cacao Extract Suppresses Differentiation of Adipocytes through Downregulation of PPAR and C/EBPs. <i>Journal of Nutritional Science and Vitaminology</i> , 2018 , 64, 151-160	1.1	7
59	Preventive Effects of Black Soybean Seed Coat Polyphenols against DNA Damage in Salmonella typhimurium. <i>Food Science and Technology Research</i> , 2013 , 19, 685-690	0.8	7
58	Concentration of Catechins and Caffeine in Black Tea Affects Suppression of Fat Accumulation and Hyperglycemia in High-fat Diet-fed Mice. <i>Food Science and Technology Research</i> , 2011 , 17, 353-359	0.8	7
57	The heterocyclic amine, 3-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole induces apoptosis in cocultures of rat parenchymal and nonparenchymal liver cells. <i>Toxicology and Applied Pharmacology</i> , 2001 , 177, 59-67	4.6	7
56	The mechanisms of ameliorating effect of a green tea polyphenol on diabetic nephropathy based on diacylglycerol kinase. <i>Scientific Reports</i> , 2020 , 10, 11790	4.9	7
55	Adenosine isolated from Grifola gargal promotes glucose uptake via PI3K and AMPK signalling pathways in skeletal muscle cells. <i>Journal of Functional Foods</i> , 2017 , 33, 268-277	5.1	6
54	Caffeine-Stimulated Intestinal Epithelial Cells Suppress Lipid Accumulation in Adipocytes. <i>Journal of Nutritional Science and Vitaminology</i> , 2017 , 63, 331-338	1.1	6
53	Theophylline suppresses interleukin-6 expression by inhibiting glucocorticoid receptor signaling in pre-adipocytes. <i>Archives of Biochemistry and Biophysics</i> , 2018 , 646, 98-106	4.1	6
52	Effects of biosurfactants on assays of PCB congeners in transgenic arabidopsis plants carrying a recombinant guinea pig AhR-mediated GUS reporter gene expression system. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2010 , 45, 773-9	2.2	6
51	Effects of dietary lipid peroxidation products on hormonal responses in primary cultured hepatocytes of rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 1997 , 61, 2089-94	2.1	6

50	Decreased expression of Bcl-x protein during hepatocarcinogenesis induced exogenously and endogenously in rats. <i>Japanese Journal of Cancer Research</i> , 2001 , 92, 1270-7		6
49	3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1) induces caspase-dependent apoptosis in mononuclear cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2001 , 1539, 44-57	4.9	6
48	Insights into the potential benefits of black soybean (L.) polyphenols in lifestyle diseases. <i>Food and Function</i> , 2020 , 11, 7321-7339	6.1	6
47	Black Soybean Improves Vascular Function and Blood Pressure: A Randomized, Placebo Controlled, Crossover Trial in Humans. <i>Nutrients</i> , 2020 , 12,	6.7	6
46	4-Hydroxyderricin and xanthoangelol isolated from <i>Angelica keiskei</i> prevent dexamethasone-induced muscle loss. <i>Food and Function</i> , 2020 , 11, 5498-5512	6.1	5
45	Amelioration of diabetic nephropathy by oral administration of d- α -tocopherol and its mechanisms. <i>Bioscience, Biotechnology and Biochemistry</i> , 2018 , 82, 65-73	2.1	5
44	Luteolin suppresses TCDD-induced wasting syndrome in a cultured adipocyte model. <i>Pesticide Biochemistry and Physiology</i> , 2015 , 120, 14-20	4.9	5
43	Suppressive Effects of Flavonoids on Activation of the Aryl Hydrocarbon Receptor Induced by Dioxins. <i>ACS Symposium Series</i> , 2008 , 369-374	0.4	5
42	3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1) triggers apoptosis by DNA double-strand breaks caused by inhibition of topoisomerase I. <i>Carcinogenesis</i> , 2004 , 25, 1149-55	4.6	5
41	Black tea extract suppresses transformation of aryl hydrocarbon receptor induced by dioxin. <i>BioFactors</i> , 2004 , 21, 367-9	6.1	5
40	Detection of biomarkers for apoptosis in rat liver after perfusion with 3-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1). <i>Bioscience, Biotechnology and Biochemistry</i> , 2000 , 64, 2021-4	2.1	5
39	Hepatic Phosphoglucosyltransferase Activity as a Marker of Oxidative Stress Induced by Pro-oxidative Drugs. <i>Bioscience, Biotechnology and Biochemistry</i> , 1994 , 58, 55-9	2.1	5
38	Enzymatically synthesized glycogen protects inflammation induced by urban particulate matter in normal human epidermal keratinocytes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 29-35	3.1	5
37	Kaempferol modulates TCDD- and t-BHQ-induced drug-metabolizing enzymes and luteolin enhances this effect. <i>Food and Function</i> , 2020 , 11, 3668-3680	6.1	5
36	Cacao polyphenols regulate the circadian clock gene expression and through glucagon-like peptide-1 secretion. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 53-60	3.1	4
35	The Ashitaba () Chalcones 4-hydroxyderricin and Xanthoangelol Suppress Melanomagenesis By Targeting BRAF and PI3K. <i>Cancer Prevention Research</i> , 2018 , 11, 607-620	3.2	4
34	Application of lipid extracts from <i>Solidago canadensis</i> to phytomonitoring of PCB126 in transgenic <i>Arabidopsis</i> plants. <i>Science of the Total Environment</i> , 2014 , 491-492, 240-5	10.2	4
33	Recombinant human AhR-mediated GUS reporter gene assays for PCB congeners in transgenic tobacco plants in comparison with recombinant mouse and guinea pig AhRs. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2010 , 45, 711-8	2.2	4

32	3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole induces apoptosis and necrosis with activation of different caspases in rat splenocytes. <i>Bioscience, Biotechnology and Biochemistry</i> , 2004 , 68, 964-7	2.1	4
31	Apoptosis in the thymus after intraperitoneal injection of rats with Trp-P-1. <i>Environmental and Molecular Mutagenesis</i> , 2002 , 40, 175-83	3.2	4
30	Xenobiotic tolerance of primary cultured hepatocytes in rats fed a high-fat or high-protein diet. <i>Journal of Nutritional Science and Vitaminology</i> , 1998 , 44, 89-102	1.1	4
29	Bisacurone suppresses hepatic lipid accumulation through inhibiting lipogenesis and promoting lipolysis. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 43-52	3.1	4
28	The cacao procyanidin extract-caused anti-hyperglycemic effect was changed by the administration timings. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 61-66	3.1	4
27	A Frequent Drinking of Green Tea Lowers the Levels of Endogenous Oxidative Stress in Small Intestines, Erythrocytes and Kidneys in Rats. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2006 , 39, 32-39	3.1	4
26	Kaempferol Promotes Glucose Uptake in Myotubes through a JAK2-Dependent Pathway. <i>Journal of Agricultural and Food Chemistry</i> , 2020 , 68, 13720-13729	5.7	4
25	Daily consumption of black soybean (<i>Glycine max</i> L.) seed coat polyphenols attenuates dyslipidemia in apolipoprotein E-deficient mice. <i>Journal of Functional Foods</i> , 2020 , 72, 104054	5.1	3
24	Assays of polychlorinated biphenyl congeners and co-contaminated heavy metals in the transgenic Arabidopsis plants carrying the recombinant guinea pig aryl hydrocarbon receptor-mediated Eglucuronidase reporter gene expression system. <i>Journal of Environmental Science and Health - Part B Pesticides, Food Contaminants, and Agricultural Wastes</i> , 2012 , 47, 925-32	2.2	3
23	Green Tea Extracts Prevent the Dioxin Toxicity through the Suppression of Transformation of the Aryl Hydrocarbon Receptor. <i>ACS Symposium Series</i> , 2003 , 119-127	0.4	3
22	Suppression of Cytochrome P4501 a Subfamily in Mouse Liver by Oral Intake of Polysaccharides from Mushrooms, <i>Lentinus edodes</i> and <i>Agaricus blazei</i> . <i>ACS Symposium Series</i> , 2003 , 235-248	0.4	3
21	Up-regulation of CD13/aminopeptidase N induced by phorbol ester is involved in redox regulation and tumor necrosis factor alpha production in HL-60 cells. <i>Inflammation</i> , 2002 , 26, 175-81	5.1	3
20	Evoking cytochrome P450 1A activity interferes with apoptosis induced by 3-amino-1,4-dimethyl-5H-pyrido [4,3-b]indole (Trp-P-1) in rat hepatocytes under the ex vivo system. <i>Bioscience, Biotechnology and Biochemistry</i> , 2002 , 66, 356-62	2.1	3
19	Screening of indigenous plants from Japan for modulating effects on transformation of the aryl hydrocarbon receptor. <i>Asian Pacific Journal of Cancer Prevention</i> , 2006 , 7, 208-20	1.7	3
18	Antagonistic effect of the Ainu-selected traditional beneficial plants on the transformation of an aryl hydrocarbon receptor. <i>Journal of Food Science</i> , 2012 , 77, C420-9	3.4	2
17	Heterocyclic Amines Induce Apoptosis in Hepatocytes. <i>ACS Symposium Series</i> , 1998 , 88-100	0.4	2
16	Enzymatically synthesized glycogen prevents ultraviolet B-induced cell damage in normal human epidermal keratinocytes. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 36-42	3.1	2
15	Enzymatically synthesized glycogen inhibited degranulation and inflammatory responses through stimulation of intestine. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2020 , 67, 67-73	3.1	2

14	Modulation of Drug-Metabolizing Enzymes and Transporters by Polyphenols as an Anticarcinogenic Effect 2014 , 1127-1135		1
13	Preventive Effects of Food Components on Caspase-8-Mediated Apoptosis Induced by Dietary Carcinogen, Trp-P-1, in Rat Mononuclear Cells. <i>ACS Symposium Series</i> , 2003 , 128-140	0.4	1
12	Tea Extracts Modulate a Glucose Transport System in 3T3-L1 Adipocytes. <i>ACS Symposium Series</i> , 2003 , 224-234	0.4	1
11	3-amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (trp-P-1) is incorporated into rat splenocytes, thymocytes, and hepatocytes through monoamine transporters and induces apoptosis. <i>Bioscience, Biotechnology and Biochemistry</i> , 2002 , 66, 1205-12	2.1	1
10	Theobromine enhances the conversion of white adipocytes into beige adipocytes in a PPAR α activation-dependent manner. <i>Journal of Nutritional Biochemistry</i> , 2021 , 100, 108898	6.3	1
9	5-Aminolevulinic acid combined with ferrous iron improves glucose tolerance in high-fat diet-fed mice via upregulation of glucose transporter 1. <i>Experimental and Therapeutic Medicine</i> , 2021 , 22, 1454	2.1	0
8	Prevention Mechanism of Lifestyle-related Diseases by Dietary Polyphenols. <i>Nihon Eiyōshokuryō Gakkai Shi = Nippon Eiyōshokuryō Gakkaishi = Journal of Japanese Society of Nutrition and Food Science</i> , 2017 , 70, 213-223	0.2	
7	Effect of Green Tea Extract on Copper Dynamics in Mouse Hair. <i>Food Science and Technology Research</i> , 2013 , 19, 123-125	0.8	
6	A Tryptophan Pyrolysis Product, 3-Amino-1,4-dimethyl-5H-pyrido[4,3-b]indole (Trp-P-1) but Not Its Metabolite Induces Apoptosis in Primary Cultured Rat Hepatocytes. <i>ACS Symposium Series</i> , 2003 , 141-151 ⁹⁻⁴		
5	Insulin-Mimetic Activity of Inositol Derivatives Depends on Phosphorylation of PKC ζ in L6 Myotubes 2010 , 327-331		
4	Low dose of luteolin activates Nrf2 in the liver of mice at start of the active phase but not that of the inactive phase 2020 , 15, e0231403		
3	Low dose of luteolin activates Nrf2 in the liver of mice at start of the active phase but not that of the inactive phase 2020 , 15, e0231403		
2	Low dose of luteolin activates Nrf2 in the liver of mice at start of the active phase but not that of the inactive phase 2020 , 15, e0231403		
1	Low dose of luteolin activates Nrf2 in the liver of mice at start of the active phase but not that of the inactive phase 2020 , 15, e0231403		