Yayoi Inomata

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/8890211/publications.pdf

Version: 2024-02-01

28	766	15	27
papers	citations	h-index	g-index
35	35	35	789
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Current situation of atmospheric nanoparticles in Fukue Island, Japan. Tellus, Series B: Chemical and Physical Meteorology, 2022, 70, 1498688.	1.6	4
2	Different Transport Behaviors between Asian Dust and Polycyclic Aromatic Hydrocarbons in Urban Areas: Monitoring in Fukuoka and Kanazawa, Japan. Applied Sciences (Switzerland), 2022, 12, 5404.	2,5	3
3	Transboundary air pollution reduction rapidly reflected in stream water chemistry in forested catchment on the sea of Japan coast in central Japan. Atmospheric Environment, 2021, 248, 118223.	4.1	11
4	Comparison of three aerosol representations of NHM-Chem (v1.0) for the simulations of air quality and climate-relevant variables. Geoscientific Model Development, 2021, 14, 2235-2264.	3.6	16
5	Particulate PAH Transport Associated with Adult Chronic Cough Occurrence Closely Connected with Meteorological Conditions: A Modelling Study. Atmosphere, 2021, 12, 1163.	2.3	1
6	Calculating source contributions to urban atmospheric polycyclic aromatic hydrocarbons and nitropolycyclic aromatic hydrocarbons using 1-nitropyrene and pyrene: An application to an Asian dust event. Chemosphere, 2021, 280, 130662.	8.2	6
7	Mass balance and latest fluxes of radiocesium derived from the fukushima accident in the western North Pacific Ocean and coastal regions of Japan. Journal of Environmental Radioactivity, 2020, 217, 106206.	1.7	32
8	Estimating transboundary transported anthropogenic sulfate deposition in Japan using the sulfur isotopic ratio. Science of the Total Environment, 2019, 691, 779-788.	8.0	7
9	NHM-Chem, the Japan Meteorological Agency's Regional Meteorology – Chemistry Model: Model Evaluations toward the Consistent Predictions of the Chemical, Physical, and Optical Properties of Aerosols. Journal of the Meteorological Society of Japan, 2019, 97, 337-374.	1.8	37
10	Fukushima radionuclides in the marine environment from coastal region of Japan to the Pacific Ocean through the end of 2016. Progress in Nuclear Science and Technology, 2019, 6, 1-7.	0.3	9
11	Estimate of Fukushima-derived radiocaesium in the North Pacific Ocean in summer 2012. Journal of Radioanalytical and Nuclear Chemistry, 2018, 318, 1587-1596.	1.5	15
12	Transport of FNPP1-derived radiocaesium from subtropical mode water in the western North Pacific Ocean to the Sea of Japan. Ocean Science, 2018, 14, 813-826.	3.4	28
13	Recirculation of FNPP1-derived radiocaesium observed in winter 2015/2016 in coastal regions of Japan. Applied Radiation and Isotopes, 2017, 126, 83-87.	1.5	26
14	Synergy between air pollution and urban meteorological changes through aerosol-radiation-diffusion feedback―A case study of Beijing in January 2013. Atmospheric Environment, 2017, 171, 98-110.	4.1	15
15	Sourceâ€"Receptor Relationship Analysis of the Atmospheric Deposition of PAHs Subject to Long-Range Transport in Northeast Asia. Environmental Science & Environmental Scienc	10.0	11
16	Transboundary transport of anthropogenic sulfur in PM2.5 at a coastal site in the Sea of Japan as studied by sulfur isotopic ratio measurement. Science of the Total Environment, 2016, 553, 617-625.	8.0	21
17	134Cs and 137Cs in the North Pacific Ocean derived from the March 2011 TEPCO Fukushima Dai-ichi Nuclear Power Plant accident, Japan. Part two: estimation of 134Cs and 137Cs inventories in the North Pacific Ocean. Journal of Oceanography, 2016, 72, 67-76.	1.7	138
18	Long-term variation of the source of sulfate deposition in a leeward area of Asian continent in view of sulfur isotopic composition. Atmospheric Environment, 2016, 140, 42-51.	4.1	24

#	Article	IF	CITATIONS
19	Spatial and temporal distributions of ¹³⁴ Cs and ¹³⁷ Cs derived from the TEPCO Fukushima Daiichi Nuclear Power Plant accident in the North Pacific Ocean by using optimal interpolation analysis. Environmental Sciences: Processes and Impacts, 2016, 18, 126-136.	3.5	37
20	Source–receptor relationships of nitrate in Northeast Asia and influence of sea salt on the long-range transport of nitrate. Atmospheric Environment, 2013, 79, 67-78.	4.1	36
21	Source contribution analysis of surface particulate polycyclic aromatic hydrocarbon concentrations in northeastern Asia by source–receptor relationships. Environmental Pollution, 2013, 182, 324-334.	7. 5	13
22	Modeling wet deposition and concentration of inorganics over Northeast Asia with MRI-PM/c. Geoscientific Model Development, 2012, 5, 1363-1375.	3.6	18
23	Development of the RAQM2 aerosol chemical transport model and predictions of the Northeast Asian aerosol mass, size, chemistry, and mixing type. Atmospheric Chemistry and Physics, 2012, 12, 11833-11856.	4.9	55
24	Optimum interpolation analysis of basin-scale 137Cs transport in surface seawater in the North Pacific Ocean. Journal of Environmental Monitoring, 2012, 14, 3146.	2.1	5
25	Emission and Atmospheric Transport of Particulate PAHs in Northeast Asia. Environmental Science & Technology, 2012, 46, 4941-4949.	10.0	99
26	Influence of blocking effect of mountain and local front on two Asian-dust events observed at Mt. Haruna and Tsukuba in Kanto, Japan, in 2007. Atmospheric Environment, 2011, 45, 4429-4441.	4.1	4
27	Dry and wet deposition of water-insoluble dust and water-soluble chemical species during spring 2007 in Tsukuba, Japan. Atmospheric Environment, 2009, 43, 4503-4512.	4.1	36
28	Analysis of 50-y record of surface ¹³⁷ Cs concentrations in the global ocean using the HAM-global database. Journal of Environmental Monitoring, 2009, 11, 116-125.	2.1	53