

# Ginette Serrero

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8907692/publications.pdf>

Version: 2024-02-01

53  
papers

2,055  
citations

279701

23  
h-index

265120

42  
g-index

53  
all docs

53  
docs citations

53  
times ranked

1529  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Resveratrol, a natural product derived from grape, exhibits antiestrogenic activity and inhibits the growth of human breast cancer cells. , 1999, 179, 297-304.  |     | 280       |
| 2  | [6] The growth of cells in serum-free hormone-supplemented media. <i>Methods in Enzymology</i> , 1979, 58, 94-109.   | 0.4 | 279       |
| 3  | PC cell-derived growth factor (PCDGF/GP88, progranulin) stimulates migration, invasiveness and VEGF expression in breast cancer cells. <i>Carcinogenesis</i> , 2004, 25, 1587-1592.  | 1.3 | 113       |
| 4  | Stimulation of adipose differentiation related protein (ADRP) expression in adipocyte precursors by long-chain fatty acids. <i>Journal of Cellular Physiology</i> , 2000, 182, 297-302.  | 2.0 | 97        |
| 5  | Isolation of myoblastic, fibro-adipogenic, and fibroblastic clonal cell lines from a common precursor and study of their requirements for growth and differentiation. <i>Experimental Cell Research</i> , 1981, 132, 313-327.            | 1.2 | 80        |
| 6  | PC Cell-Derived Growth Factor Mediates Tamoxifen Resistance and Promotes Tumor Growth of Human Breast Cancer Cells. <i>Cancer Research</i> , 2004, 64, 1737-1743.  | 0.4 | 76        |
| 7  | PC Cell-Derived Growth Factor Expression in Prostatic Intraepithelial Neoplasia and Prostatic Adenocarcinoma. <i>Clinical Cancer Research</i> , 2004, 10, 1333-1337.   | 3.2 | 75        |
| 8  | Expression of PC-cell-derived growth factor in benign and malignant human breast epithelium. <i>Human Pathology</i> , 2003, 34, 1148-1154.   | 1.1 | 71        |
| 9  | Autocrine growth factor revisited: PC-cell-derived growth factor (progranulin), a critical player in breast cancer tumorigenesis. <i>Biochemical and Biophysical Research Communications</i> , 2003, 308, 409-413.                       | 1.0 | 71        |
| 10 | Proepithelin Regulates Prostate Cancer Cell Biology by Promoting Cell Growth, Migration, and Anchorage-Independent Growth. <i>American Journal of Pathology</i> , 2009, 174, 1037-1047.  | 1.9 | 66        |
| 11 | An in vitro model to study adipose differentiation in serum-free medium. <i>Analytical Biochemistry</i> , 1982, 120, 351-359.  | 1.1 | 64        |
| 12 | Increased Serum GP88 (Progranulin) Concentrations in Rheumatoid Arthritis. <i>Inflammation</i> , 2014, 37, 1806-1813.  | 1.7 | 61        |
| 13 | The granulin-epithelin precursor/PC-cell-derived growth factor is a growth factor for epithelial ovarian cancer. <i>Clinical Cancer Research</i> , 2003, 9, 44-51.   | 3.2 | 58        |
| 14 | GP88 (progranulin): a novel tissue and circulating biomarker for non-“small cell lung carcinoma. <i>Human Pathology</i> , 2014, 45, 1893-1899.   | 1.1 | 50        |
| 15 | PC cell-derived growth factor (granulin precursor) expression and action in human multiple myeloma. <i>Clinical Cancer Research</i> , 2003, 9, 2221-8.   | 3.2 | 50        |
| 16 | Progranulin (GP88) tumor tissue expression is associated with increased risk of recurrence in breast cancer patients diagnosed with estrogen receptor positive invasive ductal carcinoma. <i>Breast Cancer Research</i> , 2012, 14, R26. | 2.2 | 47        |
| 17 | Stimulation of PC Cell-Derived Growth Factor (Epithelin/Granulin Precursor) Expression by Estradiol in Human Breast Cancer Cells. <i>Biochemical and Biophysical Research Communications</i> , 1999, 256, 204-207.                       | 1.0 | 46        |
| 18 | GP88 (PC-Cell Derived Growth Factor, progranulin) stimulates proliferation and confers letrozole resistance to aromatase overexpressing breast cancer cells. <i>BMC Cancer</i> , 2011, 11, 231.  | 1.1 | 42        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Proepithelin is an autocrine growth factor for bladder cancer. <i>Carcinogenesis</i> , 2009, 30, 861-868.   | 1.3 | 41        |
| 20 | PC Cellâ€Derived Growth Factor Confers Resistance to Dexamethasone and Promotes Tumorigenesis in Human Multiple Myeloma. <i>Clinical Cancer Research</i> , 2006, 12, 49-56.   | 3.2 | 37        |
| 21 | PC Cellâ€Derived Growth Factor Stimulates Proliferation and Confers Trastuzumab Resistance to Her-2-Overexpressing Breast Cancer Cells. <i>Clinical Cancer Research</i> , 2006, 12, 4192-4199.                            | 3.2 | 35        |
| 22 | Increased Circulating Level of the Survival Factor GP88 (Progranulin) in the Serum of Breast Cancer Patients When Compared to Healthy Subjects. <i>Breast Cancer: Basic and Clinical Research</i> , 2011, 5, BCBCR.S7224. | 0.6 | 34        |
| 23 | Differentiation of newborn rat adipocyte precursors in defined serum-free medium. <i>In Vitro Cellular &amp; Developmental Biology</i> , 1987, 23, 63-66.   | 1.0 | 29        |
| 24 | Association between increased serum GP88 (progranulin) concentrations and prognosis in patients with malignant lymphomas. <i>Clinica Chimica Acta</i> , 2017, 473, 139-146.   | 0.5 | 25        |
| 25 | Tumorigenicity associated with loss of differentiation and of response to insulin in the adipogenic cell line 1246. <i>In Vitro Cellular &amp; Developmental Biology</i> , 1985, 21, 537-540.                             | 1.0 | 23        |
| 26 | Progranulin levels in blood in Alzheimer's disease and mild cognitive impairment. <i>Annals of Clinical and Translational Neurology</i> , 2018, 5, 616-629.   | 1.7 | 23        |
| 27 | Increased cerebrospinal fluid progranulin correlates with interleukin-6 in the acute phase of neuromyelitis optica spectrum disorder. <i>Journal of Neuroimmunology</i> , 2017, 305, 175-181.                             | 1.1 | 21        |
| 28 | Multiple forms of p55PIK, a regulatory subunit of phosphoinositide 3-kinase, are generated by alternative initiation of translation. <i>Biochemical Journal</i> , 1999, 341, 831-837.                                     | 1.7 | 18        |
| 29 | Expression of GP88 (progranulin) in serum of prostate cancer patients is associated with Gleason scores and overall survival. <i>Cancer Management and Research</i> , 2018, Volume 10, 4173-4180.                         | 0.9 | 13        |
| 30 | Progranulin as a predictive factor of response to chemotherapy in advanced biliary tract carcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 1085-1092.  | 1.1 | 12        |
| 31 | Association of Serum Progranulin Levels With Disease Progression, Therapy Response and Survival in Patients With Metastatic Breast Cancer. <i>Clinical Breast Cancer</i> , 2020, 20, 220-227.                             | 1.1 | 10        |
| 32 | Higher levels of progranulin in cerebrospinal fluid of patients with lymphoma and carcinoma with CNS metastasis. <i>Journal of Neuro-Oncology</i> , 2018, 137, 455-462.   | 1.4 | 9         |
| 33 | Expression of GP88 (Progranulin) Protein Is an Independent Prognostic Factor in Prostate Cancer Patients. <i>Cancers</i> , 2019, 11, 2029.  | 1.7 | 9         |
| 34 | Anti-progranulin/GP88 antibody AG01 inhibits triple negative breast cancer cell proliferation and migration. <i>Breast Cancer Research and Treatment</i> , 2021, 186, 637-653.  | 1.1 | 9         |
| 35 | GP88 (Progranulin) Confers Fulvestrant (Faslodex, ICI 182,780) Resistance to Human Breast Cancer Cells. <i>Advances in Breast Cancer Research</i> , 2014, 03, 68-78.  | 0.1 | 9         |
| 36 | Determination of GP88 (progranulin) expression in breast tumor biopsies improves the risk predictive value of the Nottingham Prognostic Index. <i>Diagnostic Pathology</i> , 2016, 11, 71.                                | 0.9 | 8         |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Prognostic Value of Progranulin in Patients with Colorectal Cancer Treated with Curative Resection. <i>Pathology and Oncology Research</i> , 2020, 26, 397-404.  | 0.9 | 7         |
| 38 | Progranulin/GP88, A Complex and Multifaceted Player of Tumor Growth by Direct Action and via the Tumor Microenvironment. <i>Advances in Experimental Medicine and Biology</i> , 2021, 1329, 475-498.   | 0.8 | 7         |
| 39 | Signaling Pathway of GP88 (Progranulin) in Breast Cancer Cells: Upregulation and Phosphorylation of c-myc by GP88/Progranulin in Her2-Overexpressing Breast Cancer Cells. <i>Breast Cancer: Basic and Clinical Research</i> , 2015, 9s2, BCBCR.S29371.   | 0.6 | 5         |
| 40 | Expression of AR-V7 (Androgen Receptor Variant 7) Protein in Granular Cytoplasmic Structures Is an Independent Prognostic Factor in Prostate Cancer Patients. <i>Cancers</i> , 2020, 12, 2639.   | 1.7 | 5         |
| 41 | Identification of Prostaglandin F2 Receptor Negative Regulator (PTGFRN) as an internalizable target in cancer cells for antibody-drug conjugate development. <i>PLoS ONE</i> , 2021, 16, e0246197.   | 1.1 | 5         |
| 42 | Progranulin depletion inhibits proliferation via the transforming growth factor beta/SMAD family member 2 signaling axis in Kasumi-1 cells. <i>Heliyon</i> , 2021, 7, e05849.  | 1.4 | 5         |
| 43 | Insulin but not IGF-I is required for the maintenance of the adipose phenotype in the adipogenic cell line 1246. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 1999, 35, 642-646.  | 0.7 | 4         |
| 44 | Immunohistochemical Detection of Progranulin (PGRN/GP88/GEP) in Tumor Tissues as a Cancer Prognostic Biomarker. <i>Methods in Molecular Biology</i> , 2018, 1806, 107-120.   | 0.4 | 4         |
| 45 | GP88/PGRN Serum Levels Are Associated with Prognosis for Oral Squamous Cell Carcinoma Patients. <i>Biology</i> , 2021, 10, 400.  | 1.3 | 4         |
| 46 | Potential of Theranostic Target Mining in the Development of Novel Diagnostic and Therapeutic Products in Oncology: Progranulin/GP88 as a Therapeutic and Diagnostic Target for Breast and Lung Cancers. <i>Rinsho Byori the Japanese Journal of Clinical Pathology</i> , 2016, 64, 1296-1309. | 0.1 | 4         |
| 47 | A tribute to Dr. Gordon Hisashi Sato (December 24, 1927–March 31, 2017). <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2018, 54, 177-193.  | 0.7 | 3         |
| 48 | Measurement of Circulating Progranulin (PGRN/GP88/GEP) by Enzyme-Linked Immunosorbent Assay and Application in Human Diseases. <i>Methods in Molecular Biology</i> , 2018, 1806, 95-105.   | 0.4 | 3         |
| 49 | Clinicopathological characteristics and outcomes of gastrointestinal stromal tumors with high progranulin expression. <i>PLoS ONE</i> , 2021, 16, e0245153.  | 1.1 | 3         |
| 50 | Combination of GP88 Expression in Tumor Cells and Tumor-Infiltrating Immune Cells Is an Independent Prognostic Factor for Bladder Cancer Patients. <i>Cells</i> , 2021, 10, 1796.  | 1.8 | 3         |
| 51 | Stimulation of adipose differentiation related protein (ADRP) expression in adipocyte precursors by long-chain fatty acids. , 2000, 182, 297.  |     | 1         |
| 52 | Combined miR-486 and GP88 (Progranulin) Serum Levels Are Suggested as Supportive Biomarkers for Therapy Decision in Elderly Prostate Cancer Patients. <i>Life</i> , 2022, 12, 732.   | 1.1 | 1         |
| 53 | Serum GP88 as a predictive biomarker for hepatocellular carcinoma in patients with viral hepatitis C after direct-acting antiviral agents. <i>Annals of Clinical Biochemistry</i> , 2021, 58, 000456322110367.   | 0.8 | 0         |