

Ginette Serrero

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8907692/publications.pdf>

Version: 2024-02-01

53
papers

2,055
citations

279701

23
h-index

265120

42
g-index

53
all docs

53
docs citations

53
times ranked

1529
citing authors

#	ARTICLE	IF	CITATIONS
1	Combined miR-486 and GP88 (Progranulin) Serum Levels Are Suggested as Supportive Biomarkers for Therapy Decision in Elderly Prostate Cancer Patients. <i>Life</i> , 2022, 12, 732.	1.1	1
2	Clinicopathological characteristics and outcomes of gastrointestinal stromal tumors with high progranulin expression. <i>PLoS ONE</i> , 2021, 16, e0245153.	1.1	3
3	Identification of Prostaglandin F2 Receptor Negative Regulator (PTGFRN) as an internalizable target in cancer cells for antibody-drug conjugate development. <i>PLoS ONE</i> , 2021, 16, e0246197.	1.1	5
4	Progranulin depletion inhibits proliferation via the transforming growth factor beta/SMAD family member 2 signaling axis in Kasumi-1 cells. <i>Heliyon</i> , 2021, 7, e05849.	1.4	5
5	Anti-progranulin/GP88 antibody AG01 inhibits triple negative breast cancer cell proliferation and migration. <i>Breast Cancer Research and Treatment</i> , 2021, 186, 637-653.	1.1	9
6	GP88/PGRN Serum Levels Are Associated with Prognosis for Oral Squamous Cell Carcinoma Patients. <i>Biology</i> , 2021, 10, 400.	1.3	4
7	Combination of GP88 Expression in Tumor Cells and Tumor-Infiltrating Immune Cells Is an Independent Prognostic Factor for Bladder Cancer Patients. <i>Cells</i> , 2021, 10, 1796.	1.8	3
8	Serum GP88 as a predictive biomarker for hepatocellular carcinoma in patients with viral hepatitis C after direct-acting antiviral agents. <i>Annals of Clinical Biochemistry</i> , 2021, 58, 000456322110367.	0.8	0
9	Progranulin/GP88, A Complex and Multifaceted Player of Tumor Growth by Direct Action and via the Tumor Microenvironment. <i>Advances in Experimental Medicine and Biology</i> , 2021, 1329, 475-498.	0.8	7
10	Prognostic Value of Progranulin in Patients with Colorectal Cancer Treated with Curative Resection. <i>Pathology and Oncology Research</i> , 2020, 26, 397-404.	0.9	7
11	Association of Serum Progranulin Levels With Disease Progression, Therapy Response and Survival in Patients With Metastatic Breast Cancer. <i>Clinical Breast Cancer</i> , 2020, 20, 220-227.	1.1	10
12	Expression of AR-V7 (Androgen Receptor Variant 7) Protein in Granular Cytoplasmic Structures Is an Independent Prognostic Factor in Prostate Cancer Patients. <i>Cancers</i> , 2020, 12, 2639.	1.7	5
13	Expression of GP88 (Progranulin) Protein Is an Independent Prognostic Factor in Prostate Cancer Patients. <i>Cancers</i> , 2019, 11, 2029.	1.7	9
14	Higher levels of progranulin in cerebrospinal fluid of patients with lymphoma and carcinoma with CNS metastasis. <i>Journal of Neuro-Oncology</i> , 2018, 137, 455-462.	1.4	9
15	Progranulin levels in blood in Alzheimer's disease and mild cognitive impairment. <i>Annals of Clinical and Translational Neurology</i> , 2018, 5, 616-629.	1.7	23
16	A tribute to Dr. Gordon Hisashi Sato (December 24, 1927â€“March 31, 2017). <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2018, 54, 177-193.	0.7	3
17	Expression of GP88 (progranulin) in serum of prostate cancer patients is associated with Gleason scores and overall survival. <i>Cancer Management and Research</i> , 2018, Volume 10, 4173-4180.	0.9	13
18	Measurement of Circulating Progranulin (PGRN/GP88/GEP) by Enzyme-Linked Immunosorbent Assay and Application in Human Diseases. <i>Methods in Molecular Biology</i> , 2018, 1806, 95-105.	0.4	3

#	ARTICLE	IF	CITATIONS
19	Immunohistochemical Detection of Progranulin (PGRN/GP88/GEP) in Tumor Tissues as a Cancer Prognostic Biomarker. <i>Methods in Molecular Biology</i> , 2018, 1806, 107-120.	0.4	4
20	Association between increased serum GP88 (progranulin) concentrations and prognosis in patients with malignant lymphomas. <i>Clinica Chimica Acta</i> , 2017, 473, 139-146.	0.5	25
21	Increased cerebrospinal fluid progranulin correlates with interleukin-6 in the acute phase of neuromyelitis optica spectrum disorder. <i>Journal of Neuroimmunology</i> , 2017, 305, 175-181.	1.1	21
22	Determination of GP88 (progranulin) expression in breast tumor biopsies improves the risk predictive value of the Nottingham Prognostic Index. <i>Diagnostic Pathology</i> , 2016, 11, 71.	0.9	8
23	Progranulin as a predictive factor of response to chemotherapy in advanced biliary tract carcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 78, 1085-1092.	1.1	12
24	Potential of Theranostic Target Mining in the Development of Novel Diagnostic and Therapeutic Products in Oncology: Progranulin/GP88 as a Therapeutic and Diagnostic Target for Breast and Lung Cancers. <i>Rinsho Byori the Japanese Journal of Clinical Pathology</i> , 2016, 64, 1296-1309.	0.1	4
25	Signaling Pathway of GP88 (Progranulin) in Breast Cancer Cells: Upregulation and Phosphorylation of c-myc by GP88/Progranulin in Her2-Overexpressing Breast Cancer Cells. <i>Breast Cancer: Basic and Clinical Research</i> , 2015, 9s2, BCBCR.S29371.	0.6	5
26	Increased Serum GP88 (Progranulin) Concentrations in Rheumatoid Arthritis. <i>Inflammation</i> , 2014, 37, 1806-1813.	1.7	61
27	GP88 (progranulin): a novel tissue and circulating biomarker for non-small cell lung carcinoma. <i>Human Pathology</i> , 2014, 45, 1893-1899.	1.1	50
28	GP88 (Progranulin) Confers Fulvestrant (Faslodex, ICI 182,780) Resistance to Human Breast Cancer Cells. <i>Advances in Breast Cancer Research</i> , 2014, 03, 68-78.	0.1	9
29	Progranulin (GP88) tumor tissue expression is associated with increased risk of recurrence in breast cancer patients diagnosed with estrogen receptor positive invasive ductal carcinoma. <i>Breast Cancer Research</i> , 2012, 14, R26.	2.2	47
30	GP88 (PC-Cell Derived Growth Factor, progranulin) stimulates proliferation and confers letrozole resistance to aromatase overexpressing breast cancer cells. <i>BMC Cancer</i> , 2011, 11, 231.	1.1	42
31	Increased Circulating Level of the Survival Factor GP88 (Progranulin) in the Serum of Breast Cancer Patients When Compared to Healthy Subjects. <i>Breast Cancer: Basic and Clinical Research</i> , 2011, 5, BCBCR.S7224.	0.6	34
32	Proepithelin is an autocrine growth factor for bladder cancer. <i>Carcinogenesis</i> , 2009, 30, 861-868.	1.3	41
33	Proepithelin Regulates Prostate Cancer Cell Biology by Promoting Cell Growth, Migration, and Anchorage-Independent Growth. <i>American Journal of Pathology</i> , 2009, 174, 1037-1047.	1.9	66
34	PC Cell-Derived Growth Factor Confers Resistance to Dexamethasone and Promotes Tumorigenesis in Human Multiple Myeloma. <i>Clinical Cancer Research</i> , 2006, 12, 49-56.	3.2	37
35	PC Cell-Derived Growth Factor Stimulates Proliferation and Confers Trastuzumab Resistance to Her-2-Overexpressing Breast Cancer Cells. <i>Clinical Cancer Research</i> , 2006, 12, 4192-4199.	3.2	35
36	PC Cell-Derived Growth Factor Expression in Prostatic Intraepithelial Neoplasia and Prostatic Adenocarcinoma. <i>Clinical Cancer Research</i> , 2004, 10, 1333-1337.	3.2	75

#	ARTICLE	IF	CITATIONS
37	PC cell-derived growth factor (PCDGF/GP88, progranulin) stimulates migration, invasiveness and VEGF expression in breast cancer cells. <i>Carcinogenesis</i> , 2004, 25, 1587-1592.	1.3	113
38	PC Cell-Derived Growth Factor Mediates Tamoxifen Resistance and Promotes Tumor Growth of Human Breast Cancer Cells. <i>Cancer Research</i> , 2004, 64, 1737-1743.	0.4	76
39	Expression of PC-cell-derived growth factor in benign and malignant human breast epithelium. <i>Human Pathology</i> , 2003, 34, 1148-1154.	1.1	71
40	Autocrine growth factor revisited: PC-cell-derived growth factor (progranulin), a critical player in breast cancer tumorigenesis. <i>Biochemical and Biophysical Research Communications</i> , 2003, 308, 409-413.	1.0	71
41	The granulin-epithelin precursor/PC-cell-derived growth factor is a growth factor for epithelial ovarian cancer. <i>Clinical Cancer Research</i> , 2003, 9, 44-51.	3.2	58
42	PC cell-derived growth factor (granulin precursor) expression and action in human multiple myeloma. <i>Clinical Cancer Research</i> , 2003, 9, 2221-8.	3.2	50
43	Stimulation of adipose differentiation related protein (ADRP) expression in adipocyte precursors by long-chain fatty acids. <i>Journal of Cellular Physiology</i> , 2000, 182, 297-302.	2.0	97
44	Stimulation of adipose differentiation related protein (ADRP) expression in adipocyte precursors by long-chain fatty acids. , 2000, 182, 297.		1
45	Insulin but not IGF-I is required for the maintenance of the adipose phenotype in the adipogenic cell line 1246. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 1999, 35, 642-646.	0.7	4
46	Resveratrol, a natural product derived from grape, exhibits antiestrogenic activity and inhibits the growth of human breast cancer cells. , 1999, 179, 297-304.		280
47	Stimulation of PC Cell-Derived Growth Factor (Epithelin/Granulin Precursor) Expression by Estradiol in Human Breast Cancer Cells. <i>Biochemical and Biophysical Research Communications</i> , 1999, 256, 204-207.	1.0	46
48	Multiple forms of p55PIK, a regulatory subunit of phosphoinositide 3-kinase, are generated by alternative initiation of translation. <i>Biochemical Journal</i> , 1999, 341, 831-837.	1.7	18
49	Differentiation of newborn rat adipocyte precursors in defined serum-free medium. <i>In Vitro Cellular & Developmental Biology</i> , 1987, 23, 63-66.	1.0	29
50	Tumorigenicity associated with loss of differentiation and of response to insulin in the adipogenic cell line 1246. <i>In Vitro Cellular & Developmental Biology</i> , 1985, 21, 537-540.	1.0	23
51	An in vitro model to study adipose differentiation in serum-free medium. <i>Analytical Biochemistry</i> , 1982, 120, 351-359.	1.1	64
52	Isolation of myoblastic, fibro-adipogenic, and fibroblastic clonal cell lines from a common precursor and study of their requirements for growth and differentiation. <i>Experimental Cell Research</i> , 1981, 132, 313-327.	1.2	80
53	[6] The growth of cells in serum-free hormone-supplemented media. <i>Methods in Enzymology</i> , 1979, 58, 94-109.	0.4	279