

Mario Anders

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8909194/publications.pdf>

Version: 2024-02-01

26
papers

1,170
citations

430442

18
h-index

552369

26
g-index

26
all docs

26
docs citations

26
times ranked

1420
citing authors

#	ARTICLE	IF	CITATIONS
1	The Barrett's-associated variants at <i>GDF7</i> and <i>TBX5</i> also increase esophageal adenocarcinoma risk. <i>Cancer Medicine</i> , 2016, 5, 888-891.	1.3	21
2	Supportive evidence for <i>FOXP1</i> , <i>BARX1</i> , and <i>FOXF1</i> as genetic risk loci for the development of esophageal adenocarcinoma. <i>Cancer Medicine</i> , 2015, 4, 1700-1704.	1.3	26
3	Endoscopic mucosal resection of large colorectal adenomas: Only for expert centers?. <i>United European Gastroenterology Journal</i> , 2015, 3, 171-173.	1.6	2
4	Lectin Histochemistry Shows WGA, PHA-L and HPA Binding Increases During Progression of Human Colorectal Cancer. <i>Anticancer Research</i> , 2015, 35, 5333-9.	0.5	7
5	Long-term recurrence of neoplasia and Barrett's epithelium after complete endoscopic resection. <i>Gut</i> , 2014, 63, 1535-1543.	6.1	68
6	Subsquamous Extension of Intestinal Metaplasia Is Detected in 98% of Cases of Neoplastic Barrett's Esophagus. <i>Clinical Gastroenterology and Hepatology</i> , 2014, 12, 405-410.	2.4	36
7	Microarray meta-analysis defines global angiogenesis-related gene expression signatures in human carcinomas. <i>Molecular Carcinogenesis</i> , 2013, 52, 29-38.	1.3	18
8	A New Endoscopically Implantable Device (SatiSphere) for Treatment of Obesity—Efficacy, Safety, and Metabolic Effects on Glucose, Insulin, and GLP-1 Levels. <i>Obesity Surgery</i> , 2013, 23, 1727-1733.	1.1	39
9	Expression of the coxsackie adenovirus receptor in neuroendocrine lung cancers and its implications for oncolytic adenoviral infection. <i>Cancer Gene Therapy</i> , 2013, 20, 25-32.	2.2	19
10	Increased T-Helper 2 Cytokines in Bile From Patients With IgG4-Related Cholangitis Disrupt the Tight Junction—Associated Biliary Epithelial Cell Barrier. <i>Gastroenterology</i> , 2013, 144, 1116-1128.	0.6	53
11	Human Prostate Cancer in a Clinically Relevant Xenograft Mouse Model: Identification of $\beta^2(1,6)$ -Branched Oligosaccharides as a Marker of Tumor Progression. <i>Clinical Cancer Research</i> , 2012, 18, 1364-1373.	3.2	72
12	Endoscopic Detection of Colorectal Adenomas: Standards and Sophisticated Methods. <i>Digestive Diseases</i> , 2012, 30, 68-73.	0.8	1
13	Impact of the coxsackievirus and adenovirus receptor on the adenoma—carcinoma sequence of colon cancer. <i>British Journal of Cancer</i> , 2011, 104, 1426-1433.	2.9	26
14	Downregulation of the coxsackie and adenovirus receptor in cancer cells by hypoxia depends on HIF-1. <i>Cancer Gene Therapy</i> , 2010, 17, 141-146.	2.2	27
15	Probe-based confocal laser endomicroscopy compared with standard four-quadrant biopsy for evaluation of neoplasia in Barrett's esophagus. <i>Endoscopy</i> , 2010, 42, 435-440.	1.0	119
16	Endoscopic closure of GI fistulae by using an over-the-scope clip (with videos). <i>Gastrointestinal Endoscopy</i> , 2010, 72, 1289-1296.	0.5	114
17	Sodium Butyrate Increases Expression of the Coxsackie and Adenovirus Receptor in Colon Cancer Cells. <i>Cancer Investigation</i> , 2010, 28, 268-274.	0.6	8
18	Loss of Coxsackie and adenovirus receptor downregulates β -catenin expression. <i>British Journal of Cancer</i> , 2009, 101, 1574-1579.	2.9	14

#	ARTICLE	IF	CITATIONS
19	Expression and function of the coxsackie and adenovirus receptor in Barrett's esophagus and associated neoplasia. <i>Cancer Gene Therapy</i> , 2009, 16, 508-515.	2.2	14
20	Endoscopy in Barrett's oesophagus: adherence to standards and neoplasia detection in the community practice versus hospital setting. <i>Journal of Internal Medicine</i> , 2008, 264, 370-378.	2.7	9
21	Miniprobe confocal laser microscopy for the detection of invisible neoplasia in patients with Barrett's oesophagus. <i>Gut</i> , 2008, 57, 1648-1653.	6.1	192
22	Expression of EpCam and Villin in Barrett's Esophagus and in Gastric Cardia. <i>Disease Markers</i> , 2008, 24, 287-292.	0.6	9
23	Transforming Growth Factor- β 2 Receptor Inhibition Enhances Adenoviral Infectability of Carcinoma Cells via Up-Regulation of Coxsackie and Adenovirus Receptor in Conjunction with Reversal of Epithelial-Mesenchymal Transition. <i>Cancer Research</i> , 2006, 66, 1648-1657.	0.4	68
24	Disruption of 3D tissue integrity facilitates adenovirus infection by deregulating the coxsackievirus and adenovirus receptor. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 1943-1948.	3.3	95
25	Inhibition of the Raf/MEK/ERK pathway up-regulates expression of the coxsackievirus and adenovirus receptor in cancer cells. <i>Cancer Research</i> , 2003, 63, 2088-95.	0.4	68
26	Decreased expression of CD44 splicing variants in advanced colorectal carcinomas. <i>European Journal of Cancer</i> , 1998, 34, 1607-1611.	1.3	45