

Luigi Tornillo

List of Publications by Citations

Source: <https://exaly.com/author-pdf/8909307/luigi-tornillo-publications-by-citations.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

123
papers

6,118
citations

47
h-index

75
g-index

126
ext. papers

6,896
ext. citations

5.9
avg, IF

4.97
L-index

#	Paper	IF	Citations
123	Long noncoding RNA HOTTIP/HOXA13 expression is associated with disease progression and predicts outcome in hepatocellular carcinoma patients. <i>Hepatology</i> , 2014 , 59, 911-23	11.2	333
122	Clinical impact of programmed cell death ligand 1 expression in colorectal cancer. <i>European Journal of Cancer</i> , 2013 , 49, 2233-42	7.5	319
121	Diagnostic value of HSP70, glypican 3, and glutamine synthetase in hepatocellular nodules in cirrhosis. <i>Hepatology</i> , 2007 , 45, 725-34	11.2	318
120	High frequency of tumor-infiltrating FOXP3(+) regulatory T cells predicts improved survival in mismatch repair-proficient colorectal cancer patients. <i>International Journal of Cancer</i> , 2010 , 126, 2635-43	7.5	240
119	Glypican 3 expression in human nonneoplastic, preneoplastic, and neoplastic tissues: a tissue microarray analysis of 4,387 tissue samples. <i>American Journal of Clinical Pathology</i> , 2008 , 129, 899-906	1.9	193
118	The homeobox intestinal differentiation factor CDX2 is selectively expressed in gastrointestinal adenocarcinomas. <i>Modern Pathology</i> , 2004 , 17, 1392-9	9.8	158
117	Human and mouse VEGFA-amplified hepatocellular carcinomas are highly sensitive to sorafenib treatment. <i>Cancer Discovery</i> , 2014 , 4, 730-43	24.4	137
116	Hepatoid adenocarcinoma with liver metastasis mimicking hepatocellular carcinoma: an immunohistochemical and molecular study of eight cases. <i>American Journal of Surgical Pathology</i> , 2003 , 27, 1302-12	6.7	136
115	YAP promotes proliferation, chemoresistance, and angiogenesis in human cholangiocarcinoma through TEAD transcription factors. <i>Hepatology</i> , 2015 , 62, 1497-510	11.2	133
114	Is the improved prognosis of p16 positive oropharyngeal squamous cell carcinoma dependent of the treatment modality?. <i>International Journal of Cancer</i> , 2010 , 126, 1256-62	7.5	122
113	Enhanced expression of ANO1 in head and neck squamous cell carcinoma causes cell migration and correlates with poor prognosis. <i>PLoS ONE</i> , 2012 , 7, e43265	3.7	112
112	Loss of raf-1 kinase inhibitor protein expression is associated with tumor progression and metastasis in colorectal cancer. <i>American Journal of Clinical Pathology</i> , 2007 , 127, 820-7	1.9	107
111	Expression of voltage-gated potassium channels in human and mouse colonic carcinoma. <i>Clinical Cancer Research</i> , 2007 , 13, 824-31	12.9	104
110	Notch2 signaling promotes biliary epithelial cell fate specification and tubulogenesis during bile duct development in mice. <i>Hepatology</i> , 2009 , 50, 871-9	11.2	99
109	The Interplay Between Neutrophils and CD8 T Cells Improves Survival in Human Colorectal Cancer. <i>Clinical Cancer Research</i> , 2017 , 23, 3847-3858	12.9	97
108	Hepatocyte Paraffin 1 Expression in Human Normal and Neoplastic Tissues. <i>American Journal of Clinical Pathology</i> , 2004 , 122, 721-727	1.9	92
107	Constitutive Notch2 signaling induces hepatic tumors in mice. <i>Hepatology</i> , 2013 , 57, 1607-19	11.2	90

106	Histopathologic features and microsatellite instability of cancers of the papilla of vater and their precursor lesions. <i>American Journal of Surgical Pathology</i> , 2009 , 33, 691-704	6.7	90
105	HER2, TOP2A, CCND1, EGFR and C-MYC oncogene amplification in colorectal cancer. <i>Journal of Clinical Pathology</i> , 2007 , 60, 768-72	3.9	90
104	Disruption of Notch1 induces vascular remodeling, intussusceptive angiogenesis, and angiosarcomas in livers of mice. <i>Gastroenterology</i> , 2012 , 142, 967-977.e2	13.3	89
103	EphB2 expression across 138 human tumor types in a tissue microarray: high levels of expression in gastrointestinal cancers. <i>Clinical Cancer Research</i> , 2005 , 11, 6450-8	12.9	76
102	High myeloperoxidase positive cell infiltration in colorectal cancer is an independent favorable prognostic factor. <i>PLoS ONE</i> , 2013 , 8, e64814	3.7	75
101	Loss of p16 protein defines high-risk patients with gastrointestinal stromal tumors: a tissue microarray study. <i>Clinical Cancer Research</i> , 2005 , 11, 638-45	12.9	75
100	Tumor budding in colorectal cancer revisited: results of a multicenter interobserver study. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2015 , 466, 485-93	5.1	73
99	NK cells and T cells cooperate during the clinical course of colorectal cancer. <i>OncolImmunology</i> , 2014 , 3, e952197	7.2	73
98	Tumor infiltration by FcBIII (CD16)+ myeloid cells is associated with improved survival in patients with colorectal carcinoma. <i>International Journal of Cancer</i> , 2011 , 128, 2663-72	7.5	73
97	Loss of the CBX7 protein expression correlates with a more aggressive phenotype in pancreatic cancer. <i>European Journal of Cancer</i> , 2010 , 46, 1438-44	7.5	73
96	HLA class II antigen expression in colorectal carcinoma tumors as a favorable prognostic marker. <i>Neoplasia</i> , 2014 , 16, 31-42	6.4	72
95	Calcifying fibrous tumor of the stomach: clinicopathologic and molecular study of seven cases with literature review and reappraisal of histogenesis. <i>American Journal of Surgical Pathology</i> , 2010 , 34, 271-8	6.7	69
94	The RNA-binding protein RBM3 is required for cell proliferation and protects against serum deprivation-induced cell death. <i>Pediatric Research</i> , 2010 , 67, 35-41	3.2	65
93	The loss of the CBX7 gene expression represents an adverse prognostic marker for survival of colon carcinoma patients. <i>European Journal of Cancer</i> , 2010 , 46, 2304-13	7.5	65
92	Screening for ALK in non-small cell lung carcinomas: 5A4 and D5F3 antibodies perform equally well, but combined use with FISH is recommended. <i>Lung Cancer</i> , 2015 , 89, 104-9	5.9	62
91	Microscopic gastrointestinal stromal tumors in esophageal and intestinal surgical resection specimens: a clinicopathologic, immunohistochemical, and molecular study of 19 lesions. <i>American Journal of Surgical Pathology</i> , 2008 , 32, 867-73	6.7	62
90	Galectin-1 and its involvement in hepatocellular carcinoma aggressiveness. <i>Molecular Medicine</i> , 2010 , 16, 102-15	6.2	61
89	Prognostic significance of mammalian sterile20-like kinase 1 in colorectal cancer. <i>Modern Pathology</i> , 2007 , 20, 331-8	9.8	61

88	Patterns of gene amplification in gastrointestinal stromal tumors (GIST). <i>Laboratory Investigation</i> , 2005 , 85, 921-31	5.9	55
87	The HOX gene network in hepatocellular carcinoma. <i>International Journal of Cancer</i> , 2011 , 129, 2577-87	7.5	54
86	Loss of APAF-1 expression is associated with tumour progression and adverse prognosis in colorectal cancer. <i>European Journal of Cancer</i> , 2007 , 43, 1101-7	7.5	54
85	Marked genetic similarities between hepatitis B virus-positive and hepatitis C virus-positive hepatocellular carcinomas. <i>Journal of Pathology</i> , 2000 , 192, 307-12	9.4	52
84	Interferon- β -stimulated genes, but not USP18, are expressed in livers of patients with acute hepatitis C. <i>Gastroenterology</i> , 2012 , 143, 777-786.e6	13.3	51
83	Clinical features and genotype-phenotype correlations in children with progressive familial intrahepatic cholestasis type 3 related to ABCB4 mutations. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2011 , 52, 73-83	2.8	51
82	Differential significance of tumour infiltrating lymphocytes in sporadic mismatch repair deficient versus proficient colorectal cancers: a potential role for dysregulation of the transforming growth factor-beta pathway. <i>European Journal of Cancer</i> , 2007 , 43, 624-31	7.5	51
81	NY-ESO-1/LAGE-1 coexpression with MAGE-A cancer/testis antigens: a tissue microarray study. <i>International Journal of Cancer</i> , 2005 , 115, 960-6	7.5	51
80	Chromosomal alterations in hepatocellular nodules by comparative genomic hybridization: high-grade dysplastic nodules represent early stages of hepatocellular carcinoma. <i>Laboratory Investigation</i> , 2002 , 82, 547-53	5.9	50
79	Silencing of the SEC62 gene inhibits migratory and invasive potential of various tumor cells. <i>International Journal of Cancer</i> , 2011 , 128, 2284-95	7.5	49
78	Defective infiltration of natural killer cells in MICA/B-positive renal cell carcinoma involves beta(2)-integrin-mediated interaction. <i>Neoplasia</i> , 2009 , 11, 662-71	6.4	49
77	Close association between HER-2 amplification and overexpression in human tumors of non-breast origin. <i>Modern Pathology</i> , 2007 , 20, 192-8	9.8	49
76	Melanoma cells inhibit NK cell functions. <i>Cancer Research</i> , 2012 , 72, 5428-9; author reply 5430	10.1	47
75	HOX D13 expression across 79 tumor tissue types. <i>International Journal of Cancer</i> , 2009 , 125, 1532-41	7.5	46
74	GM-CSF Production by Tumor Cells Is Associated with Improved Survival in Colorectal Cancer. <i>Clinical Cancer Research</i> , 2014 , 20, 3094-106	12.9	45
73	Multiple sporadic gastrointestinal stromal tumors (GISTs) of the proximal stomach are caused by different somatic KIT mutations suggesting a field effect. <i>American Journal of Surgical Pathology</i> , 2008 , 32, 1553-9	6.7	45
72	Overexpression of the receptor for hyaluronic acid mediated motility is an independent adverse prognostic factor in colorectal cancer. <i>Modern Pathology</i> , 2006 , 19, 1302-9	9.8	45
71	Chromosomal imbalances in small cell carcinomas of the urinary bladder. <i>Journal of Pathology</i> , 1999 , 189, 230-5	9.4	43

70	Tenascin-W, a new marker of cancer stroma, is elevated in sera of colon and breast cancer patients. <i>International Journal of Cancer</i> , 2008 , 122, 2454-61	7.5	42
69	Bioreactor-engineered cancer tissue-like structures mimic phenotypes, gene expression profiles and drug resistance patterns observed "in vivo". <i>Biomaterials</i> , 2015 , 62, 138-46	15.6	41
68	Clinical significance of cell cycle- and apoptosis-related markers in biliary tract cancer: a tissue microarray-based approach revealing a distinctive immunophenotype for intrahepatic and extrahepatic cholangiocarcinomas. <i>American Journal of Clinical Pathology</i> , 2008 , 130, 780-6	1.9	41
67	Activation of pro-inflammatory and anti-inflammatory cytokines in host organs during chronic allograft rejection: role of endothelin receptor signaling. <i>American Journal of Transplantation</i> , 2005 , 5, 1042-9	8.7	40
66	MAGE-A Antigens and Cancer Immunotherapy. <i>Frontiers in Medicine</i> , 2017 , 4, 18	4.9	39
65	Role of the mitogen-activated protein kinase and phosphoinositide 3-kinase/AKT pathways downstream molecules, phosphorylated extracellular signal-regulated kinase, and phosphorylated AKT in colorectal cancer-a tissue microarray-based approach. <i>Human Pathology</i> , 2006 , 37, 1022-31	3.7	37
64	Malignant gastrointestinal leiomyosarcoma and gastrointestinal stromal tumor with prominent osteoclast-like giant cells. <i>Archives of Pathology and Laboratory Medicine</i> , 2004 , 128, 440-3	5	37
63	PED is overexpressed and mediates TRAIL resistance in human non-small cell lung cancer. <i>Journal of Cellular and Molecular Medicine</i> , 2008 , 12, 2416-26	5.6	35
62	MAGE-A10 is a nuclear protein frequently expressed in high percentages of tumor cells in lung, skin and urothelial malignancies. <i>International Journal of Cancer</i> , 2011 , 129, 1137-48	7.5	34
61	Comparative genomic hybridization analysis of hepatoblastoma reveals high frequency of X-chromosome gains and similarities between epithelial and stromal components. <i>Human Pathology</i> , 2003 , 34, 864-71	3.7	31
60	A spectrum of histopathologic findings in autoimmune liver disease. <i>American Journal of Clinical Pathology</i> , 2000 , 114, 705-11	1.9	31
59	Hepatocyte paraffin 1 expression in human normal and neoplastic tissues: tissue microarray analysis on 3,940 tissue samples. <i>American Journal of Clinical Pathology</i> , 2004 , 122, 721-7	1.9	31
58	High expression of HOXA13 correlates with poorly differentiated hepatocellular carcinomas and modulates sorafenib response in in vitro models. <i>Laboratory Investigation</i> , 2018 , 98, 95-105	5.9	30
57	OX40 expression enhances the prognostic significance of CD8 positive lymphocyte infiltration in colorectal cancer. <i>Oncotarget</i> , 2015 , 6, 37588-99	3.3	29
56	Frequency, phenotype, and genotype of minute gastrointestinal stromal tumors in the stomach: an autopsy study. <i>Human Pathology</i> , 2011 , 42, 1849-54	3.7	28
55	Co-overexpression of p21 and Ki-67 in head and neck squamous cell carcinoma relative to a significantly poor prognosis. <i>Head and Neck</i> , 2011 , 33, 267-73	4.2	27
54	Systematic assessment of the prognostic impact of membranous CD44v6 protein expression in colorectal cancer. <i>Histopathology</i> , 2009 , 55, 564-75	7.3	27
53	Effect of EpCAM, CD44, CD133 and CD166 expression on patient survival in tumours of the ampulla of Vater. <i>Journal of Clinical Pathology</i> , 2012 , 65, 140-5	3.9	27

52	Multiple sporadic gastrointestinal stromal tumours arising at different gastrointestinal sites: pattern of involvement of the muscularis propria as a clue to independent primary GISTs. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2009 , 455, 101-8	5.1	26
51	HMGA1 Expression in Human Hepatocellular Carcinoma Correlates with Poor Prognosis and Promotes Tumor Growth and Migration in in vitro Models. <i>Neoplasia</i> , 2016 , 18, 724-731	6.4	25
50	Prognostic value of cell cycle and apoptosis regulatory proteins in mismatch repair-proficient colorectal cancer: a tissue microarray-based approach. <i>American Journal of Clinical Pathology</i> , 2007 , 127, 114-23	1.9	24
49	The clinical impact of p16 status in fine-needle aspirates of cervical lymph node metastasis of head and neck squamous cell carcinomas. <i>European Archives of Oto-Rhino-Laryngology</i> , 2013 , 270, 661-7	3.5	23
48	Is immunohistochemical epidermal growth factor receptor expression overestimated as a prognostic factor in head-neck squamous cell carcinoma? A retrospective analysis based on a tissue microarray of 365 carcinomas. <i>Human Pathology</i> , 2008 , 39, 1527-34	3.7	23
47	Differential cell cycle and proliferation marker expression in ductal pancreatic adenocarcinoma and pancreatic intraepithelial neoplasia (PanIN). <i>Pathology</i> , 2010 , 42, 229-34	1.6	19
46	VEGFA gene locus (6p12) amplification identifies a small but highly aggressive subgroup of colorectal cancer [corrected] patients. <i>Modern Pathology</i> , 2011 , 24, 1404-12	9.8	19
45	Skp2 expression is associated with high risk and elevated Ki67 expression in gastrointestinal stromal tumours. <i>BMC Cancer</i> , 2008 , 8, 134	4.8	19
44	Endothelin inhibition delays onset of hyperglycemia and associated vascular injury in type I diabetes: evidence for endothelin release by pancreatic islet beta-cells. <i>Biochemical and Biophysical Research Communications</i> , 2005 , 334, 689-95	3.4	19
43	Transcriptional regulation of vascular bone morphogenetic protein by endothelin receptors in early autoimmune diabetes mellitus. <i>Life Sciences</i> , 2006 , 78, 2213-8	6.8	19
42	VEGFA gene locus analysis across 80 human tumour types reveals gene amplification in several neoplastic entities. <i>Angiogenesis</i> , 2014 , 17, 519-27	10.6	16
41	Low prevalence of SV40 in Swiss mesothelioma patients after elimination of false-positive PCR results. <i>Lung Cancer</i> , 2007 , 57, 282-91	5.9	16
40	Hepatocellular Carcinoma Xenografts Established From Needle Biopsies Preserve the Characteristics of the Originating Tumors. <i>Hepatology Communications</i> , 2019 , 3, 971-986	6	15
39	An unusual association of malignant gastrointestinal neuroectodermal tumor (clear cell sarcoma-like) and Ewing sarcoma. <i>Pathology Research and Practice</i> , 2015 , 211, 688-92	3.4	14
38	Absence of myeloperoxidase and CD8 positive cells in colorectal cancer infiltrates identifies patients with severe prognosis. <i>Oncolmmunology</i> , 2015 , 4, e1050574	7.2	14
37	Clinicopathological Features and Metastatic Pattern of Hepatocellular Carcinoma: An Autopsy Study of 398 Patients. <i>Pathobiology</i> , 2016 , 83, 301-7	3.6	14
36	Colorectal carcinoma infiltration by myeloperoxidase-expressing neutrophil granulocytes is associated with favorable prognosis. <i>Oncolmmunology</i> , 2013 , 2, e25990	7.2	14
35	KRAS mutation testing in colorectal cancer: comparison of the results obtained using 3 different methods for the analysis of codons G12 and G13. <i>Diagnostic Molecular Pathology</i> , 2012 , 21, 14-23		13

34	Immunophenotyping and oncogene amplifications in tumors of the papilla of Vater. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2008 , 453, 579-88	5.1	13
33	Vascular endothelial growth factor A amplification in colorectal cancer is associated with reduced M1 and M2 macrophages and diminished PD-1-expressing lymphocytes. <i>PLoS ONE</i> , 2017 , 12, e0175563	3.7	12
32	Gastrointestinal stromal tumor - an evolving concept. <i>Frontiers in Medicine</i> , 2014 , 1, 43	4.9	12
31	Acinic cell carcinoma of the breast arising in microglandular adenosis. <i>Case Reports in Pathology</i> , 2013 , 2013, 736048	0.9	11
30	DNA methylation instability by BRAF-mediated TET silencing and lifestyle-exposure divides colon cancer pathways. <i>Clinical Epigenetics</i> , 2019 , 11, 196	7.7	11
29	Human Papillomavirus (HPV) Detection in Cytologic Specimens: Similarities and Differences of Available Methodology. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2017 , 25, 184-189	1.9	10
28	High frequency of CD8 positive lymphocyte infiltration correlates with lack of lymph node involvement in early rectal cancer. <i>Disease Markers</i> , 2014 , 2014, 792183	3.2	10
27	Downregulation of the endothelial genes Notch1 and ephrinB2 in patients with nodular regenerative hyperplasia. <i>Liver International</i> , 2014 , 34, 594-603	7.9	10
26	Coexistence of primary gastric giant cell-rich leiomyosarcoma and gastrointestinal stromal tumor: report of a very rare combination and review of the literature. <i>International Journal of Surgical Pathology</i> , 2012 , 20, 74-8	1.2	10
25	Combined histomorphologic and immunohistochemical phenotype to predict the presence of vascular invasion in colon cancer. <i>Diseases of the Colon and Rectum</i> , 2009 , 52, 1114-21	3.1	10
24	Identification of New Players in Hepatocarcinogenesis: Limits and Opportunities of Using Tissue Microarray (TMA). <i>Microarrays (Basel, Switzerland)</i> , 2014 , 3, 91-102		9
23	Chromosomal instability in gastric mucosa-associated lymphoid tissue lymphomas: a fluorescent in situ hybridization study using a tissue microarray approach. <i>Human Pathology</i> , 2008 , 39, 536-42	3.7	9
22	Hepatic Notch1 deletion predisposes to diabetes and steatosis via glucose-6-phosphatase and perilipin-5 upregulation. <i>Laboratory Investigation</i> , 2016 , 96, 972-80	5.9	8
21	Hands-On Experience: Accreditation of Pathology Laboratories according to ISO 15189. <i>Pathobiology</i> , 2017 , 84, 121-129	3.6	7
20	Case report: appearance of an intestinal metastasis from intrahepatic cholangiocarcinoma occurring 5 years after resection of the primary tumor. <i>European Journal of Gastroenterology and Hepatology</i> , 2010 , 22, 892-4	2.2	7
19	Low Expression of Programmed Death 1 (PD-1), PD-1 Ligand 1 (PD-L1), and Low CD8+ T Lymphocyte Infiltration Identify a Subgroup of Patients With Gastric and Esophageal Adenocarcinoma With Severe Prognosis. <i>Frontiers in Medicine</i> , 2020 , 7, 144	4.9	6
18	Expression of CD24, P-cadherin and S100A4 in tumors of the ampulla of Vater. <i>Modern Pathology</i> , 2009 , 22, 306-13	9.8	6
17	PipelIT: A Singularity Container for Molecular Diagnostic Somatic Variant Calling on the Ion Torrent Next-Generation Sequencing Platform. <i>Journal of Molecular Diagnostics</i> , 2019 , 21, 884-894	5.1	5

16	Incarcerated umbilical hernia of unexpected origin: a primitive neuroectodermal tumor with early recurrence. <i>Journal of Clinical Oncology</i> , 2014 , 32, e3-6	2.2	5
15	KIT, PDGFRalpha and EGFR analysis in nephroblastoma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2008 , 452, 637-50	5.1	5
14	Molecular Profile of Gastrointestinal Stromal Tumors in Sixty-Eight Patients from a Single Swiss Institution. <i>Pathobiology</i> , 2020 , 87, 171-178	3.6	3
13	SH2D4A is frequently downregulated in hepatocellular carcinoma and cirrhotic nodules. <i>European Journal of Cancer</i> , 2014 , 50, 731-8	7.5	3
12	Clinicopathologic and immunohistochemical study of surgically treated primary gastric MALT lymphoma. <i>Journal of Surgical Oncology</i> , 2003 , 83, 106-11	2.8	3
11	Infiltration by IL22-Producing T Cells Promotes Neutrophil Recruitment and Predicts Favorable Clinical Outcome in Human Colorectal Cancer. <i>Cancer Immunology Research</i> , 2020 , 8, 1452-1462	12.5	3
10	Sensitive detection methods are key to identify secondary EGFR c.2369C>T p.(Thr790Met) in non-small cell lung cancer tissue samples. <i>BMC Cancer</i> , 2020 , 20, 366	4.8	2
9	Re-Punching Tissue Microarrays Is Possible: Why Can This Be Useful and How to Do It. <i>Microarrays (Basel, Switzerland)</i> , 2015 , 4, 245-54		2
8	Impaired vascular function in normoglycemic mice prone to autoimmune diabetes: role of nitric oxide. <i>European Journal of Pharmacology</i> , 2007 , 557, 161-7	5.3	2
7	Nestin and CD34 expression in colorectal cancer predicts improved overall survival. <i>Acta Oncologica</i> , 2021 , 60, 727-734	3.2	2
6	Biology of gastrointestinal stromal tumour and mechanisms of imatinib resistance. <i>Diagnostic Histopathology</i> , 2013 , 19, 203-210	0.7	1
5	Diffuse minute clear cell proliferation in kidney: case report and review of literature. <i>Urology</i> , 2009 , 73, 443.e9-11	1.6	1
4	The Genomic Landscape of Serrated Lesion of the Colorectum: Similarities and Differences With Tubular and Tubulovillous Adenomas. <i>Frontiers in Oncology</i> , 2021 , 11, 668466	5.3	0
3	Salmonella enterocolitis and hemorrhagic shock. <i>Gastrointestinal Endoscopy</i> , 2007 , 65, 1077-8; discussion 1078	5.2	
2	CDX2 Immunostaining in Primary and Secondary Ovarian Carcinomas. <i>Handbook of Immunohistochemistry and in Situ Hybridization of Human Carcinomas</i> , 2005 , 4, 393-397		
1	Medikamentöser toxischer Leberschaden (DILI). <i>Pathologie</i> , 2020 , 117-157	0	