

Fernanda Genre

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/8910190/fernanda-genre-publications-by-year.pdf>

Version: 2024-04-27

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

49
papers

603
citations

15
h-index

22
g-index

60
ext. papers

746
ext. citations

3.8
avg, IF

3.05
L-index

#	Paper	IF	Citations
49	Vaspin in atherosclerotic disease and cardiovascular risk in axial spondyloarthritis: a genetic and serological study. <i>Arthritis Research and Therapy</i> , 2021 , 23, 111	5.7	2
48	BAFF, APRIL and BAFFR on the pathogenesis of Immunoglobulin-A vasculitis. <i>Scientific Reports</i> , 2021 , 11, 11510	4.9	2
47	Endothelial Progenitor Cells: Relevant Players in the Vasculopathy and Lung Fibrosis Associated with the Presence of Interstitial Lung Disease in Systemic Sclerosis Patients. <i>Biomedicines</i> , 2021 , 9,	4.8	1
46	HLA association with the susceptibility to anti-synthetase syndrome. <i>Joint Bone Spine</i> , 2021 , 88, 105115	2.9	3
45	Integrative Analysis Reveals a Molecular Stratification of Systemic Autoimmune Diseases. <i>Arthritis and Rheumatology</i> , 2021 , 73, 1073-1085	9.5	27
44	Role of the IL33 and IL1RL1 pathway in the pathogenesis of Immunoglobulin A vasculitis. <i>Scientific Reports</i> , 2021 , 11, 16163	4.9	1
43	Omentin: a biomarker of cardiovascular risk in individuals with axial spondyloarthritis. <i>Scientific Reports</i> , 2020 , 10, 9636	4.9	6
42	Disease Activity Influences Cardiovascular Risk Reclassification Based on Carotid Ultrasound in Patients with Psoriatic Arthritis. <i>Journal of Rheumatology</i> , 2020 , 47, 1344-1353	4.1	8
41	Influence of MUC5B gene on antisynthetase syndrome. <i>Scientific Reports</i> , 2020 , 10, 1415	4.9	7
40	Endothelial Progenitor Cells as a Potential Biomarker in Interstitial Lung Disease Associated with Rheumatoid Arthritis. <i>Journal of Clinical Medicine</i> , 2020 , 9,	5.1	5
39	Reclassification into very-high cardiovascular risk after carotid ultrasound in patients with axial spondyloarthritis. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38, 724-731	2.2	4
38	Role of IRF5 in the pathogenesis of immunoglobulin-A vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2020 , 38 Suppl 124, 182-187	2.2	
37	Rapid beneficial effect of the IL-6 receptor blockade on insulin resistance and insulin sensitivity in non-diabetic patients with rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2019 , 37, 465-473	2.2	18
36	Relative Risk Chart Score for the Assessment of the Cardiovascular Risk in Young Patients with Ankylosing Spondylitis. <i>International Journal of Rheumatology</i> , 2018 , 2018, 1847894	2	3
35	Association of circulating calprotectin with lipid profile in axial spondyloarthritis. <i>Scientific Reports</i> , 2018 , 8, 13728	4.9	7
34	Detection of high cardiovascular risk patients with ankylosing spondylitis based on the assessment of abdominal aortic calcium as compared to carotid ultrasound. <i>Arthritis Research and Therapy</i> , 2018 , 20, 195	5.7	1
33	Cardiovascular risk stratification in axial spondyloarthritis: carotid ultrasound is more sensitive than coronary artery calcification score to detect high-cardiovascular risk axial spondyloarthritis patients. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 73-80	2.2	15

32	Implication of osteoprotegerin and sclerostin in axial spondyloarthritis cardiovascular disease: study of 163 Spanish patients. <i>Clinical and Experimental Rheumatology</i> , 2018 , 36, 302-309	2.2	5
31	Significant sE-Selectin levels reduction after 6 months of anti-TNF- α therapy in non-diabetic patients with moderate-to-severe psoriasis. <i>Journal of Dermatological Treatment</i> , 2017 , 28, 726-730	2.8	9
30	Protein tyrosine phosphatase non-receptor 22 and C-Src tyrosine kinase genes are down-regulated in patients with rheumatoid arthritis. <i>Scientific Reports</i> , 2017 , 7, 10525	4.9	10
29	A genome-wide association study suggests the HLA Class II region as the major susceptibility locus for IgA vasculitis. <i>Scientific Reports</i> , 2017 , 7, 5088	4.9	31
28	Asymmetric dimethylarginine but not osteoprotegerin correlates with disease severity in patients with moderate-to-severe psoriasis undergoing anti-tumor necrosis factor- α therapy. <i>Journal of Dermatology</i> , 2016 , 43, 389-94	1.6	10
27	Expression of osteoprotegerin and its ligands, RANKL and TRAIL, in rheumatoid arthritis. <i>Scientific Reports</i> , 2016 , 6, 29713	4.9	25
26	Cardiovascular risk assessment in patients with rheumatoid arthritis: The relevance of clinical, genetic and serological markers. <i>Autoimmunity Reviews</i> , 2016 , 15, 1013-1030	13.6	78
25	Proprotein convertase subtilisin/kexin type 9 in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2016 , 34, 1013-1019	2.2	9
24	HLA-DRB1 association with Henoch-Schonlein purpura. <i>Arthritis and Rheumatology</i> , 2015 , 67, 823-827	9.5	27
23	Anti-TNF- α therapy reduces endothelial cell activation in non-diabetic ankylosing spondylitis patients. <i>Rheumatology International</i> , 2015 , 35, 2069-78	3.6	19
22	Role of PTPN22 and CSK gene polymorphisms as predictors of susceptibility and clinical heterogeneity in patients with Henoch-Schönlein purpura (IgA vasculitis). <i>Arthritis Research and Therapy</i> , 2015 , 17, 286	5.7	8
21	Osteoprotegerin concentrations relate independently to established cardiovascular disease in rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2015 , 42, 39-45	4.1	23
20	Lack of association between JAK3 gene polymorphisms and cardiovascular disease in Spanish patients with rheumatoid arthritis. <i>BioMed Research International</i> , 2015 , 2015, 318364	3	5
19	Association of HLA-B*41:02 with Henoch-Schönlein Purpura (IgA Vasculitis) in Spanish individuals irrespective of the HLA-DRB1 status. <i>Arthritis Research and Therapy</i> , 2015 , 17, 102	5.7	21
18	Relationship between endothelial dysfunction and osteoprotegerin, vitamin D, and bone mineral density in patients with rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, 241-9	2.2	6
17	TNF-related apoptosis-inducing ligand and cardiovascular disease in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2015 , 33, 491-7	2.2	6
16	Antitumour necrosis factor α treatment reduces retinol-binding protein 4 serum levels in non-diabetic ankylosing spondylitis patients. <i>Annals of the Rheumatic Diseases</i> , 2014 , 73, 941-3	2.4	12
15	Patients with ankylosing spondylitis and low disease activity because of anti-TNF-alpha therapy have higher TRAIL levels than controls: a potential compensatory effect. <i>Mediators of Inflammation</i> , 2014 , 2014, 798060	4.3	6

14	Adipokines, biomarkers of endothelial activation, and metabolic syndrome in patients with ankylosing spondylitis. <i>BioMed Research International</i> , 2014 , 2014, 860651	3	28
13	IGF-1 and ADMA levels are inversely correlated in nondiabetic ankylosing spondylitis patients undergoing anti-TNF-alpha therapy. <i>BioMed Research International</i> , 2014 , 2014, 671061	3	4
12	Independent relationship of osteoprotegerin concentrations with endothelial activation and carotid atherosclerosis in patients with severe rheumatoid arthritis. <i>Journal of Rheumatology</i> , 2014 , 41, 429-36	4.1	26
11	Osteoprotegerin CGA haplotype protection against cerebrovascular complications in anti-CCP negative patients with rheumatoid arthritis. <i>PLoS ONE</i> , 2014 , 9, e106823	3.7	9
10	Osteoprotegerin correlates with disease activity and endothelial activation in non-diabetic ankylosing spondylitis patients undergoing TNF- α antagonist therapy. <i>Clinical and Experimental Rheumatology</i> , 2014 , 32, 640-6	2.2	12
9	The ZC3HC1 rs11556924 polymorphism is associated with increased carotid intima-media thickness in patients with rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2013 , 15, R152	5.7	21
8	Antitumour necrosis factor- α therapy modulates angiopoietin-2 serum levels in non-diabetic ankylosing spondylitis patients. <i>Annals of the Rheumatic Diseases</i> , 2013 , 72, 1265-7	2.4	9
7	SMAD3 rs17228212 gene polymorphism is associated with reduced risk to cerebrovascular accidents and subclinical atherosclerosis in anti-CCP negative Spanish rheumatoid arthritis patients. <i>PLoS ONE</i> , 2013 , 8, e77695	3.7	11
6	Adiponectin and resistin serum levels in non-diabetic ankylosing spondylitis patients undergoing TNF- α antagonist therapy. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 365-71	2.2	13
5	Apelin serum levels in non-diabetic ankylosing spondylitis patients undergoing TNF- α antagonist therapy. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 532-7	2.2	4
4	Leptin and visfatin serum levels in non-diabetic ankylosing spondylitis patients undergoing TNF- α antagonist therapy. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 538-45	2.2	15
3	Asymmetric dimethylarginine serum levels in non-diabetic ankylosing spondylitis patients undergoing TNF- α antagonist therapy. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 749-55	2.2	9
2	Angiopoietin-2 serum levels correlate with severity, early onset and cardiovascular disease in patients with rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 761-6	2.2	16
1	Correlation between insulin resistance and serum ghrelin in non-diabetic ankylosing spondylitis patients undergoing anti-TNF- α therapy. <i>Clinical and Experimental Rheumatology</i> , 2013 , 31, 913-8	2.2	4