

Lars Erik Holmer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/8918285/publications.pdf>

Version: 2024-02-01

148
papers

3,350
citations

136885

32
h-index

206029

48
g-index

154
all docs

154
docs citations

154
times ranked

1376
citing authors

#	ARTICLE	IF	CITATIONS
1	Fossil brachiopod identification using a new deep convolutional neural network. <i>Gondwana Research</i> , 2022, 105, 290-298.	3.0	10
2	Brachiopods from the Latham Shale Lagerstätte (Cambrian Series 2, Stage 4) and Cadiz Formation (Miaolingian, Wuliuan), California. <i>Journal of Paleontology</i> , 2022, 96, 61-80.	0.5	3
3	Submarine metalliferous carbonate mounds in the Cambrian of the Baltoscandian Basin induced by vent networks and water column stratification. <i>Scientific Reports</i> , 2022, 12, 8475.	1.6	1
4	The oldest Cambrian trilobite “ brachiopod association in South China. <i>Gondwana Research</i> , 2021, 89, 147-167.	3.0	16
5	First report of acrotretoid brachiopod shell beds in the lower Cambrian (Stage 4) Guanshan Biota of eastern Yunnan, South China. <i>Journal of Paleontology</i> , 2021, 95, 40-55.	0.5	6
6	Early Cambrian (Stage 4) brachiopods from the Shipai Formation in the Three Gorges area of South China. <i>Journal of Paleontology</i> , 2021, 95, 497-526.	0.5	10
7	Silurian (Aeronian) rhynchonelliform brachiopods of Shabdjereh, south-west Central Iran and their significance for early spiriferide evolution. <i>Journal of Systematic Palaeontology</i> , 2021, 19, 191-219.	0.6	2
8	Burrows filled with faecal pellets from the Cambrian (Stage 4) Guanshan biota of South China and their palaeoecological implications. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021, 567, 110249.	1.0	3
9	Go large or go conical: allometric trajectory of an early Cambrian acrotretide brachiopod. <i>Palaeontology</i> , 2021, 64, 727-741.	1.0	4
10	Using laser microprobe analysis to assess potential relationships between Cambrian tomotiids and organophosphatic brachiopods. <i>Journal of Analytical and Applied Pyrolysis</i> , 2021, 158, 105277.	2.6	1
11	Biomacromolecules in recent phosphate-shelled brachiopods: identification and characterization of chitin matrix. <i>Journal of Materials Science</i> , 2021, 56, 19884-19898.	1.7	3
12	Possible drill holes and pseudoborings in oboloid shells from the Cambrian/Ordovician boundary beds of Estonia and the uppermost Cambrian of NW Russia. <i>Historical Biology</i> , 2021, 33, 3579-3584.	0.7	3
13	First Report of Small Shelly Fossils from the Cambrian Miaolingian Limestones (Zhangxia and) Tj ETQq1 1 0.784314 rgBT /Overlock 10	0.8	2
14	Early Cambrian organophosphatic brachiopods from the Xinji Formation, at Shuiyu section, Shanxi Province, North China. <i>Palaeoworld</i> , 2020, 29, 512-533.	0.5	14
15	Characterization of organophosphatic brachiopod shells: spectroscopic assessment of collagen matrix and biomineral components. <i>RSC Advances</i> , 2020, 10, 38456-38467.	1.7	5
16	Ontogeny and evolutionary significance of a new acrotretide brachiopod genus from Cambrian Series 2 of South China. <i>Journal of Systematic Palaeontology</i> , 2020, 18, 1569-1588.	0.6	8
17	First report of brachiopods with soft parts from the Lower Cambrian Latham Shale (Series 2, Stage 4), California. <i>Science Bulletin</i> , 2020, 65, 1543-1546.	4.3	4
18	The oldest “ <i>Lingulellotreta</i> ” (Lingulata, Brachiopoda) from China and its phylogenetic significance: integrating new material from the Cambrian Stage 3 “4 Lagerstätten in eastern Yunnan, South China. <i>Journal of Systematic Palaeontology</i> , 2020, 18, 945-973.	0.6	15

#	ARTICLE	IF	CITATIONS
19	Brachiopods from the Byrd Group (Cambrian Series 2, Stage 4) Central Transantarctic Mountains, East Antarctica: biostratigraphy, phylogeny and systematics. <i>Papers in Palaeontology</i> , 2020, 6, 349-383.	0.7	15
20	Linguliform brachiopods from the Cambrian (Guzhangian) Karpinsk Formation of Novaya Zemlya. <i>Papers in Palaeontology</i> , 2020, 6, 571-592.	0.7	2
21	Shell structure, ornamentation and affinity of the problematic early Cambrian brachiopod <i>Heliomedusa orientalis</i> . <i>Lethaia</i> , 2020, 53, 574-587.	0.6	9
22	Glendonite occurrences in the Tremadocian of Baltica: first Early Palaeozoic evidence of massive ikaite precipitation at temperate latitudes. <i>Scientific Reports</i> , 2019, 9, 7205.	1.6	19
23	Cambrian rhynchonelliform nisusoid brachiopods: phylogeny and distribution. <i>Papers in Palaeontology</i> , 2019, 5, 559-575.	0.7	7
24	Mollusks from the upper Shackleton Limestone (Cambrian Series 2), Central Transantarctic Mountains, East Antarctica. <i>Journal of Paleontology</i> , 2019, 93, 437-459.	0.5	17
25	The problematic lingulate brachiopod <i>Aulonotreta</i> from the Ordovician (Dapingian–Darriwilian) of Baltoscandia. <i>Estonian Journal of Earth Sciences</i> , 2019, 68, 206.	0.4	3
26	Evolutionary significance of a middle Cambrian (Series 3) in situ occurrence of the pedunculate rhynchonelliform brachiopod <i>Nisusia sulcata</i> . <i>Lethaia</i> , 2018, 51, 424-432.	0.6	6
27	Earliest ontogeny of early Cambrian acrotretoid brachiopods – first evidence for metamorphosis and its implications. <i>BMC Evolutionary Biology</i> , 2018, 18, 42.	3.2	24
28	The attachment strategies of Cambrian kutorginate brachiopods: the curious case of two pedicle openings and their phylogenetic significance. <i>Journal of Paleontology</i> , 2018, 92, 33-39.	0.5	8
29	Post-metamorphic allometry in the earliest acrotretoid brachiopods from the lower Cambrian (Series 1) of the Burgess Shale. <i>Palaeontology</i> , 2017, 60, 269-279.	1.0	11
30	Do brachiopods show substrate-related phenotypic variation? A case study from the Burgess Shale. <i>Palaeontology</i> , 2017, 60, 269-279.	1.0	11
31	Ecology, biofacies, biogeography and systematics of micromorphic lingulate brachiopods from the Ordovician (Darriwilian–Sandbian) of south-central China. <i>Papers in Palaeontology</i> , 2017, 3, 317-361.	0.7	4
32	Unusual pitted Ordovician brachiopods from the East Baltic: the significance of coarsely pitted ornamentations in linguliforms. <i>Papers in Palaeontology</i> , 2017, 3, 387-399.	0.7	5
33	Pentameroid brachiopod <i>Karlsorus</i> new genus from the upper Wenlock (Silurian) Slite Beds, Gotland, Sweden. <i>Journal of Paleontology</i> , 2017, 91, 911-918.	0.5	2
34	Brachiopods: origin and early history. <i>Palaeontology</i> , 2017, 60, 609-631.	1.0	39
35	Gene Expression Patterns in Brachiopod Larvae Refute the Brachiopod-Fold Hypothesis. <i>Frontiers in Cell and Developmental Biology</i> , 2017, 5, 74.	1.8	4
36	The Cambrian brachiopod fauna from the first-trilobite age Shuijingtuo Formation in the Three Gorges area of China. <i>Palaeoworld</i> , 2016, 25, 333-355.	0.5	40

#	ARTICLE	IF	CITATIONS
37	Reassessment of the early Triassic lingulid brachiopod <i>Lingula borealis</i> Bittner, 1899 and related problems of lingulid taxonomy. <i>Gff</i> , 2016, 138, 519-525.	0.4	5
38	EXCEPTIONALLY PRESERVED <i>MICKWITZIA</i> FROM THE INDIAN SPRINGS LAGERSTÄTTE (CAMBRIAN STAGE 3), NEVADA AND IMPLICATIONS FOR EARLY CAMBRIAN BRACHIOPOD EVOLUTION. , 2016, , .		0
39	The early Cambrian tommotiid <i>Kulparina rostrata</i> from South Australia. <i>Journal of Paleontology</i> , 2015, 89, 920-932.	0.5	8
40	Exceptionally preserved <i>Mickwitzia</i> from the Indian Springs Lagerstätte (Cambrian Stage 3), Nevada. <i>Journal of Paleontology</i> , 2015, 89, 933-955.	0.5	8
41	Review of the Ordovician stratigraphy and fauna of the Anarak Region in Central Iran. <i>Acta Geologica Polonica</i> , 2015, 65, 403-435.	0.9	9
42	Competition and mimicry: the curious case of chaetae in brachiopods from the middle Cambrian Burgess Shale. <i>BMC Evolutionary Biology</i> , 2015, 15, 42.	3.2	24
43	Himalayan Cambrian brachiopods. <i>Papers in Palaeontology</i> , 2015, 1, 345-399.	0.7	52
44	First report of linguloid brachiopods with soft parts from the lower Cambrian (Series 2, Stage 4) of the Three Gorges area, South China. <i>Annales De Paleontologie</i> , 2015, 101, 167-177.	0.1	18
45	Taxonomy, morphology, shell structure and early ontogeny of <i>Pelmanotreta</i> nom. nov. from the lower Cambrian of Siberia. <i>Gff</i> , 2015, 137, 1-8.	0.4	15
46	Survival on a soft seafloor: life strategies of brachiopods from the Cambrian Burgess Shale. <i>Earth-Science Reviews</i> , 2015, 151, 266-287.	4.0	30
47	An early Cambrian agglutinated tubular lophophorate with brachiopod characters. <i>Scientific Reports</i> , 2014, 4, 4682.	1.6	40
48	New U-Pb zircon ages of the Sandbian (Upper Ordovician) K-bentonite in Baltoscandia (Estonia) <i>Tj ETQq 0 0 0 rgBT /Overlo</i>	0.4	14
49	<i>aterimitra pyramidalis</i> from South Australia: scleritome, shell structure and evolution of a lower Cambrian stem group brachiopod. <i>Palaeontology</i> , 2014, 57, 417-446.	1.0	27
50	Oldest glosselline linguliform brachiopod with soft parts from the Lower Cambrian of Yunnan, Southern China. <i>Gff</i> , 2014, 136, 539-547.	0.4	9
51	Ordovician "Silurian Chileida" First Post-Cambrian Records of an Enigmatic Group of Brachiopoda. <i>Journal of Paleontology</i> , 2014, 88, 488-496.	0.5	6
52	Adaptive strategies and environmental significance of lingulid brachiopods across the late Permian extinction. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014, 399, 373-384.	1.0	36
53	Brachiopods hitching a ride: an early case of commensalism in the middle Cambrian Burgess Shale. <i>Scientific Reports</i> , 2014, 4, 6704.	1.6	32
54	Metamorphosis in Craniiformea revisited: <i>Novocrania anomala</i> shows delayed development of the ventral valve. <i>Zoomorphology</i> , 2013, 132, 379-387.	0.4	16

#	ARTICLE	IF	CITATIONS
55	A sclerite-bearing stem group entoproct from the early Cambrian and its implications. <i>Scientific Reports</i> , 2013, 3, 1066.	1.6	37
56	Chapter 10 Biogeography of Ordovician linguliform and craniiform brachiopods. <i>Geological Society Memoir</i> , 2013, 38, 117-126.	0.9	18
57	Silurian craniide brachiopods from <sc>G</sc>otland. <i>Palaeontology</i> , 2013, 56, 1029-1044.	1.0	7
58	Morphology, ontogeny and affinities of the <sc>H</sc>irnantian triplisiid brachiopod <i>S</i><sc>treptis undifera</sc> from <sc>B</sc>altoscandia. <i>Palaeontology</i> , 2013, 56, 961-970.	1.0	3
59	Cambrian (Furongian) rhynchonelliform brachiopods from the Eastern Alborz Mountains, Iran. <i>Bulletin of Geosciences</i> , 2013, , 525-538.	0.5	10
60	The problematic early Cambrian fossil <i>Tumulduria incomperta</i> represents the detached ventral interarea of a paterinid brachiopod. <i>Acta Palaeontologica Polonica</i> , 2012, , .	0.4	0
61	Earliest ontogeny of Early Palaeozoic Craniiformea: compelling evidence for lecithotrophy. <i>Lethaia</i> , 2012, 45, 566-573.	0.6	10
62	Peduncular attached secondary tiering acrotretoid brachiopods from the Chengjiang fauna: Implications for the ecological expansion of brachiopods during the Cambrian explosion. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012, 323-325, 60-67.	1.0	36
63	An obolellate brachiopod with soft-part preservation from the Early Cambrian Chengjiang fauna of China. <i>Journal of Paleontology</i> , 2011, 85, 460-463.	0.5	18
64	First record of repaired durophagous shell damages in Early Cambrian lingulate brachiopods with preserved pedicles. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2011, 302, 206-212.	1.0	24
65	First record of a bivalved larval shell in Early Cambrian tomotiids and its phylogenetic significance. <i>Palaeontology</i> , 2011, 54, 235-239.	1.0	24
66	Scleritome construction, biofacies, biostratigraphy and systematics of the tomotiid <i>Eccentrotheca helenia</i> sp. nov. from the Early Cambrian of South Australia. <i>Palaeontology</i> , 2011, 54, 253-286.	1.0	68
67	Lower palaeozoic stratigraphy of Murchisonfjorden and Sparreneset, Nordaustlandet, Svalbard. <i>Geografiska Annaler, Series A: Physical Geography</i> , 2011, 93, 209-226.	0.6	9
68	The exceptionally preserved Early Cambrian stem rhynchonelliform brachiopod <i>Longtancunella</i> and its implications. <i>Lethaia</i> , 2011, 44, 490-495.	0.6	22
69	Relic aragonite from Ordovician–Silurian brachiopods: Implications for the evolution of calcification. <i>Geology</i> , 2011, 39, 967-970.	2.0	43
70	Taxonomy and biostratigraphy of Ordovician brachiopods from northeastern Ny Friesland, Spitsbergen. <i>Zootaxa</i> , 2011, 3076, 1.	0.2	14
71	Earliest ontogeny of Early Palaeozoic Craniiformea: implications for brachiopod phylogeny. <i>Lethaia</i> , 2010, 43, 323-333.	0.6	16
72	First record of the brachiopod <i>Lingulella waptaensis</i> with pedicle from the Middle Cambrian Burgess Shale. <i>Acta Zoologica</i> , 2010, 91, 150-162.	0.6	17

#	ARTICLE	IF	CITATIONS
73	<i>Setatella significans</i> , a new name for mickwitziid stem group brachiopods from the lower Cambrian of Greenland and Labrador. <i>Gff</i> , 2010, 132, 117-122.	0.4	15
74	Soft-Part Preservation in a Linguliform Brachiopod from the Lower Cambrian Wulongqing Formation (Guanshan Fauna) of Yunnan, South China. <i>Acta Palaeontologica Polonica</i> , 2010, 55, 495-505.	0.4	23
75	Diversity fluctuations and biogeography of Ordovician brachiopod faunas in northeastern Spitsbergen. <i>Bulletin of Geosciences</i> , 2010, , 497-504.	0.5	14
76	Homologous skeletal secretion in tomotiids and brachiopods. <i>Geology</i> , 2009, 37, 1143-1146.	2.0	47
77	The scleritome of <i>Paterimitra</i> : an Early Cambrian stem group brachiopod from South Australia. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2009, 276, 1651-1656.	1.2	54
78	Architecture and function of the lophophore in the problematic brachiopod <i>Heliomedusa orientalis</i> (Early Cambrian, South China). <i>Geobios</i> , 2009, 42, 649-661.	0.7	34
79	Early ontogeny and soft tissue preservation in siphonotretide brachiopods: New data from the Cambrian-Ordovician of Iran. <i>Gondwana Research</i> , 2009, 16, 151-161.	3.0	30
80	First report of the early Cambrian stem group brachiopod <i>Mickwitzia</i> from East Gondwana. <i>Gondwana Research</i> , 2009, 16, 145-150.	3.0	21
81	The first occurrence of a lingulid brachiopod from the Cretaceous of Sergipe, Brazil, with a restudy of <i>Lingula bagualensis</i> Wilckens, 1905 from southern Patagonia. <i>Palaontologische Zeitschrift</i> , 2009, 83, 255-266.	0.8	8
82	THE ENIGMATIC EARLY CAMBRIAN <i>SALANYGOLINA</i> A STEM GROUP OF RHYNCHONELLIFORM CHILEATE BRACHIOPODS?. <i>Palaeontology</i> , 2009, 52, 1-10.	1.0	45
83	Gondwanan faunal signatures from Early Palaeozoic terranes of Kazakhstan and Central Asia: evidence and tectonic implications. <i>Geological Society Special Publication</i> , 2009, 325, 23-64.	0.8	40
84	The Early Cambrian tomotioid <i>Micrina</i> , a sessile bivalved stem group brachiopod. <i>Biology Letters</i> , 2008, 4, 724-728.	1.0	82
85	Middle Cambrian to Lower Ordovician faunas from the Chingiz Mountain Range, central Kazakhstan. <i>Alcheringa</i> , 2008, 32, 443-463.	0.5	12
86	The scleritome of <i>Eccentrotheca</i> from the Lower Cambrian of South Australia: Lophophorate affinities and implications for tomotioid phylogeny. <i>Geology</i> , 2008, 36, 171.	2.0	105
87	Columnar shell structures in early linguloid brachiopods new data from the Middle Cambrian of Sweden. <i>Earth and Environmental Science Transactions of the Royal Society of Edinburgh</i> , 2007, 98, 221-232.	0.3	16
88	Earliest ontogeny of Middle Ordovician rhynchonelliform brachiopods (<i>Clitambonitoidea</i> and <i>Tj</i> ETQq0 0 0 rgBT /Overlock 10, If 50 142	0.6	37
89	A spinose stem group brachiopod with pedicle from the Middle Cambrian Burgess Shale. <i>Acta Zoologica</i> , 2006, 87, 273-290.	0.6	43
90	Proposed stratotype for the base of the highest Cambrian stage at the first appearance datum of <i>Cordylodus andresi</i> , Lawson Cove section, Utah, USA. <i>Palaeoworld</i> , 2006, 15, 384-405.	0.5	31

#	ARTICLE	IF	CITATIONS
91	New and poorly known acrotretid brachiopods (Class Lingulata) from the Cedaria-Crepicephalus zone (late Middle Cambrian) of the Great Basin, USA. <i>Geobios</i> , 2006, 39, 125-154.	0.7	19
92	ENDOSYMBIOSIS IN ORDOVICIAN SILURIAN CORALS AND STROMATOPOROIDS: A NEW LINGULID AND ITS TRACE FROM EASTERN CANADA. <i>Journal of Paleontology</i> , 2006, 80, 750-759.	0.5	28
93	The Lower Cambrian brachiopod <i>Kyrshabaktella</i> and associated shelly fossils from the Harkless Formation, southern Nevada. <i>Gff</i> , 2006, 128, 327-337.	0.4	27
94	Early Middle Ordovician (Billingen Volkhov stages) Orthide and Protorthide brachiopods from the East Baltic. <i>Gff</i> , 2006, 128, 339-348.	0.4	0
95	EARLY CAMBRIAN BRACHIOPODS FROM NORTH-EAST GREENLAND. <i>Palaeontology</i> , 2005, 48, 325-345.	1.0	48
96	Discovery of a new type of shell structure within the organophosphatic brachiopods and the status of the family Curticiidae. <i>Gff</i> , 2005, 127, 7-16.	0.4	7
97	Neodymium isotopic composition of Cambrian Ordovician biogenic apatite in the Baltoscandian Basin: implications for palaeogeographical evolution and patterns of biodiversity. <i>Geological Magazine</i> , 2005, 142, 419-439.	0.9	40
98	LOWER ORDOVICIAN (TREMADOCIAN) LINGULATE BRACHIOPODS FROM THE HOUSE AND FILLMORE FORMATIONS, IBEX AREA, WESTERN UTAH, USA. <i>Journal of Paleontology</i> , 2005, 79, 884-906.	0.5	33
99	The oldest-known metazoan parasite?. <i>Journal of Paleontology</i> , 2004, 78, 1214-1216.	0.5	15
100	22. Tube-Shaped Incertae Sedis. , 2004, , 214-222.		13
101	Conodont biostratigraphy and faunal assemblages in radiolarian ribbon-banded cherts of the Burubaital Formation, West Balkhash Region, Kazakhstan. <i>Geological Magazine</i> , 2004, 141, 699-715.	0.9	34
102	Early Cambrian lingulate brachiopods from the Shaanxi Province, China. <i>Gff</i> , 2004, 126, 193-211.	0.4	49
103	Chemico-structure of the organophosphatic shells of siphonotretide brachiopods. <i>Palaeontology</i> , 2004, 47, 1313-1337.	1.0	19
104	THE OLDEST-KNOWN METAZOAN PARASITE?. <i>Journal of Paleontology</i> , 2004, 78, 1214-1216.	0.5	27
105	The brachiopod fold: a neglected body plan hypothesis. <i>Palaeontology</i> , 2003, 46, 59-65.	1.0	55
106	Faunal composition and dynamics in unconsolidated sediments: a case study from the Middle Ordovician of the East Baltic. <i>Geological Magazine</i> , 2003, 140, 31-44.	0.9	25
107	Understanding linguloid brachiopods: Obolus and Ungula as examples. <i>Carnets De Geologie</i> , 2003, , .	0.4	3
108	LINGULATE BRACHIOPODS FROM THE CAMBRIAN-ORDOVICIAN BOUNDARY BEDS OF UTAH. <i>Journal of Paleontology</i> , 2002, 76, 211-228.	0.5	17

#	ARTICLE	IF	CITATIONS
109	Brachiopods: Cambrian-Tremadoc precursors to Ordovician radiation events. Geological Society Special Publication, 2002, 194, 13-23.	0.8	37
110	Lingulate brachiopods from the Cambrian-Ordovician boundary beds of Utah. Journal of Paleontology, 2002, 76, 211-228.	0.5	24
111	Shell Structure And Inferred Growth, Functions And Affinities Of The Sclerites Of The Problematic Micrina. Palaeontology, 2002, 45, 845-873.	1.0	78
112	A Stem Group Brachiopod From The Lower Cambrian: Support For A Micrina (Halkieriid) Ancestry. Palaeontology, 2002, 45, 875-882.	1.0	95
113	Spatial variations in faunal composition, Middle Ordovician, Volkhov Stage, East Baltic. Gff, 2001, 123, 65-72.	0.4	11
114	Phylogeny and Classification: Linguliformea and Craniiformea. The Paleontological Society Papers, 2001, 7, 11-26.	0.8	14
115	The Hunneberg Stage (Ordovician) in the area east of St. Petersburg, north-western Russia. Palaeontologische Zeitschrift, 2001, 74, 543-561.	0.8	28
116	Functional morphology of articulatory structures and implications for patterns of musculature in Cambrian rhynchonelliform brachiopods. Systematics Association Special Volume, 2001, , 163-176.	0.2	3
117	MAN1, an Inner Nuclear Membrane Protein That Shares the LEM Domain with Lamina-associated Polypeptide 2 and Emerin. Journal of Biological Chemistry, 2000, 275, 4840-4847.	1.6	289
118	Redescription of the Ordovician acrotretoid brachiopod <i>Conotreta</i> Walcott, 1889. Gff, 2000, 122, 313-318.	0.4	5
119	Early Ordovician organophosphatic brachiopods with Baltoscandian affinities from the Alay Range, southern Kyrgyzstan. Gff, 2000, 122, 367-375.	0.4	13
120	Organophosphatic brachiopods: Patterns of biodiversification and extinction in the Early Palaeozoic. Geobios, 1999, 32, 145-163.	0.7	54
121	Cambrian phosphatic brachiopods from the Precordillera of western Argentina. Gff, 1999, 121, 227-242.	0.4	13
122	A human HP1 pseudogene maps to chromosome 11p14. Somatic Cell and Molecular Genetics, 1998, 24, 353-356.	0.7	3
123	The Human Lamin B Receptor/Sterol Reductase Multigene Family. Genomics, 1998, 54, 469-476.	1.3	133
124	Nd isotope composition and rare earth element distribution in early Paleozoic biogenic apatite from Baltoscandia: A signature of Iapetus ocean water. Geology, 1998, 26, 1083.	2.0	38
125	Early Cambrian Lingulellotreta (Lingulata, Brachiopoda) from South Kazakhstan (Malyi Karatau Range) and South China (Eastern Yunnan). Journal of Paleontology, 1997, 71, 577-584.	0.5	35
126	Late Ordovician and early Silurian Trimerellide brachiopods from Kazakhstan. Journal of Paleontology, 1997, 71, 584-598.	0.5	25

#	ARTICLE	IF	CITATIONS
127	Middle Ordovician (Llanvirn) unguulate brachiopods and conodonts from the Malyi Karatau Range, Kazakhstan. <i>Palaontologische Zeitschrift</i> , 1996, 70, 481-495.	0.8	6
128	Late Ordovician brachiopod assemblage of Hiberno-Salairian type from Central Kazakhstan. <i>Gff</i> , 1996, 118, 83-96.	0.4	17
129	The elkaniide brachiopod <i>Volborthia</i> from the lower ordovician of Baltoscandia. <i>Palaontologische Zeitschrift</i> , 1995, 69, 213-221.	0.8	3
130	Review of the Cambrian acrotretid brachiopod <i>Neotreta</i> . <i>Alcheringa</i> , 1994, 18, 345-357.	0.5	7
131	Ceratretide brachiopods from the lower and middle Cambrian of Sweden, Kazakhstan, and Siberia. <i>Gff</i> , 1994, 116, 203-210.	0.4	4
132	Cambrian-Ordovician lingulate brachiopods from Scandinavia, Kazakhstan, and South Ural Mountains. <i>Lethaia</i> , 1994, 27, 1-156.	0.6	13
133	Revision of the type species of <i>Acrotreta</i> and related lingulate brachiopods. <i>Journal of Paleontology</i> , 1994, 68, 433-450.	0.5	18
134	Phylogenetic analysis of higher taxa of Brachiopoda. <i>Lethaia</i> , 1993, 26, 1-5.	0.6	75
135	Phylogenetic analysis and classification of the Brachiopoda - reply and comments. <i>Lethaia</i> , 1993, 26, 385-386.	0.6	8
136	Lingulate brachiopods from the Cambrian-Ordovician boundary beds in Sweden. <i>Gff</i> , 1993, 115, 215-237.	0.4	15
137	The Lower Ordovician brachiopod genus <i>Lamanskya</i> and the Family Elkaniidae. <i>Transactions of the Royal Society of Edinburgh: Earth Sciences</i> , 1993, 84, 151-160.	1.0	6
138	Phyletic relationships within the Brachiopoda. <i>Gff</i> , 1991, 113, 84-86.	0.4	7
139	The acrotretacean brachiopod <i>Ceratreta tanneri</i> (Metzger) from the Upper Cambrian of Baltoscandia. <i>Gff</i> , 1990, 112, 249-263.	0.4	13
140	Discinacean brachiopods from the Ordovician Kullberg and Boda limestones of Dalarna, Sweden. <i>Gff</i> , 1987, 109, 317-326.	0.4	9
141	Ordovician mazuelloids and other microfossils from Västergötland. <i>Gff</i> , 1987, 109, 67-71.	0.4	9
142	Inarticulate brachiopods around the Middle-Upper Ordovician boundary in Västergötland. <i>Gff</i> , 1986, 108, 97-126.	0.4	23
143	Lower Viruan discontinuity surfaces in central Sweden. <i>Gff</i> , 1983, 105, 29-42.	0.4	16
144	Camenellan tommotiids from the Cambrian Series 2 of East Antarctica: biostratigraphy, palaeobiogeography, and systematics. <i>Acta Palaeontologica Polonica</i> , 0, 66, .	0.4	3

#	ARTICLE	IF	CITATIONS
145	The oldest brachiopods from the lower Cambrian of South Australia. Acta Palaeontologica Polonica, 0, , .	0.4	3
146	Oldest mickwitziid brachiopod from the Terreneuvian of southern France. Acta Palaeontologica Polonica, 0, , .	0.4	0
147	The new stem-group brachiopod Oymurania from the lower Cambrian of Siberia. Acta Palaeontologica Polonica, 0, , .	0.4	1
148	Cambrian (Stage 4 to Wuliuan) brachiopods from Sonora, Mexico. Journal of Paleontology, 0, , 1-21.	0.5	2